



1.3 PRODUCT INFORMATION

1.3.1 Summary of Product Characteristics

1.	NAME OF THE MEDICINAL PRODUCT Losartan Potassium Tablets USP 25 mg
1.1	International Non-Proprietary Name (INN) Losartan Potassium Tablets USP 25 mg
1.2	Strength 25 mg
1.3	Pharmaceutical form Film-Coated Tablets
2.	QUALITATIVE AND QUANTITATIVE COMPOSITION Each film coated tablet contains: Losartan Potassium USP 25 mg Excipients q.s. Colour: Approved colours used
3.	PHARMACEUTICAL FORM Brick red coloured, round biconvex, film coated tablets plain on both sides.
4.	CLINICAL PARTICULARS
4.1	Therapeutic indications <ul style="list-style-type: none">• Treatment of essential hypertension in adults and in children and adolescents 6-18 years of age.• Treatment of renal disease in adult patients with hypertension and type 2 diabetes mellitus with proteinuria ≥ 0.5 g/day as part of an antihypertensive treatment.• Treatment of chronic heart failure in adult patients when treatment with Angiotensin converting enzyme (ACE) inhibitors is not considered suitable due to incompatibility, especially cough, or contraindication. Patients with heart failure who have been stabilised with an ACE inhibitor should not be switched to losartan. The patients should have a left ventricular ejection fraction $\leq 40\%$ and should be clinically stable and on an established treatment regimen for chronic heart failure.• Reduction in the risk of stroke in adult hypertensive patients with left ventricular hypertrophy documented by ECG.
4.2	Posology and method of administration



Posology

Hypertension

The usual starting and maintenance dose is 50mg once daily for most patients. The maximal antihypertensive effect is attained 3-6 weeks after initiation of therapy. Some patients may receive an additional benefit by increasing the dose to 100mg once daily (in the morning). Losartan potassium film-coated tablets may be administered with other antihypertensive agents, especially with diuretics (e.g. hydrochlorothiazide).

Paediatric population

There are limited data on the efficacy and safety of losartan in children and adolescents aged 6-16 years old for the treatment of hypertension. Limited pharmacokinetic data are available in hypertensive children above one month of age. For patients who can swallow tablets, the recommended dose is 25mg once daily in patients >20 to <50kg. In exceptional cases the dose can be increased to a maximum of 50mg once daily. Dosage should be adjusted according to blood pressure response.

In patients >50kg, the usual dose is 50mg once daily. In exceptional cases the dose can be adjusted to a maximum of 100mg once daily. Doses above 1.4mg/kg (or in excess of 100mg) daily have not been studied in pediatric patients. Losartan is not recommended for use in children under 6 years old, as limited data are available in these patient groups. It is not recommended in children with glomerular filtration rate <30ml/min/1.73m², as no data are available.

Losartan is also not recommended in children with hepatic impairment.

Hypertensive type II diabetic patients with proteinuria \geq 0.5 g/day

The usual starting dose is 50 mg once daily. The dose may be increased to 100 mg once daily based on blood pressure response from one month after initiation of therapy onwards. Losartan Actavis may be administered with other antihypertensive agents (e.g. diuretics, calcium channel blockers, alpha- or beta-blockers, and centrally acting agents) as well as with insulin and other commonly used hypoglycemic agents (e.g. sulfonylureas, glitazones and glucosidase inhibitors).

Reduction in the risk of stroke in hypertensive patients with left ventricular hypertrophy documented by ECG. The usual starting dose is 50 mg of Losartan Actavis once daily. A low dose of hydrochlorothiazide should be added and/ or the dose of Losartan Actavis should be increased to 100 mg once daily based on blood pressure response.



	<p>Use in patients with intravascular volume depletion:</p> <p>For patients with intravascular volume-depletion (e.g. those treated with high-dose diuretics), a starting dose of 25mg once daily should be considered (see section 4.4).</p> <p>Use in patients with renal impairment and haemodialysis patients:</p> <p>No initial dosage adjustment is necessary in patients with renal impairment and in haemodialysis patients.</p> <p>Use in patients with hepatic impairment:</p> <p>A lower dose should be considered for patients with a history of hepatic impairment. There is no therapeutic experience in patients with severe hepatic impairment. Therefore, losartan is contraindicated in patients with severe hepatic impairment (see sections 4.3 and 4.4).</p> <p>Use in Elderly</p> <p>Although consideration should be given to initiating therapy with 25mg in patients over 75 years of age, dosage adjustment is not usually necessary for the elderly.</p> <p><u>Method of administration</u></p> <p>Losartan potassium film-coated Tablets should be swallowed with a glass of water.</p> <p>Losartan potassium film-coated Tablets may be administered with or without food.</p>
4.3	<p>Contraindications</p> <ul style="list-style-type: none">- Hypersensitivity to the active substance or to any of the excipients.- 2nd and 3rd trimester of pregnancy.- Severe hepatic impairment.- The concomitant use of Losartan Potassium with aliskiren-containing products is contraindicated in patients with diabetes mellitus or renal impairment (GFR < 60 ml/min/1.73 m²).
4.4	<p>Special warnings and precautions for use</p> <p><u>Hypersensitivity</u></p> <p><i>Angiooedema.</i> Patients with a history of angiooedema (swelling of the face, lips, throat, and/ or tongue) should be closely monitored.</p> <p><u>Hypotension and Electrolyte/Fluid Imbalance</u></p> <p>Symptomatic hypotension, especially after the first dose and after increasing of the dose, may</p>



occur in patients who are volume- and/or sodium-depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. These conditions should be corrected prior to administration of Losartan potassium film-coated Tablets, or a lower starting dose should be used (see section 4.2). This also applies to children 6 to 18 years of age.

Electrolyte imbalances

Electrolyte imbalances are common in patients with renal impairment, with or without diabetes, and should be addressed. In a clinical study conducted in type 2 diabetic patients with nephropathy, the incidence of hyperkalemia was higher in the group treated with Losartan potassium film-coated Tablets as compared to the placebo group (see section 4.8). Therefore, the plasma concentrations of potassium as well as creatinine clearance values should be closely monitored, especially patients with heart failure and a Creatinine Clearance between 30-50ml/min should be closely monitored.

The concomitant use of potassium sparing diuretics, potassium supplements and potassium containing salt substitutes with losartan is not recommended.

Hepatic impairment

Based on pharmacokinetic data which demonstrate significantly increased plasma concentrations of losartan in cirrhotic patients, a lower dose should be considered for patients with a history of hepatic impairment. There is no therapeutic experience with losartan in patients with severe hepatic impairment. Therefore losartan must not be administered in patients with severe hepatic impairment.

Losartan is also not recommended in children with hepatic impairment.

Renal impairment

As a consequence of inhibiting the renin-angiotensin system, changes in renal function including renal failure have been reported (in particular, in patients whose renal function is dependent on the rennin angiotensin aldosterone system such as those with severe cardiac insufficiency or pre-existing renal dysfunction). As with other medicinal product that affect the renin-angiotensin-aldosterone system, increases in blood urea and serum creatinine have also been reported in patients with bilateral renal artery stenosis or stenosis of the artery to a solitary kidney; these



changes in renal function may be reversible upon discontinuation of therapy. Losartan should be used with caution in patients with bilateral renal artery stenosis or stenosis of the artery to a solitary kidney.

Use in paediatric patients with renal impairment

Losartan is not recommended in children with glomerular filtration rate $<30\text{ml/min}/1.73\text{m}^2$ as no data are available.

Renal function should be regularly monitored during treatment with losartan as it may deteriorate.

This applies particularly when losartan is given in the presence of other conditions (fever, dehydration) likely to impair renal function.

Concomitant use of losartan and ACE-inhibitors has shown to impair renal function. Therefore, concomitant use is not recommended.

Renal transplantation

There is no experience in patients with recent kidney transplantation.

Primary hyperaldosteronism

Patients with primary aldosteronism generally will not respond to antihypertensive medicinal product acting through inhibition of the renin-angiotensin system. Therefore, the use of losartan tablets is not recommended.

Coronary heart disease and cerebrovascular disease

As with any antihypertensive agents, excessive blood pressure decrease in patients with ischaemic cardiovascular and cerebrovascular disease could result in a myocardial infarction or stroke.

Heart failure

In patients with heart failure, with or without renal impairment, there is - as with other medicinal product acting on the renin-angiotensin system - a risk of severe arterial hypotension, and (often acute) renal impairment.

There is no sufficient therapeutic experience with losartan in patients with heart failure and concomitant severe renal impairment, in patients with severe heart failure (NYHA class IV) as



well as in patients with heart failure and symptomatic life threatening cardiac arrhythmias. Therefore, losartan should be used with caution in these patient groups. The combination of losartan with a beta-blocker should be used with caution.

Aortic and mitral valve stenosis, obstructive hypertrophic cardiomyopathy

As with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Pregnancy

AIIRAs should not be initiated during pregnancy. Unless continued AIIRAs therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately, and, if appropriate, alternative therapy should be started.

Dual blockade of the renin-angiotensin-aldosterone system (RAAS)

There is evidence that the concomitant use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren increases the risk of hypotension, hyperkalaemia and decreased renal function (including acute renal failure). Dual blockade of RAAS through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is therefore not recommended.

If dual blockade therapy is considered absolutely necessary, this should only occur under specialist supervision and subject to frequent close monitoring of renal function, electrolytes and blood pressure.

ACE-inhibitors and angiotensin II receptor blockers should not be used concomitantly in patients with diabetic nephropathy.

Other warnings and precautions

As observed for angiotensin converting enzyme inhibitors, losartan and the other angiotensin antagonists are apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of higher prevalence of low-renin states in the black hypertensive population.



4.5	<p>Interaction with other medicinal products and other forms of interaction</p> <p>Other antihypertensive agents may increase the hypotensive action of losartan. Concomitant use with other substances which may induce hypotension as an adverse reaction (like tricyclic antidepressants, antipsychotics, baclofene, amifostine): may increase the risk of hypotension.</p> <p>Losartan is predominantly metabolised by cytochrome P450 (CYP) 2C9 to the active carboxylic-acid metabolite. In a clinical trial it was found that fluconazole (inhibitor of CYP2C9) decreases the exposure to the active metabolite by approximately 50%. It was found that concomitant treatment of losartan with rifampicine (inducer of metabolism enzymes) gave a 40% reduction in plasma concentration of the active metabolite. The clinical relevance of this effect is unknown. No difference in exposure was found with concomitant treatment with fluvastatin (weak inhibitor of CYP2C9).</p> <p>As with other medicinal product that block angiotensin II or its effects, concomitant use of other medicinal product which retain potassium (e.g. potassium-sparing diuretics: amiloride, triamterene, spironolactone) or may increase potassium levels (e.g. heparin), potassium supplements or salt substitutes containing potassium may lead to increases in serum potassium. Co-medication is not advisable.</p> <p>Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with ACE inhibitors. Very rare cases have also been reported with antiotensin II receptor antagonists. Co-administration of lithium and losartan should be undertaken with caution. If this combination proves essential, serum lithium level monitoring is recommended during concomitant use.</p> <p>When angiotensin II antagonists are administered simultaneously with NSAIDs (i.e. selective COX-2 inhibitors, acetylsalicylic acid at anti-inflammatory doses and non-selective NSAIDs), attenuation of the antihypertensive effect may occur. Concomitant use of angiotensin II antagonists or diuretics and NSAIDs may lead to an increased risk of worsening of renal function, including possible acute renal failure, and an increase in serum potassium, especially in patients with poor pre-existing renal function. The combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be</p>



	<p>given to monitoring renal function after initiation of concomitant therapy, and periodically thereafter.</p> <p>Clinical trial data has shown that dual blockade of the renin-angiotensin-aldosterone-system (RAAS) through the combined use of ACE-inhibitors, angiotensin II receptor blockers or aliskiren is associated with a higher frequency of adverse events such as hypotension, hyperkalaemia and decreased renal function (including acute renal failure) compared to the use of a single RAAS-acting agent</p>
4.6	<p>Pregnancy and lactation</p> <p><u>Pregnancy</u></p> <p>Epidemiological evidence regarding the risk of teratogenicity following exposure to ACE inhibitors during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Whilst there is no controlled epidemiological data on the risk with Angiotensin II Receptor Inhibitors (AIIRAs), similar risks may exist for this class of drugs. Unless continued AIIRA therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with AIIRAs should be stopped immediately and, if appropriate, alternative therapy should be started.</p> <p>Exposure to AIIRAs therapy during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia)</p> <p>Should exposure to AIIRAs have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.</p> <p>Infants whose mothers have taken AIIRAs should be closely observed for hypotension.</p> <p><u>Lactation</u></p> <p>Because no information is available regarding the use of losartan during breastfeeding, losartan is not recommended and alternative treatments with better established safety profiles during breast-feeding are preferable, especially while nursing a newborn or preterm infant.</p>



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

<p>4.7</p>	<p>Effects on ability to drive and use machines</p> <p>No studies on the effects on the ability to drive and use machines have been performed. However, when driving vehicles or operating machinery it must be borne in mind that dizziness or drowsiness may occasionally occur when taking antihypertensive therapy, in particular during initiation of treatment or when the dose is increased.</p>												
<p>4.8</p>	<p>Undesirable effects</p> <p>Losartan has been evaluated in clinical studies as follows:</p> <ul style="list-style-type: none"> - in a controlled clinical trials in > 3000 adult patients 18 years of age and older for essential hypertension, - in a controlled clinical trial in 177 hypertensive paediatric patients 6 to 16 years of age - in a controlled clinical trial in > 9000 hypertensive patients 55 to 80 years of age with left ventricular hypertrophy - in a controlled clinical trial in > 7700 adult patients with chronic heart failure- <p>- in a controlled clinical trial in > 1500 type 2 diabetic patients 31 years of age and older with proteinuria</p> <p>In these clinical trials, the most common adverse reaction was dizziness.</p> <p>The frequency of adverse events listed below is defined using the following convention:</p> <p>Very common ($\geq 1/10$); common ($\geq 1/100$ to, $< 1/10$); uncommon ($\geq 1/1,000$ to, $< 1/100$); rare ($\geq 1/10,000$ to, $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data).</p> <p>Table 1. The frequency of adverse reactions identified from placebo-controlled clinical studies and post marketing experience</p> <table border="1" data-bbox="315 1591 1430 1877"> <thead> <tr> <th data-bbox="315 1591 604 1654">Adverse reaction</th> <th colspan="4" data-bbox="604 1591 1276 1654">Frequency of adverse reaction by indication</th> <th data-bbox="1276 1591 1430 1654">Other</th> </tr> </thead> <tbody> <tr> <td data-bbox="315 1654 604 1877"></td> <td data-bbox="604 1654 760 1877">Hypertension</td> <td data-bbox="760 1654 906 1877">Hypertensive patients with left-</td> <td data-bbox="906 1654 1081 1877">Chronic Heart Failure</td> <td data-bbox="1081 1654 1276 1877">Hypertension and type 2 diabetes with renal disease</td> <td data-bbox="1276 1654 1430 1877">Post-marketing experience</td> </tr> </tbody> </table>	Adverse reaction	Frequency of adverse reaction by indication				Other		Hypertension	Hypertensive patients with left-	Chronic Heart Failure	Hypertension and type 2 diabetes with renal disease	Post-marketing experience
Adverse reaction	Frequency of adverse reaction by indication				Other								
	Hypertension	Hypertensive patients with left-	Chronic Heart Failure	Hypertension and type 2 diabetes with renal disease	Post-marketing experience								

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

			ventricular hypertrophy			
Blood and lymphatic system disorders						
anaemia				common		frequency not known
thrombocytopenia						frequency not known
Immune system disorders						
Hypersensitivity reactions, anaphylactic reactions, angioedema*, and vasculitis**						rare
Psychiatric disorders						
depression						frequency not known
Nervous system disorders						
dizziness	common	common	common	common		
somnolence	uncommon					
headache	uncommon		uncommon			
sleep disorders	uncommon					
paraesthesia			rare			
migraine						frequency not known

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

dysgeusia					frequency not known
Ear and labyrinth disorder					
vertigo	common	common			
tinnitus					frequency not known
Cardiac disorders					
palpitations	uncommon				
angina pectoris	uncommon				
syncope			rare		
atrial fibrillation			rare		
cerebrovascular accident			rare		
Vascular disorders					
(orthostatic) hypotension (including dose- related orthostatic effects)	uncommon		common	common	
Respiratory, thoracic and mediastinal disorders					
dyspnoea			uncommon		
cough			uncommon		frequency not known
Gastrointestinal disorders					
abdominal pain	uncommon				
obstipation	uncommon				
diarrhoea			uncommon		frequency

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

					not known
nausea			uncommon		
vomiting			uncommon		
Hepatobiliary disorders					
pancreatitis					frequency not known
hepatitis					rare
liver function abnormalities					frequency not known
Skin and subcutaneous tissue disorders					
urticaria			uncommon		frequency not known
pruritus			uncommon		frequency not known
rash	uncommon		uncommon		frequency not known
photosensitivity					frequency not known
Musculoskeletal and connective tissue disorders					
myalgia					frequency not known
arthralgia					frequency not known
rhabdomyolysis					frequency not known
Renal and urinary disorders					
renal impairment			common		

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

renal failure			common		
Reproductive system and breast disorders					
erectile dysfunction / impotence					frequency not known
General disorders and administration site conditions					
asthenia	uncommon	common	uncommon	common	
fatigue	uncommon	common	uncommon	common	
oedema	uncommon				
malaise					frequency not known
Investigations					
hyperkalaemia	common		uncommon [†]	common [‡]	
increased alanine aminotransferase (ALT) [§]	rare				
increase in blood urea, serum creatinine, and serum potassium			common		
hyponatraemia					frequency not known
hypoglycaemia				common	
<p>*Including swelling of the larynx, glottis, face, lips, pharynx, and/or tongue (causing airway obstruction); in some of these patients angioedema had been reported in the past in connection with the administration of other medicines, including ACE inhibitors</p> <p>**Including Henoch-Schönlein purpura</p> <p> Especially in patients with intravascular depletion, e.g. patients with severe heart failure or under</p>					

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



	<p>treatment with high dose diuretics</p> <p>† Common in patients who received 150 mg losartan instead of 50 mg</p> <p>‡ In a clinical study conducted in type 2 diabetic patients with nephropathy, 9.9% of patients treated with Losartan tablets developed hyperkalaemia >5.5 mmol/l and 3.4% of patients treated with placebo</p> <p>§ Usually resolved upon discontinuation</p> <p>The following additional adverse reactions occurred more frequently in patients who received losartan than placebo (frequencies not known): <u>back pain, urinary tract infection, and flu-like symptoms</u></p> <p>Renal and urinary disorders:</p> <p>As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function including renal failure have been reported in patients at risk; these changes in renal function may be reversible upon discontinuation of therapy (see section 4.4)</p> <p>Paediatric populations:</p> <p>The adverse reaction profile for paediatric patients appears to be similar to that seen in adult patients. Data in the paediatric population are limited.</p>
4.9	<p>Overdose</p> <p>Symptoms of intoxication</p> <p>Limited data are available with regard to overdose in humans. The most likely manifestations of overdose would be hypotension and tachycardia. Bradycardia could occur from parasympathetic (vagal) stimulation.</p> <p>Treatment of intoxication</p> <p>If symptomatic hypotension should occur, supportive treatment should be instituted.</p> <p>Measures are depending on the time of medicinal product intake and kind and severity of symptoms. Stabilisation of the circulatory system should be given priority. After oral intake the</p>



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

	<p>administration of a sufficient dose of activated charcoal is indicated. Afterwards, close monitoring of the vital parameters should be performed. Vital parameters should be corrected if necessary.</p> <p>Neither Losartan nor the active metabolite can be removed by haemodialysis.</p>
5.	PHARMACOLOGICAL PROPERTIES
5.1	<p>Pharmacodynamic properties</p> <p>Pharmacotherapeutic group: Angiotensin II Receptor Antagonists, ATC code: C09CA01</p> <p>Losartan is a synthetic oral angiotensin-II receptor (type AT₁) antagonist. Angiotensin II, a potent vasoconstrictor, is the primary active hormone of the renin/angiotensin system and an important determinant of the pathophysiology of hypertension. Angiotensin II binds to the AT₁ receptor found in many tissues (e.g. vascular smooth muscle, adrenal gland, kidneys and the heart) and elicits several important biological actions, including vasoconstriction and the release of aldosterone. Angiotensin II also stimulates smooth muscle cell proliferation.</p> <p>Losartan selectively blocks the AT₁ receptor. <i>In vitro</i> and <i>in vivo</i> losartan and its pharmacologically active carboxylic acid metabolite E-3174 block all physiologically relevant actions of angiotensin II, regardless of the source or route of its synthesis.</p> <p>Losartan does not have an agonist effect nor does it block other hormone receptors or ion channels important in cardiovascular regulation. Furthermore Losartan does not inhibit ACE (kininase II), the enzyme that degrades bradykinin. Consequently, there is no potentiation of undesirable bradykinin-mediated effects.</p> <p>During administration of Losartan, removal of the angiotensin II negative feedback on renin secretion leads to increased plasma renin activity (PRA). Increase in the PRA leads to an increase in angiotensin II in plasma. Despite these increases, antihypertensive activity and suppression of plasma aldosterone concentration are maintained, indicating effective angiotensin II receptor blockade. After discontinuation of Losartan, PRA and angiotensin II values fell within three days to the baseline values.</p> <p>Both Losartan and its principal active metabolite have a far greater affinity for the AT₁-receptor than for the AT₂-receptor. The active metabolite is 10- to 40- times more active than Losartan on a weight for weight basis.</p> <p><u>Hypertension Studies</u></p> <p>In controlled clinical studies, once-daily administration of Losartan to patients with mild to</p>

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



moderate essential hypertension produced statistically significant reductions in systolic and diastolic blood pressure. Measurements of blood pressure 24 hours post-dose relative to 5 – 6 hours post-dose demonstrated blood pressure reduction over 24 hours; the natural diurnal rhythm was retained. Blood pressure reduction at the end of the dosing interval was 70 – 80% of the effect seen 5-6 hours postdose.

Discontinuation of Losartan in hypertensive patients did not result in an abrupt rise in blood pressure (rebound). Despite the marked decrease in blood pressure, Losartan had no clinically significant effects on heart rate.

Losartan is equally effective in males and females, and in younger (below the age of 65 years) and older hypertensive patients.

LIFE-Study

The Losartan Intervention For Endpoint Reduction in Hypertension [LIFE] study was a randomised, triple-blind, active-controlled study in 9193 hypertensive patients aged 55 to 80 years with ECGdocumented left-ventricular hypertrophy. Patients were randomised to once daily Losartan 50mg or once daily atenolol 50mg. If goal blood pressure (< 140/90mmHg) was not reached, hydrochlorothiazide (12.5mg) was added first and, if needed, the dose of Losartan or atenolol was then increased to 100mg once daily. Other antihypertensives, with the exception of ACE-inhibitors, angiotensin II antagonists or beta-blockers were added if necessary to reach the goal blood pressure.

The mean length of follow up was 4.8 years.

The primary endpoint was the composite of cardiovascular morbidity and mortality as measured by a reduction in the combined incidence of cardiovascular death, stroke and myocardial infarction. Blood pressure was significantly lowered to similar levels in the two groups. Treatment with losartan resulted in a 13.0% risk reduction (p=0.021, 95% confidence interval 0.77-0.98) compared with atenolol for patients reaching the primary composite endpoint. This was mainly attributable to a reduction of the incidence of stroke. Treatment with losartan reduced the risk of stroke by 25% relative to atenolol (p=0.001 95% confidence interval 0.63-0.89). The rates of cardiovascular death and myocardial infarction were not significantly different between the treatment groups.

Race



In the LIFE-Study black patients treated with Losartan had a higher risk of suffering the primary combined endpoint, i.e. a cardiovascular event (e.g. cardiac infarction, cardiovascular death) and especially stroke, than the black patients treated with atenolol. Therefore the results observed with losartan in comparison with atenolol in the LIFE study with regard to cardiovascular morbidity/mortality do not apply for black patients with hypertension and left ventricular hypertrophy.

RENAAL-Study

The Reduction of Endpoints in NIDDM with the Angiotensin II Receptor Antagonist Losartan RENAAL study was a controlled clinical study conducted worldwide in 1513 Type 2 diabetic patients with proteinuria, with or without hypertension. 751 Patients were treated with Losartan. The objective of the study was to demonstrate a nephroprotective effect of Losartan potassium over and above the benefit of lowering blood pressure.

Patients with proteinuria and a serum creatinine of 1.3 – 3.0mg/dl were randomised to receive Losartan 50mg once a day, titrated if necessary, to achieve blood pressure response, or to placebo, on a background of conventional antihypertensive therapy excluding ACE-inhibitors and angiotension II antagonists.

Investigators were instructed to titrate the study medication to 100mg daily as appropriate; 72 % of patients were taking the 100mg daily dose for the majority of the time. Other antihypertensive agents (diuretics, calcium antagonists, alpha- and beta-receptor blockers and also centrally acting antihypertensives) were permitted as supplementary treatment depending on the requirement in both groups. Patients were followed up for up to 4.6 years (3.4 years on average).

The primary endpoint of the study was a composite endpoint of doubling of the serum creatinine endstage renal failure (need for dialysis or transplantation) or death.

The results showed that the treatment with Losartan (327 events) as compared with placebo (359 events) resulted in a 16.1 % risk reduction ($p = 0.022$) in the number of patients reaching the primary composite endpoint. For the following individual and combined components of the primary endpoint, the results showed a significant risk reduction in the group treated with Losartan: 25.3% risk reduction for doubling of the serum creatinine ($p = 0.006$); 28.6 % risk reduction for end-stage renal failure ($p = 0.002$); 19.9% risk reduction for end-stage renal failure or death ($p = 0.009$); 21.0% risk reduction for doubling of serum creatinine or end-stage renal



failure ($p = 0.01$).

All-cause mortality rate was not significantly different between the two treatment groups.

In this study losartan was generally well tolerated, as shown by a therapy discontinuation rate on account of adverse events that was comparable to the placebo group.

HEAAL Study

The Heart Failure Endpoint Evaluation of Angiotensin II Antagonist Losartan (HEAAL) study was a controlled clinical study conducted worldwide in 3834 patients aged 18 to 98 years with heart failure (NYHA Class II-IV) who were intolerant of ACE inhibitor treatment. Patients were randomised to receive losartan 50 mg once a day or losartan 150 mg, on a background of conventional therapy excluding ACE-inhibitors.

Patients were followed for over 4 years (median 4.7 years). The primary endpoint of the study was a composite endpoint of all cause death or hospitalization for heart failure.

The results showed that treatment with 150 mg losartan (828 events) as compared with 50 mg losartan (889 events) resulted in a 10.1% risk reduction ($p=0.027$, 95% confidence interval 0.82-0.99) in the number of patients reaching the primary composite endpoint. This was mainly attributable to a reduction of the incidence of hospitalization for heart failure. Treatment with 150 mg losartan reduced the risk of hospitalization for heart failure by 13.5% relative to 50 mg losartan ($p=0.025$, 95% confidence interval 0.76-0.98). The rate of all cause death was not significantly different between the treatment groups. Renal impairment, hypotension, and hyperkalaemia were more common in the 150 mg group than in the 50 mg group, but these adverse events did not lead to significantly more treatment discontinuations in the 150 mg group.

ELITE I and ELITE II Study

In the ELITE Study carried out over 48 weeks in 722 patients with heart failure (NYHA Class II-IV), no difference was observed between the patients treated with Losartan and those treated with captopril was observed with regard to the primary endpoint of a long-term change in renal function. The observation of the ELITE I Study, that, compared with captopril, Losartan reduced the mortality risk, was not confirmed in the subsequent ELITE II Study, which is described in the following.

In the ELITE II Study Losartan 50mg once daily (starting dose 12.5mg, increased to 25mg, then 50mg once daily) was compared with captopril 50mg three times daily (starting dose 12.5mg,



increased to 25mg and then to 50mg three times daily). The primary endpoint of this prospective study was the all-cause mortality.

In this study 3152 patients with heart failure (NYHA Class II-IV) were followed for almost two years (median: 1.5 years) in order to determine whether Losartan is superior to captopril in reducing all cause mortality. The primary endpoint did not show any statistically significant difference between Losartan and captopril in reducing all-cause mortality.

In both comparator-controlled (not placebo-controlled) clinical studies on patients with heart failure the tolerability of Losartan was superior to that of captopril, measured on the basis of a significantly lower rate of discontinuations of therapy on account of adverse events and a significantly lower frequency of cough.

An increased mortality was observed in ELITE II in the small subgroup (22% of all HF patients) taking beta-blockers at baseline.

ONTARGET, VA NEPHRON-D and ALTITUDE study

Two large randomised, controlled trials (ONTARGET (ONgoing Telmisartan Alone and in combination with Ramipril Global Endpoint Trial) and VA NEPHRON-D (The Veterans Affairs Nephropathy in Diabetes)) have examined the use of the combination of an ACE-inhibitor with an angiotensin II receptor blocker.

ONTARGET was a study conducted in patients with a history of cardiovascular or cerebrovascular disease, or type 2 diabetes mellitus accompanied by evidence of end-organ damage. VA NEPHRON-D was a study in patients with type 2 diabetes mellitus and diabetic nephropathy.

These studies have shown no significant beneficial effect on renal and/or cardiovascular outcomes and mortality, while an increased risk of hyperkalaemia, acute kidney injury and/or hypotension as compared to monotherapy was observed. Given their similar pharmacodynamic properties, these results are also relevant for other ACE-inhibitors and angiotensin II receptor blockers.

ACE-inhibitors and angiotensin II receptor blockers should therefore not be used concomitantly in patients with diabetic nephropathy.

ALTITUDE (Aliskiren Trial in Type 2 Diabetes Using Cardiovascular and Renal Disease Endpoints) was a study designed to test the benefit of adding aliskiren to a standard therapy of an ACE-inhibitor or an angiotensin II receptor blocker in patients with type 2 diabetes mellitus and



chronic kidney disease, cardiovascular disease, or both. The study was terminated early because of an increased risk of adverse outcomes. Cardiovascular death and stroke were both numerically more frequent in the aliskiren group than in the placebo group and adverse events and serious adverse events of interest (hyperkalaemia, hypotension and renal dysfunction) were more frequently reported in the aliskiren group than in the placebo group.

Paediatric population

The antihypertensive effect of Losartan potassium film-coated Tablets was established in a clinical study involving 177 hypertensive pediatric patients 6 to 16 years of age with a body weight > 20kg and a glomerular filtration rate >30ml/min/1.73m². Patients who weighted >20kg to <50kg received either 2.5, 25 or 50mg of losartan daily and patients who weighted >50kg received either 5, 50 or 100mg of losartan daily. At the end of three weeks, losartan administration once daily lowered trough blood pressure in a dose-dependent manner.

Overall, there was a dose-response. The dose-response relationship became very obvious in the low dose group compared to the middle dose group (period I: -6.2 mmHg vs. -11.65mmHg), but was attenuated when comparing the middle dose group with the high dose group (period I: -11.65mmHg vs. -12.21mmHg). The lowest doses studied, 2.5mg and 5mg, corresponding to an average daily dose of 0.07mg/ kg, did not appear to offer consistent antihypertensive efficacy.

These results were confirmed during period II of the study where patients were randomized to continue losartan or placebo, after three weeks of treatment. The difference in blood pressure increase as compared to placebo was largest in the middle dose group (6.70mm Hg middle dose vs. 5.38mmHg high dose). The rise in trough diastolic blood pressure was the same in patients receiving placebo and in those continuing losartan at the lowest dose in each group, again suggesting that the lowest dose in each group did not have significant antihypertensive effect.

Long-term effects of losartan on growth, puberty and general development have not been studied. The long-term efficacy of antihypertensive therapy with losartan in childhood to reduce cardiovascular morbidity and mortality has also not been established.

In hypertensive (N=60) and normotensive (N=246) children with proteinuria, the effect of losartan on proteinuria was evaluated in a 12-week placebo- and active-controlled (amlodipine) clinical study. Proteinuria was defined as urinary protein/creatinine ratio of ≥ 0.3 . The hypertensive patients (ages 6 through 18 years) were randomized to receive either losartan (n=30)



or amlodipine (n=30). The normotensive patients (ages 1 through 18 years) were randomized to receive either losartan (n=122) or placebo (n=124). Losartan was given at doses of 0.7 mg/kg to 1.4 mg/kg (up to maximum dose of 100 mg per day). Amlodipine was given at doses of 0.05 mg/kg to 0.2 mg/kg (up to a maximum dose of 5 mg per day).

Overall, after 12 weeks of treatment, patients receiving losartan experienced a statistically significant reduction from baseline in proteinuria of 36% versus 1% increase in placebo/amlodipine group ($p \leq 0.001$). Hypertensive patients receiving losartan experienced a reduction from baseline proteinuria of -41.5% (95% CI -29.9;-51.1) versus +2.4% (95% CI -22.2; 14.1) in the amlodipine group. The decline in both systolic blood pressure and diastolic blood pressure was greater in the losartan group (-5.5/-3.8 mmHg) versus the amlodipine group (-0.1/+0.8 mmHg). In normotensive children a small decrease in blood pressure was observed in the losartan group (-3.7/-3.4 mmHg) compared to placebo. No significant correlation between the decline in proteinuria and blood pressure was noted, however it is possible that the decline in blood pressure was responsible, in part, for the decline in proteinuria in the losartan treated group. Long-term effects of losartan in children with proteinuria were studied for up to 3 years in the open-label safety extension phase of the same study, in which all patients completing the 12-week base study were invited to participate. A total of 268 patients entered the open-label extension phase and were re-randomized to losartan (N=134) or enalapril (N=134) and 109 patients had ≥ 3 years of follow-up (pre-specified termination point of ≥ 100 patients completing 3 years of follow-up in the extension period). The dose ranges of losartan and enalapril, given according to investigator discretion, were 0.30 to 4.42 mg/kg/day and 0.02 to 1.13 mg/kg/day, respectively. The maximum daily doses of 50 mg for < 50 kg body weight and 100 mg > 50 kg were not exceeded for most patients during the extension phase of the study.

In summary, the results of the safety extension show that losartan was well-tolerated and led to sustained decreases in proteinuria with no appreciable change in glomerular filtration rate (GFR) over 3 years. For normotensive patients (n=205), enalapril had a numerically greater effect compared to losartan on proteinuria (-33.0% (95%CI -47.2;-15.0) vs -16.6% (95%CI -34.9; 6.8)) and on GFR (9.4(95%CI 0.4; 18.4) vs -4.0(95%CI -13.1; 5.0) ml/min/1.73m²). For hypertensive patients (n=49), losartan had a numerically greater effect on proteinuria (-44.5% (95%CI -64.8; -12.4) vs -39.5% (95%CI -62.5; -2.2)) and GFR (18.9(95%CI 5.2; 32.5) vs -13.4(95%CI -27.3;

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



	0.6)) ml/min/1.73m ² .
5.2	<p>Pharmacokinetic properties</p> <p><u>Absorption</u></p> <p>Following oral administration, losartan is well absorbed and undergoes first-pass metabolism, forming an active carboxylic acid metabolite and other inactive metabolites. The systemic bioavailability of losartan tablets is approximately 33%. Mean peak concentrations of losartan and its active metabolite are reached in 1 hour and in 3-4 hours, respectively.</p> <p><u>Distribution</u></p> <p>Both losartan and its active metabolite are $\geq 99\%$ bound to plasma proteins, primarily albumin. The volume of distribution of losartan is 34 litres.</p> <p><u>Biotransformation</u></p> <p>About 14% of an intravenously- or orally-administered dose of losartan is converted to its active metabolite. Following oral and intravenous administration of ¹⁴C-labeled losartan potassium, circulating plasma radioactivity primarily is attributed to losartan and its active metabolite. Minimal conversion of losartan to its active metabolite was seen in about one percent of individuals studied.</p> <p>In addition to the active metabolite, inactive metabolites are formed.</p> <p><u>Elimination</u></p> <p>Plasma clearance of losartan and its active metabolite is about 600mL/min and 50mL/min, respectively. Renal clearance of losartan and its active metabolite is about 74mL/min and 26mL/min, respectively. When losartan is administered orally, about 4% of the dose is excreted unchanged in the urine, and about 6% of the dose is excreted in the urine as active metabolite. The pharmacokinetics of losartan and its active metabolite are linear with oral losartan potassium doses up to 200mg.</p> <p>Following oral administration, plasma concentrations of losartan and its active metabolite decline polyexponentially with a terminal half-life of about 2 hours and 6-9 hours, respectively. During oncedaily dosing with 100 mg, neither losartan nor its active metabolite accumulates significantly in plasma.</p> <p>Both biliary and urinary excretion contribute to the elimination of losartan and its metabolites.</p> <p>Following an oral dose/intravenous administration of ¹⁴C-labeled losartan in man, about 35%/43% of radioactivity is recovered in the urine and 58%/50% in the faeces.</p> <p><u>Characteristics in Patients</u></p> <p>In elderly hypertensive patients the plasma concentrations of losartan and its active metabolite do not differ essentially from those found in young hypertensive patients.</p> <p>In female hypertensive patients the plasma levels of losartan were up to twice as high as in male hypertensive patients, while the plasma levels of the active metabolite did not differ between men and women.</p>

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

	<p>In patients with mild to moderate alcohol-induced hepatic cirrhosis, the plasma levels of losartan and its active metabolite after oral administration were respectively 5 and 1.7 times higher than in young male volunteers (see section 4.2 and 4.4).</p> <p>Plasma concentrations of Losartan are not altered in patients with a creatinine clearance above 10ml/minute. Compared to patients with normal renal function, the AUC for Losartan is about 2-times higher in haemodialysis dialysis patients.</p> <p>The plasma concentrations of the active metabolite are not altered in patients with renal impairment or in heamodialysis patients.</p> <p>Neither Losartan nor the active metabolite can be removed by haemodialysis.</p> <p><u>Pharmacokinetics in paediatric patients</u></p> <p>The pharmacokinetics of losartan have been investigated in 50 hypertensive paediatric patients > 1 month to < 16 years of age following once daily oral administration of approximately 0.54 to 0.77mg/kg of losartan (mean doses).</p> <p>The results showed that the active metabolite is formed from losartan in all age groups. The results showed roughly similar pharmacokinetic parameters of losartan following oral administration in infants and toddlers, preschool children, school age children and adolescents. The pharmacokinetic parameters for the metabolite differed to a greater extent between the age groups. When comparing preschool children with adolescents these differences became statistically significant. Exposure in infants/ toddlers was comparatively high.</p>
5.3	<p>Preclinical safety data</p> <p>Preclinical data reveal no special hazard for humans based on conventional studies of general pharmacology, genotoxicity and carcinogenic potential. In repeated dose toxicity studies, the administration of losartan induced a decrease in the red blood cell parameters (erythrocytes, haemoglobin, haematocrit), a rise in urea-N in the serum and occasional rises in serum creatinine, a decrease in heart weight (without a histological correlate) and gastrointestinal changes (mucous membrane lesions, ulcers, erosions, haemorrhages). Like other substances that directly affect the renin-angiotensin system, losartan has been shown to induce adverse effects on the late foetal development, resulting in foetal death and malformations.</p>
6.	<p>PHARMACEUTICAL PARTICULARS</p>

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

6.1	List of Excipients		
	S.N	Ingredients	Specs.
	1	Microcrystalline Cellulose	USP
	2	Croscarmellose Sodium	USP
	3	Sodium Starch Glycol	USP
	4	Talc	USP
	5	Colloidal Silicon Dioxide	USP
	6	Magnesium Stearate	USP
7	AF Coat Non Colouring Agent Aqueous Extra	-----	
6.2	Incompatibilities Not applicable		
6.3	Shelf life 36 months		
6.4	Special precautions for storage Store at a temperature 30 ⁰ C in a dry place. Protect from light.		
6.5	Nature and contents of container 14x2x10 Packed in a carton along with Pack insert.		
6.6	Special precautions for Disposal/ handling This medicinal product does not require any special storage conditions.		
7.	REGISTRANT FIRST J POCCO PHARM LTD. 89 DENIS OSADEBE WAY ASABA DELTA STATE NIG		
8.	MANUFACTURER Scott-Edil Pharmacia Ltd. 56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA		
9.	DATE OF REVISION OF THE TEXT To be given after registration approval from PPB.		

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg



SCOTT-EDIL Pharmacia Ltd.

56 EPIP, Phase-I Jharmajri, Baddi, Distt. Solan (HP) INDIA

10.	DOSIMETRY (IF APPLICABLE) Not applicable
11.	INSTRUCTIONS FOR PREPARATION OF RADIOPHARMACEUTICALS (IF APPLICABLE) Not Applicable

MODULE 1: Administrative and Prescribing Information

Dossier for Losartan Potassium Tablets USP 25 mg