

VALUMORAL TABLETS (Trypsin + Chymotrypsin)

- Prescribing Information

- Overview

Composition

Each enteric-coated tablet contains: Trypsin plus Chymotrypsin ..50,000 armour units enzymatic activity (Trypsin and Chymotrypsin are present in the ratio of 6:1 approximately)

Dosage Form

Enteric-coated tablet for oral use.

Pharmacology

The cells in the pancreas synthesize and produce digestive enzymes that breakdown fats (lipases), starches (amylases) and proteins (proteases). Pancreatic proteases can be divided into several families of enzymes that differ in structure and catalytic effect in how they interact with the peptide bonds of proteins. Trypsin and chymotrypsin are two types of proteases originally synthesized in the pancreas in the inactive form of zymogen precursors (trypsinogen and chymotrypsinogen) for the purpose of stopping unnecessary cellular activity and controlling when and where enzyme activity occurs. Zymogens are then carried either into the bloodstream or the intestines where they are excreted or are converted by the process of proteolysis into the active enzymes that aid digestion. When taking the trypsin-chymotrypsin combination, the active proteolytic enzymes are being ingested and used in addition to the inactive forms the body naturally produces. Trypsin and chymotrypsin give the body the extra boost it might need for smoother digestion of proteins as well as for reducing inflammation and fighting infection.

Fibrinolytic Activity

When fibrin clots have stopped bleeding, the body's own fibrinolytic agent, plasmin, breaks the fibrin barrier. The liver, in response to trauma, releases acute-phase reactants (APRs) that inhibit plasmin (and its fibrinolytic action). Chymotrypsin and trypsin together break down the fibrin barrier, thus improving and restoring circulation, resolving oedema, haematoma and pain, promoting phagocytosis to remove the debris, and accelerate recovery.

There are reports suggesting that the chymotrypsin-trypsin combination helps modulate the process of inflammation. Thus, the trypsin-chymotrypsin combination reduces the pro-inflammatory mediators and fastens the healing process.

Smoothens the Process of Digestion

Trypsin helps to break down large protein molecules by cutting protein chains at specific sites. The large protein molecule is actually a chain of smaller units called amino acids, which are linked, end-to-end, in chains of hundreds. There are 20 different amino acids from which these protein chains are made. The specific site along the protein chain where trypsin is active is one with the amino acids, lysine and arginine, which are two of the smaller amino acids.

The enzyme, chymotrypsin, also cuts the larger protein chain, but at sites that are different from where trypsin cuts it. Chymotrypsin makes its cut at positions along the protein chain that contain very large amino acids such as phenylalanine, tyrosine and tryptophan. Otherwise, it is very similar to trypsin.

In some individuals, the production of these digestive enzymes is deficient, resulting in the inability to completely digest food. This can result in abdominal pain, indigestion, gas and malnutrition. This condition is treatable with trypsin-chymotrypsin enzyme supplements.

Indications

Trypsin-chymotrypsin tablets are indicated in post-operative wounds, oedema and haematoma, prevention of inflammation of the surgical stitches, pelvic inflammatory disease, caesarean section, episiotomy, abdominal hysterectomy, tooth extraction, peri-apical abscess, maxillofacial surgery, post-traumatic oedema, soft tissue injury, fractures and dislocation, sports injuries, and sprains and strains.

Treatment should begin as soon as possible after injury occurs.

Dosage And Administration

One tablet, four times a day, half an hour before meals or as directed by the physician. The tablet must be swallowed whole to preserve the enteric coating.

Contraindications

are contraindicated in patients with severe liver problems, kidney impairment, peptic ulcer, high vitreous pressure, and hypersensitivity.

- Hypersensitivity to the ingredients of VALUMORAL Tablets.
- VALUMORAL Tablets

Warnings and Precautions

General

- Rarely, chymotrypsin might cause an allergic reaction when taken by mouth. Symptoms include itching, shortness of breath, swelling of the lips or throat, shock, loss of consciousness, and death.
- Not to be used in patients with severe hepatic impairment or renal damage or irregularities of the blood clotting mechanism.

Drug Interactions

Herbal Supplements/Alcohol

Systemic proteases may increase the effectiveness of herbal supplements. Chymotrypsin is also known to interact with alcohol.

Antibiotics

Administration of the trypsin-chymotrypsin combination (intramuscularly) has been found to increase the levels of orally administered semi-synthetic penicillin antibiotics in the blood serum and organs of rats.

Chymotrypsin is known to interact with chloramphenicol.

Anticoagulants

The trypsin-chymotrypsin combination should not be administered concurrently with anticoagulants such as coumadin, heparin and clopidogrel.

Pregnancy

Not enough is known about the use of trypsin and chymotrypsin during pregnancy.

Lactation

Not enough is known about the use of trypsin and chymotrypsin during breastfeeding.

Undesirable Effects

Rarely, chymotrypsin might cause an allergic reaction when taken by mouth. Symptoms include itching, shortness of breath, swelling of the lips or throat, shock, loss of consciousness, and death.

Occasional gastric disturbance may also occur.

Overdosage

No data available.

Packaging Information

VALUMORAL TABLETS.....Strip pack of 10 tablets

EXPIRY :- 3 YEARS

SOLE AGENT : VALUELIFE PHARMACY CO. LTD.

NAFDAC REG. NO. : B4-0420



164(L) x 18(B) x 70(H)mm Prepared On : 17Nov2017

NAFDAC Reg. No: B4-0420

3 x 10 Tablets



فاليومورال
Valumoral
Trypsin-Chymotrypsin 50,000 units

Valumoral
Trypsin-Chymotrypsin 50,000 units

COMPOSITION : Each enteric sugar coated tablet contains :
Trypsin chymotrypsin (6:1)
equivalent to 50,000 USP
units of proteolytic enzyme activity,
Approved colour used.

التركيب : يحتوي كل قرص مغلف بالسكر على:
التريسينكيماتريسين (6:1) أي ما يعادل ٥٠,٠٠٠ وحدة حسب دستور
الأدوية الأمريكي في فعالية الأنزيم الحائل للبروتين.

Protect from light
Store in a cool, dry & dark place below 30°C.
Keep all medicines out of reach of children.
Dosage: As directed by the Physician.



التون المستحسن مستخدم.
الجرعة: حسب تعليمات الطبيب.
يحفظ في مكان بارد وجاف ومظلم.
تحفظ جميع الأدوية بعيدا عن متناول الأطفال.

NAFDAC Reg. No: B4-0420

3 x 10 Tablets



فاليومورال
Valumoral
Trypsin-Chymotrypsin 50,000 units

Mfg. Lic. No. : KD-354
Batch No. :
Mfg Date :
Exp Date :



Manufactured By:
Mancare
PHARMACEUTICALS PVT. LTD.
Plot No. 55, 55/1, 55/2, V.M.I.E, Dheroi Village,
Vasai (W), Dist. Thane, Maharashtra, India.
E-mail: mancars@ymail.com

Sole Agent:
Valulife Pharmacy Co. Ltd.
No. 46, Ifite Road, High Tension Street,
Aroma, Awka, Anambara state, Nigeria

60 x 162

40mm

Prepared On : 25Oct2013

Valumoral Trypsin Chymotrypsin Tablets
Valumoral
 Each enteric sugar coated tablet contains:
 Trypsin Chymotrypsin (6:1) equivalent to
 50,000 USP units of proteolytic enzyme activity,
 Approved colour used.
 Dosage : As directed by the physician.
 Store in a cool, dry & dark place.
 Keep all medicines out of reach of children.

Manufactured By:
Mancare
 PHARMACEUTICALS PVT. LTD.
 Plot No. 59, 60, 85, 86, V.M.I.E. Dowlai Village,
 Vasai (W), Dist. Thane, Maharashtra, India.
 E-mail: mancep@yahoo.co.in

Sole Agent:
Valulife Pharmacy Co. Ltd.
 No. 46, Ifite Road, High Tension Street,
 Arroma, Awka, Anambara state, Nigeria

Mfg. Lic. No.: KD-354
 Nafdac Reg. No: B4-0420

Valumoral Trypsin Chymotrypsin Tablets
Valumoral
 Each enteric sugar coated tablet contains:
 Trypsin Chymotrypsin (6:1) equivalent to
 50,000 USP units of proteolytic enzyme activity,
 Approved colour used.
 Dosage : As directed by the physician.
 Store in a cool, dry & dark place.
 Keep all medicines out of reach of children.

Manufactured By:
Mancare
 PHARMACEUTICALS PVT. LTD.
 Plot No. 59, 60, 85, 86, V.M.I.E. Dowlai Village,
 Vasai (W), Dist. Thane, Maharashtra, India.
 E-mail: mancep@yahoo.co.in

Sole Agent:
Valulife Pharmacy Co. Ltd.
 No. 46, Ifite Road, High Tension Street,
 Arroma, Awka, Anambara state, Nigeria

Mfg. Lic. No.: KD-354
 Nafdac Reg. No: B4-0420

Valumoral Trypsin Chymotrypsin Tablets
Valumoral
 Each enteric sugar coated tablet contains:
 Trypsin Chymotrypsin (6:1) equivalent to
 50,000 USP units of proteolytic enzyme activity,
 Approved colour used.
 Dosage : As directed by the physician.
 Store in a cool, dry & dark place.
 Keep all medicines out of reach of children.

Manufactured By:
Mancare
 PHARMACEUTICALS PVT. LTD.
 Plot No. 59, 60, 85, 86, V.M.I.E. Dowlai Village,
 Vasai (W), Dist. Thane, Maharashtra, India.
 E-mail: mancep@yahoo.co.in

Sole Agent:
Valulife Pharmacy Co. Ltd.
 No. 46, Ifite Road, High Tension Street,
 Arroma, Awka, Anambara state, Nigeria

Mfg. Lic. No.: KD-354
 Nafdac Reg. No: B4-0420

Valumoral Trypsin Chymotrypsin Tablets
Valumoral
 Each enteric sugar coated tablet contains:
 Trypsin Chymotrypsin (6:1) equivalent to
 50,000 USP units of proteolytic enzyme activity,
 Approved colour used.
 Dosage : As directed by the physician.
 Store in a cool, dry & dark place.
 Keep all medicines out of reach of children.

Manufactured By:
Mancare
 PHARMACEUTICALS PVT. LTD.
 Plot No. 59, 60, 85, 86, V.M.I.E. Dowlai Village,
 Vasai (W), Dist. Thane, Maharashtra, India.
 E-mail: mancep@yahoo.co.in

Sole Agent:
Valulife Pharmacy Co. Ltd.
 No. 46, Ifite Road, High Tension Street,
 Arroma, Awka, Anambara state, Nigeria

Mfg. Lic. No.: KD-354
 Nafdac Reg. No: B4-0420

For the use of Registered medical Practitioner or a Hospital or a Laboratory only.



Valumoral

Trypsin-Chymotrypsin 50,000 units

COMPOSITION

Each enteric sugar coated tablet contains:

Trypsin Chymotrypsin (6:1) Equivalent to 50,000 U.S.P units of proteolytic enzyme activity

DESCRIPTION

A proteolytic enzyme obtained by the activation of Chymotrypsinogen extracted from OX pancreas. A non steroidal anti-inflammatory(NSAID) and rheumatoid drug. Activity is expressed in terms of USP units.

USES AND ADMINISTRATION

Trypsin- Chymotrypsin is given with the intention of reducing soft tissue inflammation and oedema from abscesses and ulcer or associated with traumatic injuries and to promote liquefaction of secretions of the upper respiratory tract in patients suffering from Asthama, Bronchitis,pulmonary diseases and sinusitis.Chymotrypsin is administered by mouth in conjunction with trypsin in a combined dose of 100,000 units(USP) four times a day.

ADVERSE EFFECTS

Chymotrypsin is antigenic and severe allergic reactions have occasionally followed by the use.

PRESENTATION

3 strips of 10 tablets in a Carton & 3 Alu Alu blisters of 10 Tablets in a Carton.

STORAGE INSTRUCTIONS

Store in a cool, dry place protected from light.

KEEP ALL MEDICINE OUT OF REACH OF CHILDREN.

Mfg. Lic. No.: KD-354

Nafdac Reg. No: B4-0420



Manufactured By :

Mancare

PHARMACEUTICALS PVT. LTD.

Plot No. 59, 60, 85, 86, V.M.I.E, Dowali Village,

Vasai (W), Dist. Thane, Maharashtra, India.

E-mail: mancerp@yahoo.co.in



Sole Agent:

Valulife Pharmacy Co. Ltd.

No. 46, Ifite Road, High Tension Street,

Arroma, Awka, Anambara state,

Nigeria