

## **1. NAME OF THE MEDICINAL PRODUCT**

Haepler Blood capsules  
(Iron + Vitamins)

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each Capsule of Haepler Blood capsules contains

Vitamin B1(Thiamine Mononitrate) 1.5mg

Vitamin B2 (Riboflavin) 2mg

Vitamin B6 (Pyridoxine HCl) 2mg

Vitamin C (Sodium Ascorbate) 75mg

Vitamin B12 (Cyanocobalamin) 3mcg

Nicotinamide 20mg

Folic Acid 0.4mg

Iron (from Ferrous Fumarate) 15mg

Excipients:

For full list of excipients, see section 6.1

## **3. PHARMACEUTICAL FORM**

Oral Capsule

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Haepler Blood Capsules is a specially formulated supplement in the management of Iron deficiency anemia, prevention and treatment of malnutrition and other vitamins deficiencies. It provides vitamins and iron for maintenance of good health and vitality in adults and pregnant women..

### **4.2 Posology and Method of Administration**

#### Posology

One capsule daily for 30days or as directed by the physician.

Mode of Administration(s): Oral

### **4.3 Contraindications**

- Primary (idiopathic) or secondary iron storage disease.
- Anaemia associated with ineffective erythropoiesis, marrow hypoplasia, sideroblastic change, uncomplicated Cyanocobalamin or folate deficiency.

- Intestinal disease (oral iron may aggravate severe acute inflammatory intestinal disease and is ineffective in patients with extensive small intestinal disease eg. celiac sprue.)
- Previous hypersensitivity to any of the ingredient in the syrup.
- Known idiosyncrasy to commonly used excipients.
- Porphyria cutanea tarda.
- Uncontrolled parathyroid disease, sickle cell patient

#### **4.4 Special warnings and precautions for use**

Do not use if the tube membrane is already perforated. Do not use on inflamed or broken skin.

Avoid contact with mouth, mucous membranes and eyes and wash hands immediately after use. Discontinue use if irritation develops. If symptoms persist after four weeks, a doctor should be consulted.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

- Compounds containing Calcium and Magnesium including antacids and mineral supplements and bicarbonates, carbonates, oxalates or phosphates, may also impair the absorption of iron by the formation of insoluble complexes.
- Similarly absorption of both iron & tetracyclines is diminished when they are taken concomitantly by mouth. If treatment with both drugs is required, a time interval of about 2 to 3 hours should be allowed between them.
- Avoid milk and dairy products at least for 2 hours.
- Some agents such as Ascorbic Acid & Citric Acid may actually increase the absorption of iron.
- The response to iron may be delayed in patients receiving concomitant parenteral chloramphenicol therapy.
- Iron salts can decrease the absorption of bisphosphonates, fluoroquinolones, levodopa, methyldopa, penicillamine and tetracycline.
- Iron salts may reduce the efficacy of thyroxine.

#### **4.6 Pregnancy and lactation**

Can be used in Iron deficiency anaemia during Pregnancy and Lactation after considering risk benefit ratio

#### **4.7 Effects on Ability to drive and use machines**

Haepler<sup>®</sup> blood capsules have no influence on the ability to drive or use machines.

#### 4.8 Undesirable effects

- Ferrous Fumarate: In high or toxic doses or poisoning - Gastrointestinal irritation, abdominal pain with nausea, vomiting, and either diarrhoea or constipation. Cardiovascular disorders such as hypotension, tachycardia, and metabolic changes including acidosis and hypoglycemia. CNS depression ranges from lethargy to coma.
- Folic acid: Almost non-toxic in man and no adverse effects have been reported except a rare and doubtful allergic reaction.
- Pyridoxine may increase the peripheral metabolism of levodopa reducing therapeutic efficiency in patients with Parkinson's disease.
- Cyanocobalamin: No known side effects even with very large doses.

#### 4.9 Overdose

The most sign & Symptoms of overdosage are Gastrointestinal irritation, abdominal pain with nausea, vomiting and either diarrhoea or constipation. Cardiovascular disorders such as hypotension, tachycardia, metabolic changes including acidosis and hypoglycemia. CNS depression ranging from lethargy to coma. Vomiting is induced immediately followed by parenteral injection of desferroxamine mesylate and then gastric lavage. In the meantime give milk and/or 5% sodium bicarbonate solution by mouth. Fluid replacement is essential. Other measures include symptomatic management and therapy of metabolic and cardiovascular disorder

### 5. PHARMACOLOGICAL PROPERTIES

#### 5.1 Pharmacodynamic properties

**Pharmacotherapeutic group:** Hematinic Capsules

**ATC code:** A11JB

**Ferric Ammonium Citrate:** Ferric Ammonium Citrate is one of the best-tolerated iron supplements. It rapidly supplements elemental iron so that iron deficiency is quickly controlled, thus leading to faster correction of anaemia and replenishment of tissue iron stores.

Each 5ml of Haepler® Blood Tonic provides 69mg of Ferric Ammonium Citrate that is equivalent to 14 mg of elemental iron. Thus, when an adult takes the recommended dosage of 1 tablespoonful twice a day, he gets 28 mg of elemental iron which is sufficient amount of elemental iron per day for an adult patient suffering from mild to moderate iron deficiency.

**Folic Acid:** It is so called because it is present in green leaves, which is chemically pteroylglutamic acid. It is a growth promoter. It helps in the synthesis of Deoxyribose Nucleic Acid (DNA), the building block of life. It is essential for proper maturation of red cells. It helps in inter conversion and metabolism of amino acids.

**Cyanocobalamin:** Cyanocobalamin (Vitamin B12) is a complex organo-metallic compound in which a cobalt atom is placed within a corrin ring. It plays an important role in synthesis of DNA and maturation of red cells. It acts as a co-enzyme in certain steps necessary for genetic replication.

#### 5.2 Pharmacokinetic properties

Iron (Ferric Ammonium Citrate):

Iron irregularly and incompletely absorbed from the gastrointestinal tract, the main sites of absorption being the duodenum and jejunum. Absorption is aided by the acid section of the stomach and by some dietary acids (such as ascorbic acid) and is more readily affected when the iron is in the ferrous state or is part of the haem complex (haem-iron). Absorption is also increased in conditions of iron deficiency or in the fasting state but is decreased if the body stores are overloaded. Only about 5 to 15% of the iron ingested in food is normally absorbed.

Most absorbed iron is bound to transferrin and transported to the bone marrow where it is incorporated into haemoglobin; the remainder is contained within the storage forms,

ferritin or haemosiderin, or as myoglobin, with smaller amounts occurring in haem-containing enzymes or in plasma bound to transferrin. Only very small amounts of iron are excreted as the majority released after the destruction of the haemoglobin molecule is re-used.

Folic Acid:

Folic acid is rapidly absorbed from the gastro-intestinal tract, mainly from the duodenum and jejunum. Folic acid administered therapeutically enters the portal circulation largely unchanged since it is a poor substrate for reduction by dihydrofolate reductase. It is converted to the metabolically active form 5-methyltetrahydrofolate in the plasma and liver.

The principal storage site of folate is the liver; it is also actively concentrated in the CSF.

Folate undergoes enterohepatic circulation. Folate metabolites are eliminated in the urine and folate in excess of body requirements is excreted unchanged in the urine. Folate is distributed into breast milk. Folic acid is removed by haemodialysis.

Cyanocobalamin: Cyanocobalamin bind to intrinsic factor, a glycoprotein secreted by the gastric mucosa, and are then actively absorbed from the gastro-intestinal tract. Absorption is impaired in patients with an absence of intrinsic factor, with a malabsorption syndrome or with disease or abnormality of the gut, or after gastrectomy. Absorption from the gastro-intestinal tract can also occur by passive diffusion. Cyanocobalamin is extensively bound to specific plasma proteins called transcobalamins; transcobalamin II appears to be involved in the rapid transport of the cobalamins to tissues. Cyanocobalamin is stored in the liver, excreted in the bile and undergoes extensive enterohepatic recycling; part of an administered dose is excreted in the urine, most of it in the first 8 hours; urinary excretion, however, accounts for only a small fraction in the reduction of total body stores acquired by dietary means. Cyanocobalamin diffuses across the placenta and also appears in breast milk.

### **5.3 Preclinical safety data**

#### Toxicology

Iron: Iron is a general cellular poison and is directly corrosive to the GI mucosa. Cellular toxicity The absorption of excessive quantities of ingested iron results in systemic iron toxicity. Severe overdose causes impaired oxidative phosphorylation and mitochondrial dysfunction, which can result in cellular death. The liver is one of the organs most affected by iron toxicity, but other organs such as the heart, kidneys, lungs, and the hematologic systems also may be impaired.

**MILD TO MODERATE POISONING:** Vomiting and diarrhea may occur within 6 hours of ingestion.

**SEVERE POISONING:** Severe vomiting and diarrhea, lethargy, metabolic acidosis, shock, GI hemorrhage, coma, seizures, hepatotoxicity, and late onset GI strictures.

Folic acid: The risk of toxicity from folic acid is low, because folate is a water-soluble vitamin and is regularly removed from the body through urine. One potential issue associated with high dosages of folic acid is that it has a masking effect on the diagnosis of pernicious anaemia (vitamin B12 deficiency).

Vitamin B12:

Vitamin B12 is usually considered a non-toxic substance. Even taking it by injection at high doses does not seem to increase the risk for toxicity

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Lactose Anhydrous 200 Mesh(To Be Dried)  
Microcrystalline Cellulose (Ph 101)  
Aerosil 200  
Sodium Lauryl Sulphate  
Magnesium Stearate

### **6.2 Incompatibilities**

None known

### **6.3 Shelf life**

24 months

### **6.4 Special precautions for storage**

Store below 30°C, and protect from light and moisture.

### **6.5 Nature and contents of container <and special equipment for use, administration or implantation>**

Haepler is presented in Hard Gelatin Capsule of 3x 10 Capsule

### **6.6 Special precautions for disposal <and other handling>**

No special requirements

## **7. APPLICANT/ MANUFACTURER**

Drugfield Pharmaceuticals Limited  
Lynson Chemical Avenue Km38,  
Lagos-Abeokuta Expressway  
Sango-Otta, Ogun State, Nigeria  
Tel: +2348033513989  
Email:Info@drugfieldpharma.com

