

# SMPC FOR JESSY HEMOGLOBIN SYRUP

## 1. Name of the medicinal product

Jessy Hemoglobin Syrup

## 2. Qualitative and quantitative composition

Each 5ml contains

Hemoglobin 5%w/v

Chelated iron 0.3%w/v

Cyanocobalamin 0.0004%w/v

Malt extract 2%w/v

For a full list of excipients, see section 6.1

## 3. Pharmaceutical form

Oral Liquid

A dark brown viscous liquid with a faint chocolate flavour

## 4. Clinical particulars

### 4.1 Therapeutic indications

Jessy hemoglobin Syrup is indicated for

- Iron deficiency anaemia due to chronic blood loss, hook-worm infestation, inadequate intake of iron, etc.
- Dimorphic anaemia due to deficiency of Iron, Folic Acid and /or Vitamin B12 (Cyanocobalamin).
- Anemia of pregnancy and lactation.
- Tonic in general weakness, lack of appetite, rundown conditions and convalescence. • Post surgery and other debilitated states.

### 4.2 Posology and method of administration

For oral administration:

#### Adults

2 teaspoonful 4 times daily

#### **Children (5-12 years)**

1 teaspoonful 4 times daily

#### **Children (2-5 years)**

1 teaspoonful 2 times daily

### 4.3 Contraindications

- Primary (idiopathic) or secondary iron storage disease.

## SMPC FOR JESSY HEMOGLOBIN SYRUP

- Anaemia associated with ineffective erythropoiesis, marrow hypoplasia, sideroblastic change, uncomplicated Cyanocobalamin deficiency.
- Intestinal disease (oral iron may aggravate severe acute inflammatory intestinal disease and is ineffective in patients with extensive small intestinal disease eg. celiac sprue.) Previous hypersensitivity to any of the ingredient in the syrup.
- Known idiosyncrasy to commonly used excipients.
- Porphyria cutanea tarda.
- Uncontrolled parathyroid disease, sickle cell patients.

### 4.4 Special warnings and precautions for use

Care should be taken in patients with iron storage or iron absorption diseases such as haemochromatosis, hemoglobinopathies or existing gastro-intestinal diseases such as inflammatory bowel disease, intestinal strictures and diverticulae.

### 4.5 Interaction with other medicinal products and other forms of interaction

If you use other drugs or over the counter products at the same time, the effects of Jessy Hemoglobin Syrup may change. This may increase your risk for side-effects or cause your drug not to work properly. Tell your doctor about all the drugs, vitamins, and herbal supplements you are using, so that you doctor can help you prevent or manage drug interactions. Jessy Hemoglobin Syrup may interact with the following drugs and products:

- Arsenic trioxide
- Chloramphenicol
- Folic acid
- Furosemide
- Gabapentin
- Hydrochlorothiazide
- Levothyroxine
- Lisinopril

Compounds containing Calcium and Magnesium including antacids and mineral supplements and bicarbonates, carbonates, oxalates or phosphates, may also impair the absorption of iron by the formation of insoluble complexes.

Similarly, absorption of both iron & tetracyclines is diminished when they are taken concomitantly by mouth. If treatment with both drugs is required, a time interval of about 2 to 3 hours should be allowed between them.

- Avoid milk and dairy products at least for 2 hours.
- Some agents such as Ascorbic Acid & Citric Acid may actually increase the absorption of iron.
- The response to iron may be delayed in patients receiving concomitant parenteral chloramphenicol therapy.
- Iron salts can decrease the absorption of bisphosphonates, fluoroquinolones, levodopa, methyl dopa, penicillamine and tetracycline.
- Iron salts may reduce the efficacy of thyroxine.

# SMPC FOR JESSY HEMOGLOBIN SYRUP

## 4.6 Fertility, pregnancy and lactation

Can be used in Iron deficiency anaemia during Pregnancy and Lactation after considering risk benefit ratio

## 4.7 Effects on ability to drive and use machines

Jessy Hemoglobin Syrup has no influence on the ability to drive or use machines.

## 4.8 Undesirable effects

### Hemoglobin

These side-effects are possible, but do not always occur. Some of the side-effects may be rare but serious. Consult your doctor if you observe any of the following side-effects, especially if they do not go away.

Diarrhea

Swelling

Rapid weight gain

Itching or mild rash

Jessy Hemoglobin Syrup may also cause side-effects not listed here.

### Ferric Ammonium Citrate:

In high or toxic doses or poisoning - Gastrointestinal irritation, abdominal pain with nausea, vomiting and either diarrhoea or constipation. Cardiovascular disorders such as hypotension, tachycardia, metabolic changes including acidosis and hypoglycemia. CNS depression ranging from lethargy to coma.

### Cyanocobalamin:

No known side effects even with very large doses.

## 4.9 Overdose

The most sign & Symptoms of overdosage are Gastrointestinal irritation, abdominal pain with nausea, vomiting and either diarrhoea or constipation. Cardiovascular disorders such as hypotension, tachycardia, metabolic changes including acidosis and hypoglycemia. CNS depression ranging from lethargy to coma.

Vomiting is induced immediately followed by parenteral injection of desferroxamine mesylate and then gastric lavage. In the meantime, give milk and/or 5% sodium bicarbonate solution by mouth. Fluid replacement is essential. Other measures include symptomatic management and therapy of metabolic and cardiovascular disorders.

## 5. Pharmacological properties

### 5.1 Pharmacodynamic properties

Pharmacotherapeutics group: Hematinic syrup, ATC code: B03AE01

### Ferric Ammonium Citrate:

Ferric Ammonium Citrate is one of the best-tolerated iron supplements. It rapidly supplements elemental iron so that iron deficiency is quickly controlled, thus leading to faster correction of anaemia and replenishment of tissue iron stores.

# SMPC FOR JESSY HEMOGLOBIN SYRUP

## **Cyanocobalamin:**

Cyanocobalamin (Vitamin B12) is a complex organo-metallic compound in which a cobalt atom is placed within a corrin ring. It plays an important role in synthesis of DNA and maturation of red cells. It acts as a co-enzyme in certain steps necessary for genetic replication.

## **Pharmacokinetic properties Iron (Ferric Ammonium Citrate):**

Iron irregularly and incompletely absorbed from the gastrointestinal tract, the main sites of absorption being the duodenum and jejunum. Absorption is aided by the acid section of the stomach and by some dietary acids (such as ascorbic acid) and is more readily affected when the iron is in the ferrous state or is part of the haem complex (haem-iron). Absorption is also increased in conditions of iron deficiency or in the fasting state but is decreased if the body stores are overloaded. Only about 5 to 15% of the iron ingested in food is normally absorbed.

Most absorbed iron is bound to transferrin and transported to the bone marrow where it is incorporated into haemoglobin; the remainder is contained within the storage forms, ferritin or haemosiderin, or as myoglobin, with smaller amounts occurring in haem-containing enzymes or in plasma bound to transferrin.

Only very small amounts of iron are excreted as the majority released after the destruction of the haemoglobin molecule is re-used.

## **Cyanocobalamin:**

Cyanocobalamin bind to intrinsic factor, a glycoprotein secreted by the gastric mucosa, and are then actively absorbed from the gastro-intestinal tract. Absorption is impaired in patients with an absence of intrinsic factor, with a malabsorption syndrome or with disease or abnormality of the gut, or after gastrectomy. Absorption from the gastro-intestinal tract can also occur by passive diffusion.

Cyanocobalamin is extensively bound to specific plasma proteins called transcobalamins; transcobalamin II appears to be involved in the rapid transport of the cobalamins to tissues. Cyanocobalamin is stored in the liver, excreted in the bile and undergoes extensive enterohepatic recycling; part of an administered dose is excreted in the urine, most of it in the first 8 hours; urinary excretion, however, accounts for only a small fraction in the reduction of total body stores acquired by dietary means. Cyanocobalamin diffuses across the placenta and also appears in breast milk.

## **5.3 Preclinical safety data**

### **Toxicology Iron:**

Iron is a general cellular poison and is directly corrosive to the GI mucosa.

### **Cellular toxicity**

The absorption of excessive quantities of ingested iron results in systemic iron toxicity. Severe overdose causes impaired oxidative phosphorylation and mitochondrial dysfunction, which can result in cellular death. The liver is one of the organs most affected by iron toxicity, but other organs such as the heart, kidneys, lungs, and the hematologic systems also may be impaired.

# SMPC FOR JESSY HEMOGLOBIN SYRUP

**Mild To Moderate Poisoning:** Vomiting and diarrhea may occur within 6 hours of ingestion.

**Severe Poisoning:** Severe vomiting and diarrhea, lethargy, metabolic acidosis, shock, GI hemorrhage, coma, seizures, hepatotoxicity, and late onset GI strictures.

## **Vitamin B12:**

Vitamin B12 is usually considered a non-toxic substance. Even taking it by injection at high doses does not seem to increase the risk for toxicity.

## **Alcohol:**

Even though alcohol contributed to severe toxicity, Jessy Hemoglobin Syrup contains very low level of alcohol i.e. may not lead to toxicity even during pregnancy. As the National Institute of Health and Clinical Trial Excellence (NICE) recommends that women should avoid alcohol during the 1st trimester, women who chose to drink alcohol are advised to drink no more than 1 – 2 UK units of alcohol once or twice a week. They further state that, “Although there is uncertainty regarding a safe level of alcohol consumption in pregnancy, at this low level there is no evidence of harm to the unborn baby”.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Xanthan gum  
Ethanol  
Methyl paraben  
Propyl paraben  
Sodium benzoate  
Chocolate flavour

### **6.2 Incompatibilities**

Not applicable

### **6.3 Shelf life**

Unopened: 2 years

After opening: 6 months

### **6.4 Special precautions for storage**

Store below 30°C.

Do not refrigerate or freeze.

### **6.5 Nature and contents of container**

200ml amber-coloured Glass bottles with 28mm plastic caps

### **6.6 Special precautions for disposal and other handling**

No special requirements

# SMPC FOR JESSY HEMOGLOBIN SYRUP

**7. Marketing authorisation holder**

Jessy Pharmaceutical company Limited

**8. Marketing authorisation number(s)**

A11-1301

**9. Date of first authorisation/renewal of the authorisation**

29/09/2022

**10. Date of revision of the text**

28/09/2027