10 X 10 Capsules

JAFFKIN TRAMADOL CAPSULES 50MG

Tramadol Hydrochloride BP 50mg





IMPRIN TRAMADOL CAPSULES SOMG Tramadol Hydrochloride BP 50mg

Composition:

Each capsule contains:

Tramadol Hydrochloride BP 50mg

Excipients q.s

Approved colours used in capsule shells

Dosage: As directed by the Physician Storage: Store below 30°C in a cool and dry place. Protect from light and moisture.

10 X 10 Capsules

JAFFKIN

Tramadol Hydrochloride BP 50mg

TRAMADOL CAPSULES 50MG

Tramadol Hydrochloride BP 50mg





JAFFKIN
TRAMADOL CAPSULES 50MG



NAFDAC Reg. No.:

Batch No.: Mfg. Date:

Exp. Date:



Manufactured by:



TOWA PHARMA PVT. LTD.

Survey No-17, Jaliyanamath Village, Dehgam Rakhiyal Road, Ta. Dehgam, Dist. Gandhinagar- 382305, Guiarat State. India.

TRAMADOL CAPSULES 50MG Tramadol Hydrochloride BP 50mg

Each capsule contains: ramadol Hydrochloride BP 50mg regionats q.s

Approved colours used in capsule shells

DOSAGE: As directed by the Physician **Storage:** Store below 30°C in a cool and dry place. Protect from light and moisture. TOWA

NAFDAC Reg. No.:



JAFFKIN TRAMADOL CAPSULES 50MG

Tramadol Hydrochloride BP 50mg

Composition: Each capsule contains:

Tramadol Hydrochloride B.P. 50mg

Excipients q.s.

Approved colours used in capsules shell

Pharmacological Action: JAFFKIN TRAMADOL is an analgesic with dual mode of action. It is an opioid analgesic with low affinity for opioid receptors. It acts by blocking the re-uptake of nor-adrenaline and serotonin synapses in the descending neural pathways, which inhibit pain responses in the spinal cord. This non opioid action contributes significantly to the analgesic effect. Due to its dual mode of action, JAFFKIN TRAMADOL has pronounced opioid adverse effect at therapeutic doses.

It has little euphoric effect and therefore lower abuse potential than other opioids.

Indications: Moderate to Severe pain, diagnostic procedure and surgical pains

Dosage: 50-100mg every 6-8 Hours; not exceeding 400mg per day. Not recommended for children, pregnant woman and lactating mothers. Reduced dose may be necessary for the elderly.

Contra-Indications: History of Hypersensitivity to Tramadol or any of the components of the preparation, acute intoxication with alcohol hypnotics, analgesics, opiods or other CNS active drugs, concurrent use with MAOIs, pregnancy, Lactation and children less than 14 years.

Side Effects: Sedation, sweating, nausea, vomiting, animation, stupor.

Precaution: Impaired renal or hepatic function (reduce dose); history of epilepsy or those susceptible to seizures; may impair ability to drive or operate machinery especially in conjunction with alcohol.

Packing: 10 Capsules per blister; 100 blister per clinical carton. Store in a cool dry place, away from direct sunlight. Keep all medicines out of reach of children.

Manufactured by:

NAFDAC Reg. No.:



TOWA PHARMA PVT. LTD.

Survey No-17, Jaliyanamath Village, Dehgam Rakhiyal Road, Ta. Dehgam, Dist. Gandhinagar- 382305, Gujarat State, India.