

# Summary of Product Characteristics

## 1 NAME OF THE MEDICINAL PRODUCT

Hydrocortisone 1 % w/w Cream

## 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Hydrocortisone 1 % w/w Cream contains Hydrocortisone 1% w/w.

### Excipient(s) with known effect

Contains 0.1% w/w chlorocresol and 7.2% w/w cetostearyl alcohol.

For a full list of excipients, see section 6.1.

## 3 PHARMACEUTICAL FORM

Cream

Smooth white aqueous cream.

## 4 CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Hydrocortisone 1 % w/w Cream is recommended for the topical treatment of contact dermatitis, irritant dermatitis, insect bite reactions, mild to moderate eczema.

### 4.2 Posology and method of administration

Hydrocortisone 1 % w/w Cream is applied topically.

*Recommended dosage schedules:*

*Adults:* Use sparingly over a small area once or twice daily for a maximum period of one week.

*Children:* Hydrocortisone 1 % w/w Cream is not recommended for use in children under 12 years without medical supervision.

### 4.3 Contraindications

Hydrocortisone 1 % w/w Cream is contraindicated for use on the eyes or face, the ano-genital area or on broken or infected skin including cold sores, acne and athletes' foot.

Use in the presence of untreated infections of bacterial, viral, tuberculous or fungal origin.

Use in acne vulgaris, acne rosacea or in perioral dermatoses.

Hypersensitivity to the preparation.

Use in plaque psoriasis.

### 4.4 Special warnings and precautions for use

Hydrocortisone 1 % w/w Cream is not recommended for use in children under 12 years, except under medical supervision.

Prolonged use of uninterrupted occlusion or use with extensive occlusive dressings may suppress adrenocortical function.

Continuous application without interruption will result in local atrophy of the skin, striae, and superficial vascular dilatation, particularly on the face.

Long term continuous or inappropriate use of topical steroids can result in the development of rebound flares after stopping treatment (topical steroid withdrawal syndrome). A severe form of rebound flare can develop which takes the form of a dermatitis with intense redness, stinging and burning that can spread beyond the initial treatment area. It is more likely to occur when delicate skin sites such as the face and flexures are treated. Should there be a reoccurrence of the condition within

days to weeks after successful treatment a withdrawal reaction should be suspected. Reapplication should be with caution and specialist advise is recommended in these cases or other treatment options should be considered.

#### Visual disturbance:

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

The inactive ingredients, cetostearyl alcohol may cause local skin reactions (e.g. contact dermatitis) and chlorocresol may cause allergic reactions.

### 4.5 Interaction with other medicinal products and other forms of interaction

None known.

### 4.6 Fertility, pregnancy and lactation

#### Pregnancy:

Topical administration of corticosteroids to pregnant animals can cause abnormalities of foetal development. The relevance of this finding to human beings has not been established. The product should not be used in pregnancy unless considered essential by the physician, and then over as small an area and for as short a time as possible.

### 4.7 Effects on ability to drive and use machines

No adverse effects reported.

### 4.8 Undesirable effects

Adverse events which have been associated with topical corticosteroids are given below, tabulated by system organ class and frequency. Frequencies are defined as: Very common ( $\geq 1/10$ ); Common ( $\geq 1/100$  and  $< 1/10$ ); Uncommon ( $\geq 1/1000$  and  $< 1/100$ ); Rare ( $\geq 1/10,000$  and  $< 1/1000$ ); Very rare ( $< 1/10,000$ ); Not known (cannot be estimated from the available data). Within each frequency grouping, adverse events are presented in order of decreasing seriousness.

System Organ Class	Frequency	Adverse Events
Skin and Subcutaneous Tissue Disorders	Very rare	Local irritation, itching and burning
		Striae may occur especially in intertriginous areas.
	Not known	Urticaria
		Withdrawal reactions - redness of the skin which may extend to areas beyond the initial affected area, burning or stinging sensation, itch, skin peeling, oozing pustules. (see section 4.4)
Endocrine Disorders	Not known	Pheochromocytoma crisis
Eye Disorders	Not known	Vision, blurred (see section 4.4)

#### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRC Pharmacovigilance; [www.hpra.ie](http://www.hpra.ie).

### 4.9 Overdose

No special procedures or antidote are likely to be necessary.

Topical overdose may occur where large amounts are applied, or where smaller amounts are applied in occlusive conditions. Overdose ultimately leads to adrenocortical suppression. Treatment is by gradual reduction of the amounts used. The side effects of overuse i.e. skin atrophy, striae etc. may not be reversed.

## 5 PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

**Pharmacotherapeutic Group:** Corticosteroids, dermatological preparations; corticosteroids, weak (group I); **ATC Code:** D07AA02

Hydrocortisone, as a mild topical corticosteroid, is used in the treatment of many skin conditions such as eczema, atopic dermatitis, contact dermatitis, seborrhoeic dermatitis and some forms of psoriasis. When administered by topical application, particularly under an occlusive dressing or when the skin is broken, sufficient corticosteroid may be absorbed to give systemic effects.

### 5.2 Pharmacokinetic properties

The literature states that absorption does occur through the skin, particularly denuded skin. This absorption is not usually of clinical significance as hydrocortisone topically has only rarely been associated with side effects resulting from pituitary-adrenal suppression.

Topically applied steroids may be absorbed to a significant extent if applied to broken skin, to very large areas, or under occlusive dressings.

If systemic absorption occurs, hydrocortisone is extensively bound to plasma proteins, mainly to globulin and less so to albumin. Corticosteroids are metabolised mainly in the liver but also in the kidney, and are excreted in the urine.

### 5.3 Preclinical safety data

No further information available.

## 6 PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

White soft paraffin  
Cetomacrogol emulsifying wax (containing cetostearyl alcohol and macrogol cetostearyl ether)  
Liquid paraffin  
Chlorocresol  
Purified water

### 6.2 Incompatibilities

Not applicable.

### 6.3 Shelf life

Three years.

### 6.4 Special precautions for storage

Store below 25°C. Do not freeze.

### 6.5 Nature and contents of container

Hydrocortisone 1 % w/w Cream is packaged in internally lacquered aluminium tubes with polyethylene closures containing 15g of cream.

**6.6 Special precautions for disposal of a used medicinal product or waste materials derived from such medicinal product and other handling of the product**

No special requirements.

**7 MARKETING AUTHORISATION HOLDER**

Norbarc  
Pharmaceutical  
Ltd, No.7  
Abeokuta Street  
Ebute Metta

**Manufacturer; Marine Lifesciences,  
Plot no: 30, Phase III (Ext),HPSIDC, Baddi, Dist. Solan (H.P) India**