



Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bptbrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

1. Name of the medicinal product

1.1. Name of the medicinal product:

Generic Name/INN Name: Glimepiride & Metformin Tablets

Trade Name: GLAVAMET GR

1.2 Strength:

Glimepiride USP.....2 mg


Metformin Hydrochloride BP.....500 mg (In Extended release form)

1.3 Pharmaceutical form:

Tablets (Solid Oral Dosage form)

2. Qualitative and Quantitative composition:

Sr. No.	Ingredients	Spec.	Qty. per Tablet (in mg)	% w/w	Function
For Glimepiride 2 mg IR layer					
Materials used For Uncoated Tablets					
Materials used For Dry mixing					
1.	Glimepiride	USP	2.000	2.50	Active
2.	Microcrystalline Cellulose Grade 102	BP	67.584	84.48	Diluent
3.	Crosspovidone	BP	10.000	12.50	Disintegrant
4.	Col. Iron oxide of red IDA-LA-COL	IH	0.096	0.12	Colorant
5.	Magnesium Stearate	BP	0.320	0.40	Lubricant
Total weight : 80.000 mg					
Loss of Moisture and Solvent: 0.000 mg					
Net weight: 80.000 mg					
For Metformin HCl 500 mg ER layer					
1.	Metformin Hydrochloride	BP	500.000	72.89	Active
2.	Maize Starch	BP	25.500	3.72	Diluent
3.	Hypromellose Type 2208	USP	90.000	13.12	Sustained Release agent
Materials used For binding (Povidone slurry preparation)					
1.	Povidone K-30	BP	10.000	1.46	Binder
2.	Isopropyl Alcohol	BP	10.000	---	Solvent
3.	Purified Water	BP	67.900	---	Solvent
Materials used as Lubricants					
1.	Hypromellose Type 2208	USP	45.000	6.56	Sustained Release agent
2.	Colloidal Silicon Dioxide	BP	7.000	1.02	Glidant
3.	Magnesium Stearate	BP	8.000	1.17	Lubricant
4.	Titanium Dioxide	BP	0.500	0.07	Colorant





Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bpibrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

Total weight: 763.900 mg
Loss of Moisture and Solvent: 77.900 mg
Net weight: 686.000 mg
Net weight of Bilayer uncoated Tablet: 766.00 mg

3. Pharmaceutical form:

Dosage Form: Tablets (Solid Oral Dosage form)

Visual & Physical characteristics of the product: White & peach colour, bilayer capsule shaped biconvex, uncoated tablets with a breakline on peach coloured side.

4. Clinical particulars

4.1. Therapeutic indications:

Glavamet GR is indicated as an adjunct to diet and exercise to improve glycaemic control in patients with type-2 diabetes who are already treated with a combination of glimepiride and metformin or whose diabetes is not adequately controlled with metformin alone, or for those patients who have initially responded to glimepiride alone and require additional glycaemic control.

4.2. Posology and method of administration:

General


Dosage should be individualized on the basis of both effectiveness and tolerance. The combination should be given once daily with meals and should be started at a low dose. The initial recommended dose is one tablet once daily with breakfast or first main meal of the day.

Starting dose for patients inadequately controlled on metformin monotherapy Glavamet GR may be initiated once daily, and gradually titrated after assessing the therapeutic response.

Starting dose for patients who initially responded to glimepiride monotherapy and require additional glycaemic control Based on the initial starting dose of glimepiride (1 or 2 mg), Glavamet GR may be initiated once daily, and gradually titrated after assessing the therapeutic response.

Starting dose for patients switching from combination therapy of glimepiride plus metformin as separate tablets Glavamet GR may be initiated based on the dose of glimepiride and metformin already being take

GLAVAMET GR (Glimepiride & Metformin Tablets)





Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bplbrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

Maximum Recommended Dose: The maximum recommended dose for glimepiride is 8 mg daily. The maximum recommended daily dose for metformin sustained-release is 2000 mg in adults.

4.3. Contraindications:

Renal disease or renal dysfunction, as suggested by serum creatinine levels ≥ 1.5 mg/dL [males], ≥ 1.4 mg/dL [females] or abnormal creatinine clearance, which may also result from conditions such as cardiovascular collapse (shock), acute myocardial infarction, and septicaemia.

Hepatic impairment.

Known hypersensitivity to this product or any of its components.

Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma.

Diabetic ketoacidosis should be treated with insulin.

Patients undergoing radiologic studies involving intravascular administration of iodinated contrast materials, because the use of such products may result in acute alteration of renal function.

4.4. Special warnings and precautions for use:

Cardiac effects

The administration of oral hypoglycemic drugs (tolbutamide) has been reported to be associated with increased cardiovascular mortality as compared to treatment with diet alone or diet plus insulin. In view of close similarities between the oral hypoglycemic drugs, this warning also applies for glimepiride.

Lactic acidosis

Lactic acidosis is a rare, but serious, metabolic complication that can occur due to metformin accumulation during treatment with glimepiride and metformin combination therapy; when it occurs, it is fatal in approximately 50% of cases. When metformin is implicated as the cause of lactic acidosis, metformin plasma levels >5 $\mu\text{g/mL}$ are generally found. The reported incidence of lactic acidosis in patients receiving metformin hydrochloride is very low (approximately 0.03 cases/1000 patient-years, with approximately 0.015 fatal cases/1000 patient years). Reported cases have occurred primarily in diabetic patients with significant renal insufficiency and congestive heart failure.





Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bpibrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

Hypoglycemia

All sulphonylurea drugs are capable of producing severe hypoglycemia. Proper patient selection, dosage, and instructions are important to avoid hypoglycemic episodes.

Loss of control of blood glucose

When a patient stabilized on any diabetic regimen, is exposed to stress such as fever, trauma, infection, or surgery, a temporary loss of glycemic control may occur. At such times, it may be necessary to withhold the diabetic regime and temporarily administer insulin. The oral antidiabetic therapy may be reinstated after the acute episode is resolved.

Alcohol intake

Alcohol is known to potentiate the effect of metformin on lactate metabolism. Patients should be warned against excessive alcohol intake, while receiving metformin.

Hypoxic states

Cardiovascular collapse (shock) from whatever cause, acute congestive heart failure, acute myocardial infarction and other conditions characterized by hypoxemia have been associated with lactic acidosis and may also cause prerenal azotemia. When such events occur in patients on metformin therapy, the drug should be promptly discontinued.

Hemolytic anemia

Treatment of patients with glucose 6-phosphate dehydrogenase (G6PD) deficiency with sulphonylurea agents can lead to hemolytic anemia. Since glimepiride is a sulphonylurea agent, caution should be used in patients with G6PD deficiency and a non-sulphonylurea alternative should be considered.

Geriatric use

Metformin is known to be excreted by the kidneys, and because risk of serious adverse reactions to the drug is greater in patients with impaired renal function, glimepiride and metformin should be used only in patients with normal renal function. Because aging is associated with reduced renal function, glimepiride and metformin combination should be used with caution in the elderly.

4.5. Interaction with other medicinal products and other forms of interaction:

Cationic drugs: Certain medications used concomitantly with metformin may increase the risk of lactic acidosis. Cationic drugs that are eliminated by renal tubular secretions (e.g:

GLAVAMET GR (Glimepiride & Metformin Tablets)





SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

amiloride, digoxin, morphine, procainamide, quinidine, ranitidine, or vancomycin) may decrease metformin elimination by competing for common renal tubular transport systems. Hence, careful patient monitoring and dose adjustment of metformin/cationic drug is recommended.

Miconazole (systemic route, oromucosal gel) and Phenylbutazone (systemic route): Increases hypoglycemic effect of glimepiride.

Furosemide: A single-dose, metformin-furosemide drug interaction study in healthy subjects demonstrated that pharmacokinetic parameters of both compounds were affected by co-administration. Furosemide increased the metformin plasma and blood C_{max} by 22% and blood AUC by 15%, without any significant change in metformin renal clearance. When administered with metformin, the C_{max} and AUC of furosemide were 31% and 12% smaller, respectively, than when administered alone, and the terminal half-life was decreased by 32%, without any significant change in furosemide renal clearance.

Vitamin B12: Metformin may result in suboptimal oral vitamin B12 absorption by competitively blocking the calcium dependent binding of the intrinsic factor vitamin B12 complex to its receptor. The reaction very rarely results in pernicious anemia which is reversible with discontinuation of metformin and supplementation with vitamin B12.

Nifedipine: Nifedipine appears to enhance the absorption of metformin, it increases plasma metformin C_{max} and AUC by 20% and 9% respectively and increases the amount of metformin excreted in the urine.

Danazol: If the use of this active substance cannot be avoided, the patient's urine and blood glucose must be monitored. It may be necessary to adjust the dose of glimepiride and metformin during and after treatment with danazol.

Salicylates: If salicylates are administered or discontinued in patients receiving oral antidiabetic agents, patients should be monitored for hypoglycemia or loss of blood glucose control.

Thiazide: Interactions between thiazide diuretics and oral antidiabetic agents decreases insulin sensitivity thereby leading to glucose intolerance and hyperglycemia. Hence diabetic patients should be monitored closely.



Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bplbrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

4.6. Pregnancy and lactation:

Pregnancy

Abnormal blood glucose levels during pregnancy are associated with the higher incidence of congenital abnormalities. Most experts suggest insulin be used to maintain the blood glucose levels as close to normal as possible. The use of glimepiride and metformin combination is not recommended for use in pregnancy.

Lactation

Studies in lactating rats show that metformin is excreted into milk and reaches levels comparable to those in plasma. Similar studies have not been conducted on nursing mothers. Also Glimepiride should not be used by breast-feeding mothers. Hence, the use of glimepiride and metformin combination is not recommended for use in lactating mothers, and if the diet alone is inadequate for controlling blood glucose, insulin therapy should be considered.

4.7. Effects on ability to drive and use machines:

For Glimepiride

Alertness and reactions may be impaired due to hypo or hyperglycemia, especially when beginning or after altering treatment or when glimepiride is not taken regularly, this may, for example, affect the ability to drive or to operate machinery.

For Metformin


Metformin monotherapy does not cause hypoglycemia and therefore has no effect on the ability to drive or to use machine. However patient should be alerted to the risk of hypoglycemia when metformin is used in combination with other antidiabetic agent.

4.8. Undesirable effects:

Gastrointestinal disturbances: Nausea, diarrhea, abdominal pain, constipation, vomiting and metallic taste in mouth may occur. These reactions are generally dose related and disappear when the dose is reduced.

Dermatological effects: Rash, pruritus, urticaria, erythema & flushing.

Miscellaneous: Headache and dizziness.





SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)


Hematologic Reactions: Leukopenia, agranulocytosis, thrombocytopenia, hemolytic anemia, aplastic anemia, and pancytopenia have been reported with sulfonylureas, including glimepiride.

Metabolic Reactions: Hepatic porphyria reactions and disulfiram-like reactions have been reported with sulfonylureas, including glimepiride. Cases of hyponatremia have been reported with glimepiride and all other sulfonylureas, most often in patients who are on other medications or have medical conditions known to cause hyponatremia or increase release of antidiuretic hormone. The syndrome of inappropriate antidiuretic hormone (SIADH) secretion has been reported with sulfonylureas, including glimepiride, and it has been suggested that certain sulfonylureas may augment the peripheral (antidiuretic) action of ADH and/or increase release of ADH.

Hypoglycemia: Glimepiride appears to be associated with a low incidence of hypoglycemia. Although Glimepiride has the potential to produce adverse cardiovascular effects, glimepiride has been an established agent for the treatment of type-2 diabetes for a number of years without producing adverse cardiovascular effects.

4.9. Overdose:

Overdosage of sulfonylureas, including glimepiride, can produce hypoglycemia. Mild hypoglycemic symptoms without loss of consciousness or neurologic findings should be treated aggressively with oral glucose and adjustments in drug dosage and/or meal patterns. Close monitoring should continue until the physician is assured that the patient is out of danger. Severe hypoglycemic reactions with coma, seizure, or other neurological impairment occur infrequently, but constitute medical emergencies requiring immediate hospitalization. If hypoglycemic coma is diagnosed or suspected, the patient should be given a rapid IV injection of concentrated (50%) glucose solution. This should be followed by a continuous infusion of a more dilute (10%) glucose solution at a rate that will maintain the blood glucose level above 100 mg/dL. Patients should be closely monitored for a minimum of 24 to 48 hours, because hypoglycemia may recur after apparent clinical recovery. Lactic acidosis is a rare, but serious, metabolic complication that can occur if metformin accumulates during treatment due to overdosing. Strict monitoring should be continued until the doctor is sure that the patient is out of danger.





SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

5. Pharmacological properties:

5.1. Pharmacodynamic properties:

Pharmacotherapeutic group: Anti-diabetic drug

ATC-CODE: A10BB12 - glimepiride

A10BA02 - metformin

Glimepiride: The primary mechanism of action of glimepiride in lowering blood glucose appears to be dependent on stimulating the release of insulin from functioning pancreatic beta cells. In addition, extra-pancreatic effects may also play a role in the activity of sulphonylureas such as glimepiride.

Metformin: Metformin improves glucose tolerance in patients with type-2 diabetes (NIDDM), lowering both basal and postprandial plasma glucose. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. Hence, the combination of glimepiride and metformin sustained-release complements each other and provides better glycemic control in management of type-2 diabetes and probably in the prevention of its associated macrovascular and microvascular complications.

5.2. Pharmacokinetic properties:

Absorption

Glimepiride: After oral administration, glimepiride is completely absorbed from the GI tract. Studies have shown significant absorption of glimepiride within 1 hour after administration and peak drug levels (C_{max}) at 2 to 3 hours. When glimepiride was given with meals, the mean T_{max} (time to reach C_{max}) was slightly increased (12%) and the mean C_{max} and AUC (area under the curve) were slightly decreased (8% and 9%, respectively).

Metformin sustained-release: The absolute bioavailability of a metformin 500-mg tablet given under fasting conditions is approximately 50-60%. Following a single oral dose of metformin sustained-release, C_{max} is achieved within 4-8 hours. Peak plasma levels are approximately 20% lower compared to the same dose of metformin immediate release, however, the extent of absorption (as measured by AUC) is similar to immediate release. Both high and low fat meals had the same effect on the pharmacokinetics of sustained release.





Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bplbrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

Distribution

Glimepiride: After intravenous dosing in normal subjects, the volume of distribution (Vd) was 8.8 L (113 mL/kg). Protein binding was greater than 99.5%.

Metformin sustained-release: Metformin is negligibly bound to plasma proteins, in contrast to sulphonylureas, which are more than 90% protein bound. Metformin partitions into erythrocytes, most likely as a function of time. Distribution studies with metformin sustained release have not been conducted. At usual clinical doses and dosing schedules of immediate-release metformin, steady state plasma concentrations of metformin are reached within 24-48 hours and are generally

<1 µg/mL. During controlled clinical trials of immediate-release metformin, maximum metformin plasma levels did not exceed 5 µg/mL, even at maximum doses.

Metabolism

Glimepiride: Glimepiride is completely metabolized by oxidative biotransformation. The major metabolites are the cyclohexyl hydroxy methyl derivative (M1) and the carboxyl derivative (M2). Cytochrome P450 II C9 has been shown to be involved in the biotransformation of glimepiride to M1. M1 is further metabolized to M2 by one or several cytosolic enzymes. M1, but not M2, possesses about 1/3 of the pharmacological activity as compared to its parent; however, whether the glucose lowering effect of M1 is clinically meaningful is not clear.

Metformin sustained-release: Metabolism studies with metformin sustained release have not been conducted. However, intravenous single-dose studies in normal subjects demonstrate that metformin immediate release does not undergo hepatic metabolism or biliary excretion.

Excretion

Glimepiride: When ¹⁴C-glimepiride was given orally, approximately 60% of the total radioactivity was recovered in the urine in 7 days and M1 (predominant) and M2 accounted for 80-90% of that recovered in the urine. Approximately 40% of the total radioactivity was recovered in feces and M1 and M2 (predominant) accounted for about 70% of that recovered in feces. No parent drug was recovered from urine or feces.

GLAVAMET GR (Glimepiride & Metformin Tablets)





Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bplbrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

Metformin: Intravenous single-dose studies in normal subjects demonstrate that metformin is excreted unchanged in the urine and does not undergo hepatic metabolism or biliary excretion. Renal clearance of metformin is approximately 3.5 times greater than creatinine clearance, which indicates that tubular secretion is the major route of metformin elimination. Following oral administration, approximately 90% of the absorbed drug is eliminated via the renal route within the first 24 hours, with a plasma elimination half-life of approximately 6.2 hours. In blood, the elimination half-life is approximately 17.6 hours, suggesting that the erythrocyte mass may be a compartment of distribution.

5.3. Preclinical safety data

Not applicable.

6. Pharmaceutical particulars:

6.1. List of Excipients:

The list of excipients is as follows:

Microcrystalline Cellulose
Crosspovidone
Col. Iron oxide of red
Magnesium Stearate
Maize Starch
Hypromellose Type 2208
Povidone K-30
Isopropyl Alcohol
Purified Water
Colloidal Silicon Dioxide
Titanium Dioxide

6.2. Incompatibilities:

Not applicable

6.3. Shelf life:

24 months

6.4. Special precautions for storage:

Store below 30°C in a dry place.

6.5. Nature and contents of container:

Pack Style: 4×7 Tablets





Bharat Parenterals Limited

Registered Office & Works:
Vill. Haripura, Ta. Savli, Dist. Vadodara - 391520 (Guj.) India.
Tele Fax : (02667)-251679, 251680, 251669, 99099 28332.
E-mail: bplbrd@bplindia.in, info@bplindia.in, Web.: www.bplindia.in
CIN NO: U24231GJ1992PLC018237

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

7 tablets are packed in 1 Alu - PVC Blister. Such 4 Alu - PVC blisters are packed in one monocarton along with patient information leaflet.

6.6. Special precautions for disposal:

No special requirement.

7. Applicant:

Name and Address of Applicant

Evans Therapeutics Limited
24 Abimbola Way,
Isolo Industrail Estate,
Isolo, Lagos, Nigeria.

Name and Address of manufacturer:

M/s. Bharat Parenterals Limited
Survey No. 144 &146, Jarod Samlaya Road,
Village: Haripura, Ta. Savli, Dist. Vadodara – 391520
Gujarat, INDIA.
Tel.91-2667-251680
Fax: 91-2667-251679
E-mail: ra@bplindia.in

