

## Summary of product characteristics

### SILODOSIN CAPSULES 4 mg & 8 mg

#### 1. NAME OF THE MEDICINAL PRODUCT

Silodosin capsules 4 mg & 8 mg

#### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each hard capsule contains silodosin 4 mg , 8 mg.

For the full list of excipients, see section 6.1.

#### 3. PHARMACEUTICAL FORM

Hard gelatin capsule.

4 mg: White to off white color granular powder was filled in size “3” hard gelatin capsules with grey color body and red color cap..

8 mg: White to off white color granular powder was filled in size “2” hard gelatin capsules with golden yellow color body and golden yellow color cap.

#### 4. CLINICAL PARTICULARS

##### 4.1 Therapeutic indications

Treatment of the signs and symptoms of benign prostatic hyperplasia (BPH) in adult men.

##### 4.2 Posology and method of administration

###### Posology

The recommended dose is one capsule of Silodosin 8 mg daily. For special patient populations, one capsule of Silodosin 4 mg daily is recommended (see below).

###### ***Elderly***

No dose adjustment is required in the elderly (see section 5.2).

###### ***Renal impairment***

No dose adjustment is required for patients with mild renal impairment ( $CL_{CR} \geq 50$  to  $\leq 80$  ml/min). A starting dose of 4 mg once daily is recommended in patients with moderate renal impairment ( $CL_{CR} \geq 30$  to  $< 50$  ml/min), which may be increased to 8 mg once daily after one week of treatment, depending on the individual patient's response. The use in patients with severe renal impairment ( $CL_{CR} < 30$  ml/min) is not recommended (see sections 4.4 and 5.2).

### ***Hepatic impairment***

No dose adjustment is required for patients with mild to moderate hepatic impairment. As no data are available, the use in patients with severe hepatic impairment is not recommended (see sections 4.4 and 5.2).

### ***Paediatric population***

There is no relevant use of Silodosin in the paediatric population in the indication.

### **Method of administration**

Oral use.

The capsule should be taken with food, preferably at the same time every day. The capsule should not be broken or chewed but swallowed whole, preferably with a glass of water.

## **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

## **4.4 Special warnings and precautions for use**

### **Intraoperative Floppy Iris Syndrome (IFIS)**

IFIS (a variant of small pupil syndrome) has been observed during cataract surgery in some patients on  $\alpha_1$ -blockers or previously treated with  $\alpha_1$ -blockers. This may lead to increased procedural complications during the operation.

The initiation of therapy with silodosin is not recommended in patients for whom cataract surgery is scheduled. Discontinuing treatment with an  $\alpha_1$ -blocker 1-2 weeks prior to cataract surgery has been recommended, but the benefit and duration of stopping the therapy prior to cataract surgery has not yet been established.

During pre-operative assessment, eye surgeons and ophthalmic teams should consider whether patients scheduled for cataract surgery are being or have been treated with silodosin, in order to ensure that appropriate measures will be in place to manage IFIS during surgery.

### **Orthostatic effects**

The incidence of orthostatic effects with silodosin is very low. However, a reduction in blood pressure can occur in individual patients, leading in rare cases to syncope. At the first signs of orthostatic hypotension (such as postural dizziness), the patient should sit or lie down until the symptoms have disappeared. In patients with orthostatic hypotension, treatment with silodosin is not recommended.

### Renal impairment

The use of silodosin in patients with severe renal impairment ( $CL_{CR} < 30$  ml/min) is not recommended (see sections 4.2 and 5.2).

### Hepatic impairment

Since no data are available in patients with severe hepatic impairment, the use of silodosin in these patients is not recommended (see sections 4.2 and 5.2).

### Carcinoma of the prostate

Since BPH and prostate carcinoma may present the same symptoms and can co-exist, patients thought to have BPH should be examined prior to starting therapy with silodosin, to rule out the presence of carcinoma of the prostate. Digital rectal examination and, when necessary, determination of prostate specific antigen (PSA) should be performed before treatment and at regular intervals afterwards.

Treatment with silodosin leads to a decrease in the amount of semen released during orgasm that may temporarily affect male fertility. This effect disappears after discontinuation of silodosin (see section 4.8).

## **4.5 Interaction with other medicinal products and other forms of interaction**

Silodosin is metabolised extensively, mainly via CYP3A4, alcohol dehydrogenase and UGT2B7. Silodosin is also a substrate for P-glycoprotein. Substances that inhibit (such as ketoconazole, itraconazole, ritonavir or cyclosporine) or induce (such as rifampicin, barbiturates, carbamazepine, phenytoin) these enzymes and transporters may affect the plasma concentrations of silodosin and its active metabolite.

### Alpha-blockers

There is inadequate information about the safe use of silodosin in association with other  $\alpha$ -adrenoreceptor antagonists. Consequently, the concomitant use of other  $\alpha$ -adrenoreceptor antagonists is not recommended.

### CYP3A4 inhibitors

In an interaction study, a 3.7-fold increase in maximum silodosin plasma concentrations and a 3.1-fold increase in silodosin exposure (i.e. AUC) were observed with concurrent administration of a potent CYP3A4 inhibitor (ketoconazole 400 mg). Concomitant use with potent CYP3A4 inhibitors (such as ketoconazole, itraconazole, ritonavir or cyclosporine) is not recommended.

When silodosin was co-administered with a CYP3A4 inhibitor of moderate potency such as diltiazem, an increase in silodosin AUC of approximately 30 % was observed, but  $C_{max}$  and half-life were not affected. This change is clinically not relevant and no dose adjustment is required.

### PDE-5 inhibitors

Minimal pharmacodynamic interactions have been observed between silodosin and maximum doses of sildenafil or tadalafil. In a placebo-controlled study in 24 subjects 45-78 years of age receiving silodosin, the co-administration of sildenafil 100 mg or tadalafil 20 mg induced no clinically meaningful mean decreases in systolic or diastolic blood pressure, as assessed by orthostatic tests (standing *versus* supine). In the subjects over 65 years, the mean decreases at the various time points were between 5 and 15 mmHg (systolic) and 0 and 10 mmHg (diastolic). Positive orthostatic tests were only slightly more common during co-administration; however, no symptomatic orthostasis or dizziness occurred. Patients taking PDE-5 inhibitors concomitantly with silodosin should be monitored for possible adverse reactions.

### Antihypertensives

In the clinical study program, many patients were on concomitant antihypertensive therapy (mostly agents acting on the renin-angiotensin system, beta-blockers, calcium antagonists and diuretics) without experiencing an increase in the incidence of orthostatic hypotension. Nevertheless, caution should be exercised when starting concomitant use with antihypertensives and patients should be monitored for possible adverse reactions.

### Digoxin

Steady state levels of digoxin, a substrate of P-glycoprotein, were not significantly affected by co-administration with silodosin 8 mg once daily. No dose adjustment is required.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy and breast-feeding

Not applicable as silodosin is intended for male patients only.

### Fertility

In clinical studies, the occurrence of ejaculation with reduced or no semen has been observed during treatment with silodosin (see section 4.8), due to the pharmacodynamic properties of silodosin. Before starting treatment, the patient should be informed that this effect may occur, temporarily affecting male fertility.

## **4.7 Effects on ability to drive and use machines**

Silodosin has minor or moderate influence on the ability to drive and use machines. Patients should be informed about the possible occurrence of symptoms related to postural hypotension (such as dizziness) and should be cautioned about driving or operating machines until they know how silodosin will affect them.

## 4.8 Undesirable effects

### Summary of the safety profile

The safety of silodosin has been evaluated in four Phase II-III double-blind controlled clinical studies (with 931 patients receiving silodosin 8 mg once daily and 733 patients receiving placebo) and in two long-term open-label extension phase studies. In total, 1,581 patients have received silodosin at a dose of 8 mg once daily, including 961 patients exposed for at least 6 months and 384 patients exposed for 1 year. The most frequent adverse reactions reported with silodosin in placebo controlled clinical studies and during long-term use were ejaculatory disorders such as retrograde ejaculation and anejaculation (ejaculatory volume reduced or absent), with a frequency of 23 %. This may temporarily affect male fertility. It is reversible within a few days upon discontinuation of treatment (see section 4.4).

### Tabulated list of adverse reactions

In the table below, adverse reactions reported in all clinical studies and in the worldwide post-marketing experience for which a reasonable causal relationship exists are listed by MedDRA system organ class and frequency: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ), not known (cannot be estimated from available data). Within each frequency grouping the observed adverse reactions are presented in order of decreasing seriousness.

	Very common	Common	Uncommon	Rare	Very rare	Not known
Immune system disorders					Allergic-type reactions including facial swelling, swollen tongue and pharyngeal oedema <sup>1</sup>	
Psychiatric disorders			Libido decreased			
Nervous system disorders		Dizziness		Syncope Loss of consciousness <sup>1</sup>		

Cardiac disorders			Tachycardia <sup>1</sup>	Palpitations <sup>1</sup>		
Vascular disorders		Orthostatic hypotension	Hypotension <sup>1</sup>			
Respiratory, thoracic and mediastinal disorders		Nasal congestion				
Gastrointestinal disorders		Diarrhoea	Nausea Dry mouth			
Hepatobiliary disorders			Abnormal liver function tests <sup>1</sup>			
Skin and subcutaneous tissue disorders			Skin rash <sup>1</sup> , Pruritus <sup>1</sup> Urticaria <sup>1</sup> Drug eruption <sup>1</sup>			
Reproductive system and breast disorders	Ejaculatory disorders, including retrograde ejaculation, anejaculation		Erectile dysfunction			
Injury, poisoning and procedural complication						Intraoperative Floppy Iris Syndrome

<sup>1</sup> - adverse reactions from spontaneous reporting in the worldwide post-marketing experience (frequencies calculated from events reported in Phase I-IV clinical trials and non-interventional studies).

#### Description of selected adverse reactions

##### *Orthostatic hypotension*

The incidence of orthostatic hypotension in placebo-controlled clinical studies was 1.2 % with silodosin and 1.0 % with placebo. Orthostatic hypotension may occasionally lead to syncope (see section 4.4).

##### *Intraoperative Floppy Iris Syndrome (IFIS)*

IFIS has been reported during cataract surgery (see section 4.4).

## 4.9 Overdose

Silodosin was evaluated at doses of up to 48 mg/day in healthy male subjects. The dose-limiting adverse reaction was postural hypotension. If ingestion is recent, induction of vomiting or gastric lavage may be considered. Should overdose of silodosin lead to hypotension, cardiovascular support has to be provided. Dialysis is unlikely to be of significant benefit since silodosin is highly (96.6 %) protein bound.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Urologicals, alpha-adrenoreceptor antagonists, ATC code: G04CA04.

#### Mechanism of action

Silodosin is highly selective for  $\alpha_{1A}$ -adrenoreceptors that are primarily located in the human prostate, bladder base, bladder neck, prostatic capsule and prostatic urethra. Blockade of these  $\alpha_{1A}$ -adrenoreceptors causes smooth muscle in these tissues to relax, thus decreasing bladder outlet resistance, without affecting detrusor smooth muscle contractility. This causes an improvement of both storage (irritative) and voiding (obstructive) symptoms (Lower urinary tract symptoms, LUTS) associated with benign prostatic hyperplasia.

Silodosin has a substantially lower affinity for the  $\alpha_{1B}$ -adrenoreceptors that are primarily located in the cardiovascular system. It has been demonstrated *in vitro* that the  $\alpha_{1A}:\alpha_{1B}$  binding ratio of silodosin (162:1) is extremely high.

#### Clinical efficacy and safety

In a Phase II dose-finding, double-blind, placebo-controlled clinical study with silodosin 4 or 8 mg once daily, a greater improvement in American Urologic Association (AUA) symptom index score was observed with silodosin 8 mg ( $-6.8 \pm 5.8$ ,  $n=90$ ;  $p=0.0018$ ) and silodosin 4 mg ( $-5.7 \pm 5.5$ ,  $n=88$ ;  $p=0.0355$ ) as compared to placebo ( $-4.0 \pm 5.5$ ,  $n=83$ ).

Over 800 patients with moderate to severe symptoms of BPH (International Prostate Symptom Score, IPSS, baseline value  $\geq 13$ ) received silodosin 8 mg once daily in two Phase III placebo-controlled clinical studies conducted in the United States and in one placebo- and active-controlled clinical study conducted in Europe. In all studies, patients who did not respond to placebo during a 4-week placebo run-in phase were randomised to receive the study treatment. In

all studies, patients treated with silodosin had a greater decrease in both storage (irritative) and voiding (obstructive) symptoms of BPH as compared to placebo as assessed after 12 weeks of treatment. Data observed in the Intent-to-treat populations of each study are shown below:

Study	Treatment arm	No. of patients	IPSS Total score			IPSS Irritative symptoms		IPSS Obstructive symptoms	
			Baseline value (±SD)	Change from baseline	Difference (95 % CI) vs placebo	Change from baseline	Difference (95 % CI) vs placebo	Change from baseline	Difference (95 % CI) vs placebo
US-1	Silodosin	233	22 ± 5	-6.5	-2.8*	-2.3	-0.9*	-4.2	-1.9*
	Placebo	228	21 ± 5	-3.6	(-3.9, -1.7)	-1.4	(-1.4, -0.4)	-2.2	(-2.6, -1.2)
US-2	Silodosin	233	21 ± 5	-6.3	-2.9*	-2.4	-1.0*	-3.9	-1.8*
	Placebo	229	21 ± 5	-3.4	(-4.0, -1.8)	-1.3	(-1.5, -0.6)	-2.1	(-2.5, -1.1)
Europe	Silodosin	371	19 ± 4	-7.0	-2.3*	-2.5	-0.7°	-4.5	-1.7*
	Tamsulosin	376	19 ± 4	-6.7	(-3.2, -1.4)	-2.4	(-1.1, -0.2)	-4.2	(-2.2, -1.1)
	Placebo	185	19 ± 4	-4.7	-2.0*	-1.8	(-1.1, -0.2)	-2.9	(-2.0, -0.8)

\* p<0.001 vs Placebo; ° p =0.002 vs Placebo

In the active-controlled clinical study conducted in Europe, silodosin 8 mg once daily was shown to be non inferior to tamsulosin 0.4 mg once daily: the adjusted mean difference (95 % CI) in the IPSS Total Score between treatments in the per-protocol population was 0.4 (-0.4 to 1.1). The responder rate (i.e. improvement in the IPSS total score by at least 25 %) was significantly higher in the silodosin (68 %) and tamsulosin group (65 %), as compared to placebo (53 %).

In the long-term open-label extension phase of these controlled studies, in which patients received silodosin for up to 1 year, the symptom improvement induced by silodosin at week 12 of treatment was maintained over 1 year.

In a Phase IV clinical trial performed in Europe, with a mean baseline IPSS total score of 18.9 points, 77.1 % were responders to silodosin (as assessed by a change from baseline in the IPSS total score of at least 25 %). Approximately half of the patients reported an improvement in the most bothersome symptoms complained at baseline by the patients (i.e. nocturia, frequency, decreased stream, urgency, terminal dribbling and incomplete emptying), as assessed by the ICS-male



questionnaire.

No significant reduction in supine blood pressure was observed in all clinical studies conducted with silodosin.

Silodosin 8 mg and 24 mg daily had no statistically significant effect on ECG intervals or cardiac repolarisation relative to placebo.

#### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with Silodosin in all subsets of the paediatric population in BPH (see section 4.2 for information on paediatric use).

### **5.2 Pharmacokinetic properties**

The pharmacokinetics of silodosin and its main metabolites have been evaluated in adult male subjects with and without BPH after single and multiple administrations with doses ranging from 0.1 mg to 48 mg per day. The pharmacokinetics of silodosin is linear throughout this dose range.

The exposure to the main metabolite in plasma, silodosin glucuronide (KMD-3213G), at steady-state is about 3-fold that of the parent substance. Silodosin and its glucuronide reach steady-state after 3 days and 5 days of treatment, respectively.

#### Absorption

Silodosin administered orally is well absorbed and absorption is dose proportional. The absolute bioavailability is approximately 32 %. An *in vitro* study with Caco-2 cells showed that silodosin is a substrate for P-glycoprotein. Food decreases  $C_{\max}$  by approximately 30 %, increases  $t_{\max}$  by approximately 1 hour and has little effect on AUC.

In healthy male subjects of the target age range (n=16, mean age 55±8 years) after once-a-day oral administration of 8 mg immediately after breakfast for 7 days, the following pharmacokinetic parameters were obtained:  $C_{\max}$  87±51 ng/ml (sd),  $t_{\max}$  2.5 hours (range 1.0-3.0), AUC 433±286 ng • h/ml.

#### Distribution

Silodosin has a volume of distribution of 0.81 l/kg and is 96.6 % bound to plasma proteins. It does not distribute into blood cells.

Protein binding of silodosin glucuronide is 91 %.

#### Biotransformation

Silodosin undergoes extensive metabolism through glucuronidation (UGT2B7),

alcohol and aldehyde dehydrogenase and oxidative pathways, mainly CYP3A4. The main metabolite in plasma, the glucuronide conjugate of silodosin (KMD-3213G), that has been shown to be active *in vitro*, has an extended half-life (approximately 24 hours) and reaches plasma concentrations approximately four times higher than those of silodosin. *In vitro* data indicate that silodosin does not have the potential to inhibit or induce cytochrome P450 enzyme systems.

### Elimination

Following oral administration of <sup>14</sup>C-labelled silodosin, the recovery of radioactivity after 7 days was approximately 33.5 % in urine and 54.9 % in faeces. Body clearance of silodosin was approximately 0.28 l/h/kg. Silodosin is excreted mainly as metabolites, very low amounts of unchanged drug are recovered in urine. The terminal half-life of parent drug and its glucuronide is approximately 11 hours and 18 hours, respectively.

### Special populations

#### ***Elderly***

Exposure to silodosin and its main metabolites does not change significantly with age, even in subjects of age over 75 years.

#### ***Paediatric population***

Silodosin has not been evaluated in patients less than 18 years of age.

#### ***Hepatic impairment***

In a single-dose study, the pharmacokinetics of silodosin was not altered in nine patients with moderate hepatic impairment (Child-Pugh scores 7 to 9), compared to nine healthy subjects. Results from this study should be interpreted with caution, since enrolled patients had normal biochemistry values, indicating normal metabolic function, and they were classified as having moderate liver impairment based on ascites and hepatic encephalopathy.

The pharmacokinetics of silodosin in patients with severe hepatic impairment has not been studied.

#### ***Renal impairment***

In a single-dose study, exposure to silodosin (unbound) in subjects with mild (n=8) and moderate renal impairment (n=8) resulted, on average, in an increase of C<sub>max</sub> (1.6-fold) and AUC (1.7-fold) relative to subjects with normal renal function (n=8). In subjects with severe renal impairment (n=5) increase of exposure was 2.2-fold for C<sub>max</sub> and 3.7-fold for AUC. Exposure to the main metabolites, silodosin glucuronide

and KMD3293, was also increased.

Plasma level monitoring in a Phase III clinical study showed that levels of total silodosin after 4 weeks of treatment did not change in patients with mild impairment (n=70), compared to patients with normal renal function (n=155), while the levels were doubled on average in patients with moderate impairment (n=7).

A review of safety data of patients enrolled in all clinical studies does not indicate that mild renal impairment (n=487) poses an additional safety risk during silodosin therapy (such as an increase in dizziness or orthostatic hypotension) as compared to patients with normal renal function (n=955). Accordingly, no dose adjustment is required in patients with mild renal impairment. Since only limited experience exists in patients with moderate renal impairment (n=35), a lower starting dose of 4 mg is recommended. In patients with severe renal impairment administration of Silodosin is not recommended.

### **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, carcinogenic, mutagenic and teratogenic potential. Effects in animals (affecting the thyroid gland in rodents) were observed only at exposures considered sufficiently in excess of the maximum human exposure, indicating little relevance to clinical use.

In male rats, decreased fertility was observed from exposures which were approximately twice the exposure at the maximum recommended human dose. The observed effect was reversible.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Capsule contents

Mannitol , Pregelatinised starch , Sodium lauryl sulphate , Magnesium Stearate

#### Capsule shell contents

**4 mg:** size “3” hard gelatin capsules with grey color body and red color cap.

**Composition:** Gelatin, Methylparaben, Propylparaben, Sodium Lauryl Sulphate, Purified Water, Brilliant Blue, Carmoisine, Tartrazine, Titanium dioxide.

**8 mg:** size “2” hard gelatin capsules with golden yellow color body and golden yellow color cap.

**Composition:** Gelatin ,Methyl Paraben, Propyl Paraben, Sodium Lauryl Sulfate, Purified Water, Sunset yellow, Tartrazine, Titanium Dioxide.

**6.2 Incompatibilities**

Not applicable.

**6.3 Shelf life**

2 years.

**6.4 Special precautions for storage**

Do not store above 30°C.

**6.5 Nature and contents of container**

10's Alu – Alu strip pack.

**6.6 Special precautions for disposal**

No special requirements.

**7. Manufactured by:**


**MSN LABORATORIES PRIVATE LIMITED,**

(Formulations Division),

Plot No: 42, Anrich Industrial Estate,

IDA, Bollaram, Sangareddy District,

Telangana, India.

<b>MSN LABORATORIES PRIVATE LIMITED-FORMULATIONS DIVISION</b>	
<b>SILODOSIN CAPSULES 4 mg</b>	

### **1.3.2 Labelling (outer & inner labels)**

The artworks of container label, carton & Pack Insert for SILOTRIF 4 (Silodosin Capsules 4 mg) are enclosed in the following pages.

# Harmonization

Botswana, Nigeria, Rwanda, Ethiopia, Zambia,  
Malawi, Tanzania, Uganda, Burundi, Mauritius, Madagascar, Namibia, IVC.

## R<sub>x</sub> Silodosin Capsules 4 mg/ Silodosine 4 mg - Gélules SILOTRIF 4

Each hard gelatin capsule contains: Silodosin 4 mg/  
Chaque gélule de gélatine dure contient: Silodosine 4 mg  
Dosage: As directed by the Physician /  
Posologie: Conformément à la prescription médicale.  
Lire attentivement la notice.  
Do not store above 30°C /  
A conserver à une température ne dépassant pas 30°C.  
Protect from light and moisture /  
à l'abri de la lumière et de l'humidité.  
Keep out of reach of children /  
Tenir hors de la vue et de la portée des enfants

B12958-00

Manufactured by / Fabriqué par :  
**MSN Laboratories Private Limited**  
(Formulations Division),  
Plot No. 42, Anrich Industrial Estate,  
Bollaram, Sangareddy District - 502 325,  
Telangana, INDIA.

Mfg. Lic. No. / No. Lic. de Fabrication:  
38/MD/AP/2007/F/CC

RESPECTER LES DOSES PRESCRITES

Liste I : Uniquement sur ordonnance

## R<sub>x</sub> Silodosin Capsules 4 mg/ Silodosine 4 mg - Gélules SILOTRIF 4

Each hard gelatin capsule contains: Silodosin 4 mg/  
Chaque gélule de gélatine dure contient: Silodosine 4 mg  
Dosage: As directed by the Physician /  
Posologie: Conformément à la prescription médicale.  
Lire attentivement la notice.  
Do not store above 30°C /  
A conserver à une température ne dépassant pas 30°C.  
Protect from light and moisture /  
à l'abri de la lumière et de l'humidité.  
Keep out of reach of children /  
Tenir hors de la vue et de la portée des enfants

B12958-00

Manufactured by / Fabriqué par :  
**MSN Laboratories Private Limited**  
(Formulations Division),  
Plot No. 42, Anrich Industrial Estate,  
Bollaram, Sangareddy District - 502 325,  
Telangana, INDIA.

Mfg. Lic. No. / No. Lic. de Fabrication:  
38/MD/AP/2007/F/CC

RESPECTER LES DOSES PRESCRITES

Liste I : Uniquement sur ordonnance

## R<sub>x</sub> Silodosin Capsules 4 mg/ Silodosine 4 mg - Gélules SILOTRIF 4

Each hard gelatin capsule contains: Silodosin 4 mg/  
Chaque gélule de gélatine dure contient: Silodosine 4 mg  
Dosage: As directed by the Physician /  
Posologie: Conformément à la prescription médicale.  
Lire attentivement la notice.  
Do not store above 30°C /  
A conserver à une température ne dépassant pas 30°C.  
Protect from light and moisture /  
à l'abri de la lumière et de l'humidité.  
Keep out of reach of children /  
Tenir hors de la vue et de la portée des enfants

B12958-00

Manufactured by / Fabriqué par :  
**MSN Laboratories Private Limited**  
(Formulations Division),  
Plot No. 42, Anrich Industrial Estate,  
Bollaram, Sangareddy District - 502 325,  
Telangana, INDIA.

Mfg. Lic. No. / No. Lic. de Fabrication:  
38/MD/AP/2007/F/CC

RESPECTER LES DOSES PRESCRITES

Liste I : Uniquement sur ordonnance

## R<sub>x</sub> Silodosin Capsules 4 mg/ Silodosine 4 mg - Gélules SILOTRIF 4

Each hard gelatin capsule contains: Silodosin 4 mg/  
Chaque gélule de gélatine dure contient: Silodosine 4 mg  
Dosage: As directed by the Physician /  
Posologie: Conformément à la prescription médicale.  
Lire attentivement la notice.  
Do not store above 30°C /  
A conserver à une température ne dépassant pas 30°C.  
Protect from light and moisture /  
à l'abri de la lumière et de l'humidité.  
Keep out of reach of children /  
Tenir hors de la vue et de la portée des enfants

B12958-00

Manufactured by / Fabriqué par :  
**MSN Laboratories Private Limited**  
(Formulations Division),  
Plot No. 42, Anrich Industrial Estate,  
Bollaram, Sangareddy District - 502 325,  
Telangana, INDIA.



Mfg. Lic. No. / No. Lic. de Fabrication:  
38/MD/AP/2007/F/CC

RESPECTER LES DOSES PRESCRITES


Liste I : Uniquement sur ordonnance

Foil width 262 mm



Strip Size : 130 x 80 mm  
Foil width : 262 mm

MSN LABORATORIES PRIVATE LIMITED		PACKAGING DEVELOPMENT	
Artwork Information		Specification for Printed Foil	
		Test	Specification
Brand Name	Silotrif 4	Description	Printed Soft tempered aluminium foil with LDPE coating on the sealing side.
Generic Name	Silodosin Capsules 4 mg		
Pack Style	10's	Thickness of Aluminium	0.037 to 0.043 mm
Foil Width	262 mm	Width	262 ± 1.0 mm (261 – 263 mm)
Item Code	B12958-00	Aluminium foil GSM	108 ± 8% (99.36 - 116.64) GSM
Supersede Code	NA	VMCH Coating GSM	35 ± 8% (32.20 - 37.80) GSM
Change Part No:	–	NC Coating	NA
Version	00	Pin Holes	Nil
Date & Time	23-08-2023 / 1:48pm	Ink Adhesion Test	No Ink Lifting
Country	Harmonization Botswana, Nigeria, Rwanda, Ethiopia, Zambia, Malawi, Tanzania, Uganda, Burundi, Mauritius, Madagascar, Namibia, IVC.	Inner Core Diameter	76 ± 1 mm
Customer	NA	PRC/Non-PRC	Non-PRC (Continues text)
Font Type	Arial	Eye Mark Size	NA
Font Size (min.)	5pt	Perforation/Non-Perforation	Non-Perforation
Developed by	Sridhar	Colours	 BLACK  485 C
Reviewed by	Srilakshmi		







Harmonization  
Botswana, Nigeria, Rwanda, Ethiopia, Zambia,  
Malawi, Tanzania, Uganda, Burundi, Mauritius,  
Madagascar, Namibia, IVC.




- Braille Text embossed not to be print





Silodosin  
Capsules  
#4 mg

 LABORATORIES PRIVATE LIMITED		PACKAGING DEVELOPMENT	
Artwork Information		Specification for Printed Carton	
		Test	Specification
Brand Name	SILOTRIF 4	Substrate	Cyber XL board with reverse tuck in type
Generic Name	Silodosin Capsules 4 mg	GSM	300 ± 5%
Pack Style	10's Pack	Varnish/Lamination	UV Varnish except over printing area
Dimensions	135 x 18 x 85 mm (LxWxH)	Mode of supply	Bundles 25's, 50's & 100's
Item Code	B23292-00	Pharmacode	4301
Supersede Code	NA	Font Type	Arial
Version	00	Font Size (min.)	7pt
Date & Time	23-08-2023 / 1:57pm	Reviewed by	Srilakshmi
Country	Harmonization Botswana, Nigeria, Rwanda, Ethiopia, Zambia, Malawi, Tanzania, Uganda, Burundi, Mauritius, Madagascar, Namibia, IVC.	Colours	 PANTONE 2935C  PANTONE 485 C
Customer	—		 PANTONE 273 C  Black
Developed by	Sridhar		 PANTONE 122 C

# Harmonization



- Braille Text embossed not to be print



MSN LABORATORIES PRIVATE LIMITED		PACKAGING DEVELOPMENT	
Artwork Information		Specification for Printed Carton	
		Test	Specification
Brand Name	SILOTRIF 8	Substrate	Cyber XL board with reverse tuck in type
Generic Name	Silodosin Capsules 8 mg	GSM	300 ± 5%
Pack Style	10's Pack	Varnish/Lamination	UV Varnish except over printing area
Dimensions	150 x 18 x 97 mm (LxWxH)	Mode of supply	Bundles 25's, 50's & 100's
Item Code	B23293-00	Pharmacode	4302
Supersede Code	NA	Font Type	Arial
Version	00	Font Size (min.)	7pt
Date & Time	12-04-2023 / 12:41pm	Reviewed by	Srilakshmi
Country	Harmonization	Colours	<div><div></div> PANTONE 313 C</div> <div><div></div> PANTONE 485 C</div>
Customer	—		<div><div></div> PANTONE 273 C</div> <div><div></div> Black</div>
Developed by	Sridhar		<div><div></div> PANTONE 122 C</div>

Silodosin  
Capsules  
#8 mg