

## SUMMARY OF PRODUCT CHARACTERISTIC:

### 1. NAME OF THE MEDICINAL PRODUCT:

Loperamide Hydrochloride Capsules USP

### 2. Qualitative and Quantitative Composition

#### Composition:

Each Hard Gelatin Capsule Contains:

Loperamide Hydrochloride USP..... 2 mg

Excipients.....Q.S.

Colour: Approved Colours used in Capsules shells

For the full list of excipients, see section 6.1

### 3. Pharmaceutical Form

Solid Dosage form (Hard Gelatin, Capsules)

Loperamide Hydrochloride Capsules USP are available as, yellow coloured unsealed Hard gelatin capsules, size “4” containing almost white powder.

### 4. Clinical Particulars

#### 4.1 Therapeutic indications

For symptomatic treatment of acute diarrhoea in adults and children aged 12 years and over.

#### 4.2 Posology and method of administration

##### Posology

##### Adults and children over 12 years of age:

Two capsules to be taken initially, followed by one capsule after each loose motion, up to a maximum of six capsules in any 24 hours.

##### Children under 12 years of age:

Not recommended

##### Use in Elderly

No dose adjustment is required for the elderly.

##### Renal impairment

No dose adjustment is required for patients with renal impairment.

##### Hepatic impairment

Although no pharmacokinetic data are available in patients with hepatic impairment Loperamide should be used with caution in such patients because of reduced first pass metabolism.

## **Method of administration**

Oral use.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipient listed in section 6.1.

When inhibition of peristalsis is to be avoided due to the possible risk of significant sequelae including ileus, megacolon, toxic megacolon and certain poisonings in particular:

- Children less than 12 years of age.
- When ileus or constipation are present or when abdominal distension develops
- In patients with acute ulcerative colitis.
- In patients with bacterial enterocolitis caused by invasive organisms including Salmonella, Shigella, and Campylobacter.
- In patients with pseudomembranous colitis associated with the use of broad-spectrum antibiotics.

Loperamide hydrochloride should not be used alone in acute dysentery, which is characterised by blood in stools and elevated body temperatures.

### **4.4 Special warnings and precautions for use**

Treatment of diarrhoea with loperamide is only symptomatic. Whenever an underlying etiology can be determined, specific treatment should be given when appropriate.

The priority in acute diarrhoea is the prevention or reversal of fluid and electrolyte depletion. This is particularly important in young children and in frail and elderly patients with acute diarrhoea. Use of loperamide hydrochloride does not preclude the administration of appropriate fluid and electrolyte replacement therapy.

Since persistent diarrhoea can be an indicator of potentially more serious conditions, loperamide hydrochloride should not be used for prolonged periods until the underlying cause of the diarrhoea has been investigated.

Loperamide hydrochloride must be used with caution when the hepatic function necessary for the drug's metabolism is defective (eg in cases of severe hepatic disturbance), as this might result in a relative overdose leading to CNS toxicity

Patients with AIDS treated with loperamide hydrochloride for diarrhoea should have therapy stopped at the earliest signs of abdominal distension. There have been isolated reports of toxic megacolon in AIDS patients with infectious colitis from both viral and bacterial pathogens treated with loperamide hydrochloride.

When no clinical change is observed in the acute diarrhoea within 48 hours, the administration of loperamide must be interrupted and the patient must be advised to consult his doctor.

Treatment with Loperamide must be interrupted immediately when obstipation, abdominal distension or subileus develops

Cardiac events including QT prolongation and torsades de pointes have been reported in association with overdose. Some cases had a fatal outcome. Overdose can unmask existing Brugada syndrome. Patients should not exceed the recommended dose and/or the recommended duration of treatment.

Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine because it contains lactose

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Non-clinical data have shown that loperamide is a P-glycoprotein substrate. Furthermore, loperamide is mainly metabolised by CYP3A4 and CYP2C8. Concomitant administration of loperamide (16 mg single dose) with quinidine, or ritonavir, which are both P-glycoprotein inhibitors, resulted in a 2 to 3-fold increase in loperamide plasma levels.

The results of one published pharmacokinetic study suggested that the concomitant administration of loperamide with oral desmopressin may result in a 3-fold increase of desmopressin plasma concentrations although no clinical effects were reported.

Possible interactions may occur with drugs that delay intestinal peristalsis (for instance anticholinergic drugs) because the effects of loperamide could be enhanced.

Administration of itraconazole with loperamide (4 mg single dose) increased loperamide plasma levels 3- to 4-fold. In addition, gemfibrozil, a CYP2C8 inhibitor, increased the AUC of loperamide 2-fold. Concomitant use of itraconazole and gemfibrozil with loperamide raised the mean C<sub>max</sub> and AUC of loperamide about 2- and 13-fold, respectively. This increase did not lead to measurable CNS effects.

The concomitant administration of loperamide (16mg single dose) and ketoconazole, an inhibitor of CYP3A4 and P-glycoprotein, resulted in a 5-fold increase in loperamide plasma concentrations. This increase was not associated with increased pharmacodynamic effects as measured by pupillometry.

The clinical relevance of these pharmacokinetic interactions, when loperamide is given at recommended dosages (2 mg, up to 12 mg maximum daily dose), is unknown.

#### **4.6 Fertility, pregnancy and lactation**

## **Pregnancy**

A limited amount of data from the use of loperamide in pregnant women is available. In one of two epidemiological studies the use of loperamide during early pregnancy suggested a possible moderate increased risk for hypospadias, however, an increased risk for major malformations could not be identified. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. Safety in human pregnancy has not been established, although from animal studies there are no indications that loperamide HCl possesses any teratogenic or embryotoxic properties.

If possible the use of loperamide should be avoided during the first trimester of pregnancy, however, it may be used during the second and third trimester of pregnancy

## **Breast-feeding**

Small amounts of loperamide may appear in human breast milk. Therefore, this medicine is not recommended during breast-feeding. Women who are pregnant or breast feeding infants should therefore be advised to consult their doctor for appropriate treatment.

## **Fertility**

Only high doses of loperamide hydrochloride affected female fertility in non-clinical studies.

### **4.7 Effects on ability to drive and use machines**

Loperamide hydrochloride has moderate influence on the ability to drive and use machines. Loss of consciousness, depressed level of consciousness, tiredness, dizziness or drowsiness may occur when diarrhoea is treated with loperamide hydrochloride.

Therefore, it is advisable to use caution when driving or operating machinery.

### **4.8 Undesirable effects**

#### **Adults and children aged $\geq 12$ years**

The safety of loperamide hydrochloride was evaluated in 2755 adults and children aged  $\geq 12$  years who participated in 26 controlled and uncontrolled clinical trials of loperamide hydrochloride used for the treatment of acute diarrhoea.

The most commonly reported (i.e.  $\geq 1\%$  incidence) adverse drug reactions (ADRs) in clinical trials with loperamide hydrochloride in acute diarrhoea were: constipation (2.7%), flatulence (1.7%), headache (1.2%) and nausea (1.1%).

Table 1 displays ADRs that have been reported with the use of loperamide hydrochloride from either clinical trial (acute diarrhoea) or post-marketing experience.

The frequency categories use the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$ )

to <1/10); uncommon ( $\geq 1/1,000$  to <1/100); rare ( $\geq 1/10,000$  to <1/1,000); and very rare (<1/10,000).

**Table 1 Adverse Drug reactions**

System Organ Class	Indication		
	Common	Uncommon	Rare
Immune System Disorders			Hypersensitivity reaction <sup>a</sup> Anaphylactic reaction (including Anaphylactic shock) <sup>a</sup> Anaphylactoid reaction <sup>a</sup>
Nervous System Disorders	Headache Dizziness	Somnolence <sup>a</sup>	Loss of consciousness <sup>a</sup> Stupor <sup>a</sup> Depressed level of consciousness <sup>a</sup> Hypertonia <sup>a</sup> Coordination abnormality <sup>a</sup>
Eye Disorders			Miosis <sup>a</sup>
Gastrointestinal Disorders	Constipation Nausea Flatulence	Abdominal pain Abdominal discomfort Dry mouth Abdominal pain upper Vomiting Dyspepsia <sup>a</sup>	Ileus <sup>a</sup> (including paralytic ileus) Megacolon <sup>a</sup> (including toxic megacolon <sup>b</sup> ) Glossodynia <sup>a</sup> Abdominal distension
Skin and Subcutaneous Tissue Disorders		Rash	Bullous eruption <sup>a</sup> (including Stevens-Johnson syndrome, toxic epidermal necrolysis and erythema multiforme) Angioedema <sup>a</sup> Urticaria <sup>a</sup> Pruritus <sup>a</sup>
Renal and Urinary Disorders			Urinary retention <sup>a</sup>
General Disorders and Administration Site Conditions			Fatigue <sup>a</sup>

a: Inclusion of this term is based on post-marketing reports for loperamide hydrochloride. As the process for determining post marketing ADRs did not differentiate between chronic and acute indications or adults and children, the frequency is estimated from all clinical trials with

loperamide hydrochloride (acute and chronic), including trials in children  $\leq 12$  years (N=3683).

b: See section Special Warnings and Special Precautions for use.

### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).

## **4.9 Overdose**

### **Symptoms**

In case of overdose (including relative overdose due to hepatic dysfunction), CNS depression (stupor, coordination abnormality, somnolence, miosis, muscular hypertonia, and respiratory depression), urinary retention and ileus may occur. Children may be more sensitive to CNS effects than adults.

In individuals who have ingested overdoses of loperamide HCl, cardiac events such as QT interval prolongation, torsades de pointes, other serious ventricular arrhythmias, cardiac arrest and syncope have been observed. Fatal cases have also been reported. Overdose can unmask existing Brugada syndrome.

### **Treatment**

If symptoms of overdose occur, naloxone can be given as an antidote. Since the duration of action of loperamide is longer than that of naloxone (1 to 3 hours), repeated treatment with naloxone might be indicated. Therefore, the patient should be monitored closely for at least 48 hours in order to detect possible CNS depression.

## **5. Pharmacological properties**

### **5.1 Pharmacodynamics properties**

Pharmacotherapeutic Group: Antipropulsives;

ATC code: A07DA03

Loperamide hydrochloride is a synthetic opioid which inhibits gut motility by binding to opiate receptors in the gut wall and may also reduce gastrointestinal secretions, resulting in improvement in diarrhoea symptoms.

Loperamide also increases the tone of the anal sphincter. Onset of antidiarrhoeal effect occurred as soon as one hour after intake of a 4 mg dose of loperamide.

In a double blind randomised clinical trial in 56 patients with acute diarrhoea receiving loperamide, onset of anti-diarrhoeal action was observed within one hour following a single 4 mg dose. Clinical comparisons with other antidiarrhoeal drugs confirmed this exceptionally rapid onset of action of loperamide.

## **5.2 Pharmacokinetic properties**

**Pharmacotherapeutic Group:** Antipropulsives;

**ATC code:** A07DA03

Loperamide hydrochloride is a synthetic opioid which inhibits gut motility by binding to opiate receptors in the gut wall and may also reduce gastrointestinal secretions, resulting in improvement in diarrhoea symptoms.

Loperamide also increases the tone of the anal sphincter. Onset of antidiarrhoeal effect occurred as soon as one hour after intake of a 4 mg dose of loperamide.

In a double blind randomised clinical trial in 56 patients with acute diarrhoea receiving loperamide, onset of anti-diarrhoeal action was observed within one hour following a single 4 mg dose. Clinical comparisons with other antidiarrhoeal drugs confirmed this exceptionally rapid onset of action of loperamide.

## **5.3 Preclinical safety data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development.

Acute and chronic studies on loperamide showed no specific toxicity.

Loperamide had no effect on fertility in male rats when administered orally prior to mating at doses up to approximately 40 mg/kg. No pregnancy occurred in females dosed with approximately 40 mg/kg. Lower doses (approximately 10 and 2.5mg/kg) did not affect female fertility. In rabbits no differences in pregnancy rate were observed when females were administered orally up to 40mg/kg.

No malformations of offspring were noted in rats and rabbits dosed up to 40 mg/kg. Loperamide did not show genotoxic potential.

In an 18-month carcinogenicity study in rats, with doses up to 100 times the maximum human dose no evidence of carcinogenesis was found.

Non-clinical in vitro and in vivo evaluation of loperamide indicates no significant cardiac electrophysiological effects within its therapeutically relevant concentration range and at

significant multiples of this range (up to 47-fold. However, at extremely high concentrations associated with overdoses, loperamide has cardiac electrophysiological actions consisting of inhibition of potassium (hERG) and sodium currents, and arrhythmias.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Maize Starch	BP
Maize Starch (For Paste)	BP
Lactose Monohydrate	BP
Sodium Methyl Hydroxybenzoate	BP
Sodium Propyl Hydroxybenzoate	BP
Povidone	BP
Purified Water	BP
Purified Talc	BP
Magnesium Stearate	BP
Empty Hard Gelatin Capsules Yellow Colour Size - “4”	IH



## **6.2 Incompatibilities**

Not Applicable

## **6.3 Shelf life**

24 Months

## **6.4 Special precaution for storage**

Store at temperature should not exceed 30°C.

## **6.5 Nature and contents of the container**

### **Primary Packing materials:**

- Printed Alu foil
- PVC foil

### **Secondary Packaging Material:**

- Printed Carton
- Pack Insert

## **6.6 Special precautions for disposal and other handling**

No special requirements for disposal.

## **7. Marketing authorisation holder**

McW Healthcare Private Limited

286, 287A, 287B, Sector E, Industrial Area,

Sanwer Road, Indore (M.P.), INDIA

Website: - [www.mcwhealthcare.com](http://www.mcwhealthcare.com)

## **8. Marketing authorisation number(s)**

XXXX

## **9. Date of first authorisation/renewal of the authorisation**

Jan 2022

## **10. Date of revision of the text**

XXXX