

Summary of Product Characteristics for

GEMBIN
GEMCITABINE FOR INJECTION USP 200mg/vial, 1g/vial & 1.4g/vial

1. Name of the medicinal product

Gemcitabine for Injection USP 200mg/vial, 1g/vial & 1.4g/vial

2. Qualitative and quantitative composition

Gemcitabine for Injection USP (Lyophilized)

Each Vial Contains:

Gemcitabine Hydrochloride USP

Equivalent to Gemcitabine200 mg

Suitably Buffered

Gemcitabine for Injection USP (Lyophilized)

Each Vial Contains:

Gemcitabine Hydrochloride USP

Equivalent to Gemcitabine1 gm

Suitably Buffered

Gemcitabine for Injection USP (Lyophilized)

Gemcitabine Hydrochloride USP

Equivalent to Gemcitabine1.4 gm

Suitably Buffered

3. Pharmaceutical form

A white to off white lyophilized mass

4. Clinical particulars

4.1 Therapeutic indications

Gemcitabine is indicated for the treatment of locally advanced or metastatic bladder cancer in combination with cisplatin.

Gemcitabine is indicated for treatment of patients with locally advanced or metastatic adenocarcinoma of the pancreas.

Gemcitabine, in combination with cisplatin is indicated as first line treatment of patients with locally advanced or metastatic non-small cell lung cancer (NSCLC). Gemcitabine monotherapy can be considered in elderly patients or those with performance status 2.

Gemcitabine is indicated for the treatment of patients with locally advanced or metastatic epithelial ovarian carcinoma, in combination with carboplatin, in patients with relapsed disease following a recurrence-free interval of at least 6 months after platinum-based, first-line therapy.

Gemcitabine, in combination with paclitaxel, is indicated for the treatment of patients with unresectable, locally recurrent or metastatic breast cancer who have relapsed following adjuvant/neoadjuvant chemotherapy. Prior chemotherapy should have included an anthracycline unless clinically contraindicated.

4.2 Posology and method of administration

Gemcitabine should only be prescribed by a physician qualified in the use of anti-cancer chemotherapy.

Recommended posology

Bladder cancer

Combination use

The recommended dose for gemcitabine is 1000 mg/m², given by 30-minute infusion. The dose should be given on Days 1, 8 and 15 of each 28-day cycle in combination with cisplatin. Cisplatin is given at a recommended dose of 70 mg/m² on Day 1 following gemcitabine or day 2 of each 28-day cycle. This 4-week cycle is then repeated. Dosage reduction with each cycle or within a cycle may be applied based upon the grade of toxicity experienced by the patient.

Pancreatic cancer

The recommended dose of gemcitabine is 1000 mg/m², given by 30-minute intravenous infusion. This should be repeated once weekly for up to 7 weeks followed by a week of rest. Subsequent cycles should consist of injections once weekly for 3 consecutive weeks out of every 4 weeks. Dosage reduction with each cycle or within a cycle may be applied based upon the grade of toxicity experienced by the patient.

Non small Cell lung cancer

Monotherapy

The recommended dose of gemcitabine is 1000 mg/m², given by 30-minute intravenous infusion. This should be repeated once weekly for 3 weeks, followed by a 1-week rest period. This 4-week cycle is then repeated. Dosage reduction with each cycle or within a cycle may be applied based upon the grade of toxicity experienced by the patient.

Combination use

The recommended dose for gemcitabine is 1250 mg/m² body surface area given as a 30-minute intravenous infusion on Day 1 and 8 of the treatment cycle (21 days). Dosage reduction with each cycle or within a cycle may be applied based upon the grade of toxicity experienced by the patient. Cisplatin has been used at doses between 75-100 mg/m² once every 3 weeks.

Breast cancer

Combination use

Gemcitabine in combination with paclitaxel is recommended using paclitaxel (175 mg/m²) administered on Day 1 over approximately 3-hours as an intravenous infusion, followed by gemcitabine (1250 mg/m²) as a 30-minute intravenous infusion on Days 1 and 8 of each 21-day cycle. Dose reduction with each cycle or within a cycle may be applied based upon the grade of toxicity experienced by the patient. Patients should

have an absolute granulocyte count of at least 1,500 ($\times 10^6/l$) prior to initiation of gemcitabine + paclitaxel combination.

Ovarian cancer

Combination use

Gemcitabine in combination with carboplatin is recommended using gemcitabine 1000 mg/m² administered on Days 1 and 8 of each 21-day cycle as a 30-minute intravenous infusion. After gemcitabine, carboplatin will be given on Day 1 consistent with a target Area under curve (AUC) of 4.0 mg/ml·min. Dosage reduction with each cycle or within a cycle may be applied based upon the grade of toxicity experienced by the patient.

Monitoring for toxicity and dose modification due to toxicity

Dose modification due to non haematological toxicity

Periodic physical examination and checks of renal and hepatic function should be made to detect non-haematological toxicity. Dosage reduction with each cycle or within a cycle may be applied based upon the grade of toxicity experienced by the patient. In general, for severe (Grade 3 or 4) non-haematological toxicity, except nausea/vomiting, therapy with gemcitabine should be withheld or decreased depending on the judgement of the treating physician. Doses should be withheld until toxicity has resolved in the opinion of the physician.

For cisplatin, carboplatin, and paclitaxel dosage adjustment in combination therapy, please refer to the corresponding Summary of Product Characteristics.

Dose modification due to haematological toxicity

Initiation of a cycle

For all indications, the patient must be monitored before each dose for platelet and granulocyte counts. Patients should have an absolute granulocyte count of at least 1,500 ($\times 10^6/l$) and platelet count of 100,000 ($\times 10^6/l$) prior to the initiation of a cycle.

Within a cycle

Dose modifications of gemcitabine within a cycle should be performed according to the following tables:

Dose modification of gemcitabine within a cycle for bladder cancer, NSCLC and pancreatic cancer, given in monotherapy or in combination with cisplatin		
Absolute granulocyte count ($\times 10^6/l$)	Platelet count ($\times 10^6/l$)	Percentage of standard dose of gemcitabine (%)
> 1,000 and	> 100,000	100
500-1,000 or	50,000-100,000	75
<500 or	< 50,000	Omit dose *

*Treatment omitted will not be re-instated within a cycle before the absolute granulocyte count reaches at least 500 ($\times 10^6/l$) and the platelet count reaches 50,000 ($\times 10^6/l$).

Dose modification of gemcitabine within a cycle for breast cancer, given in combination with paclitaxel		
Absolute granulocyte count (x 10⁶/l)	Platelet count (x 10⁶/l)	Percentage of standard dose of gemcitabine (%)
≥ 1,200 and	>75,000	100
1,000- <1,200 or	50,000-75,000	75
700- <1,000 and	≥ 50,000	50
<700 or	<50,000	Omit dose*

*Treatment omitted will not be re-instated within a cycle. Treatment will start on day 1 of the next cycle once the absolute granulocyte count reaches at least 1,500 (x10⁶/l) and the platelet count reaches 100,000 (x10⁶/l).

Dose modification of gemcitabine within a cycle for ovarian cancer, given in combination with carboplatin		
Absolute granulocyte count (x 10⁶/l)	Platelet count (x 10⁶/l)	Percentage of standard dose of gemcitabine (%)
> 1,500 and	≥ 100,000	100
1000-1,500 or	75,000-100,000	50
<1000 or	< 75,000	Omit dose*

*Treatment omitted will not be re-instated within a cycle. Treatment will start on day 1 of the next cycle once the absolute granulocyte count reaches at least 1,500 (x10⁶/l) and the platelet count reaches 100,000 (x10⁶/l).

Dose modifications due to haematological toxicity in subsequent cycles, for all indications

The gemcitabine dose should be reduced to 75% of the original cycle initiation dose, in the case of the following haematological toxicities:

- Absolute granulocyte count < 500 x 10⁶/l for more than 5 days
- Absolute granulocyte count < 100 x 10⁶/l for more than 3 days
- Febrile neutropaenia
- Platelets < 25,000 x 10⁶/l
- Cycle delay of more than 1 week due to toxicity

Method of administration

Gemcitabine is tolerated well during infusion and may be administered ambulant. If extravasation occurs, generally the infusion must be stopped immediately and started again in another blood vessel. The patient should be monitored carefully after the administration.

Reconstitution: The recommended diluent for reconstitution of Gemcitabine for injection is 0.9% Sodium Chloride Injection without preservatives. Due to solubility considerations, the maximum concentration for Gemcitabine for injection upon reconstitution is 40 mg/ml. Reconstitution at concentrations greater than 40 mg/ml may result in incomplete dissolution, and should be avoided.

To reconstitute, add 5 ml of 0.9% Sodium Chloride Injection, to the 200mg vial, add 25 ml of 0.9% Sodium Chloride Injection to the 1gm vial, to 1.4 gm vial add 35 ml 0.9% Sodium Chloride Injection.

Shake to dissolve. These dilutions each yield a gemcitabine concentration of 38 mg/mL to 40 mg/ml which includes accounting for the displacement volume of the lyophilized powder.

The appropriate amount of drug may be administered as prepared or further diluted with 0.9% Sodium Chloride Injection to concentrations as low as 0.1 mg/mL.

After reconstitution with 0.9% Sodium Chloride Injection, the pH of the resulting solution lies in the range of 2.7 to 3.3. The solution should be inspected visually for particulate matter and discoloration, prior to administration, whenever solution or container permit. If particulate matter or discoloration is found, do not administer.

When prepared as directed, Gemcitabine for injection solutions are stable for 24 hours at controlled room temperature 20°C to 25°C. Do not refrigerate after reconstitution.

Discard unused portion. Solutions of reconstituted Gemcitabine for injection should not be refrigerated, as crystallization may occur.

Caution should be exercised in handling and preparing Gemcitabine for injection solutions. The use of gloves is recommended. If Gemcitabine for injection solution contacts the skin or mucosa, immediately wash the skin thoroughly with soap and water or rinse the mucosa with copious amounts of water.

Special populations:

Patients with renal or hepatic impairment

Gemcitabine should be used with caution in patients with hepatic or renal impairment as there is insufficient information from clinical studies to allow for clear dose recommendations for these patient populations.

Elderly people

Gemcitabine has been well tolerated in patients over the age of 65. There is no evidence to suggest that dose adjustments, other than those already recommended for all patients, are necessary in the elderly.

Paediatric population (< 18 years)

Gemcitabine is not recommended for use in children under 18 years of age due to insufficient data on safety and efficacy.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed.

Breast-feeding

4.4 Special warnings and precautions for use

Prolongation of the infusion time and increased dosing frequency have been shown to increase toxicity.

Haematological toxicity

Gemcitabine can suppress bone marrow function as manifested by leucopaenia, thrombocytopaenia and anaemia.

Patients receiving gemcitabine should be monitored prior to each dose for platelet, leucocyte and granulocyte counts. Suspension or modification of therapy should be considered when drug-induced bone marrow depression is detected. However, myelosuppression is short lived and usually does not result in dose reduction and rarely in discontinuation.

Peripheral blood counts may continue to deteriorate after gemcitabine administration has been stopped. In patients with impaired bone marrow function, the treatment should be started with caution. As with other cytotoxic treatments, the risk of cumulative bone-marrow suppression must be considered when gemcitabine treatment is given together with other chemotherapy.

Hepatic and renal impairment

Gemcitabine should be used with caution in patients with hepatic impairment or with impaired renal function as there is insufficient information from clinical studies to allow clear dose recommendation for this patient population.

Administration of gemcitabine in patients with concurrent liver metastases or a pre-existing medical history of hepatitis, alcoholism or liver cirrhosis may lead to exacerbation of the underlying hepatic impairment.

Laboratory evaluation of renal and hepatic function (including virological tests) should be performed periodically.

Concomitant radiotherapy

Concomitant radiotherapy (given together or ≤ 7 days apart): Toxicity has been reported (see section 4.5 for details and recommendations for use).

Live vaccinations

Yellow fever vaccine and other live attenuated vaccines are not recommended in patients treated with gemcitabine.

Nervous system

Posterior reversible encephalopathy syndrome

Reports of posterior reversible encephalopathy syndrome (PRES) with potentially severe consequences have been reported in patients receiving gemcitabine as single agent or in combination with other chemotherapeutic agents. Acute hypertension and seizure activity were reported in most gemcitabine patients experiencing PRES, but other symptoms such as headache, lethargy, confusion and blindness could also be present. Diagnosis is optimally confirmed by magnetic resonance imaging (MRI). PRES was typically reversible with appropriate supportive measures. Gemcitabine should be permanently discontinued and supportive measures implemented, including blood pressure control and anti-seizure therapy, if PRES develops during therapy.

Cardiovascular

Due to the risk of cardiac and/or vascular disorders with gemcitabine, particular caution must be exercised with patients presenting a history of cardiovascular events.

Capillary leak syndrome

Capillary leak syndrome has been reported in patients receiving gemcitabine as single agent or in combination with other chemotherapeutic agents (see section 4.8). The condition is usually treatable if recognised early and managed appropriately, but fatal cases have been reported. The condition involves systemic capillary hyperpermeability during which fluid and proteins from the intravascular space leak into the interstitium. The clinical features include generalised oedema, weight gain, hypoalbuminaemia, severe hypotension, acute renal impairment and pulmonary oedema. Gemcitabine should be discontinued and supportive measures implemented if capillary leak syndrome develops during therapy. Capillary leak syndrome can occur in later cycles and has been associated in the literature with adult respiratory distress syndrome.

Pulmonary

Pulmonary effects, sometimes severe (such as pulmonary oedema, interstitial pneumonitis or adult respiratory distress syndrome (ARDS)) have been reported in association with gemcitabine therapy. The aetiology of these effects is unknown. If such effects develop, consideration should be made to discontinuing gemcitabine therapy. Early use of supportive care measure may help ameliorate the condition.

Renal

Haemolytic uraemic syndrome

Clinical findings consistent with the haemolytic uraemic syndrome (HUS) were rarely reported (post-marketing data) in patients receiving gemcitabine (see section 4.8). HUS is a potentially life-threatening disorder. Gemcitabine should be discontinued at the first signs of any evidence of microangiopathic haemolytic anaemia, such as rapidly falling haemoglobin with concomitant thrombocytopenia, elevation of serum bilirubin, serum creatinine, blood urea nitrogen, or LDH. Renal failure may not be reversible with discontinuation of therapy and dialysis may be required.

Fertility

In fertility studies gemcitabine caused hypospermatogenesis in male mice (see section 5.3). Therefore, men being treated with gemcitabine are advised not to father a child during and up to 6 months after treatment and to seek further advice regarding cryoconservation of sperm prior to treatment because of the possibility of infertility due to therapy with gemcitabine.

4.5 Interaction with other medicinal products and other forms of interaction

No specific interaction studies have been performed

Radiotherapy

Concurrent (given together or ≤ 7 days apart) - Toxicity associated with this multimodality therapy is dependent on many different factors, including dose of gemcitabine, frequency of gemcitabine administration, dose of radiation, radiotherapy planning technique, the target tissue, and target volume. Pre-clinical and clinical studies have shown that gemcitabine has radiosensitising activity. In a single trial, where gemcitabine at a dose of 1,000 mg/m² was administered concurrently for up to 6 consecutive weeks with therapeutic thoracic radiation to patients with non-small cell lung cancer, significant toxicity in the form of severe, and potentially life threatening mucositis, especially oesophagitis, and pneumonitis was observed, particularly in

patients receiving large volumes of radiotherapy [median treatment volumes 4,795 cm³]. Studies done subsequently have suggested that it is feasible to administer gemcitabine at lower doses with concurrent radiotherapy with predictable toxicity, such as a phase II study in non-small cell lung cancer, where thoracic radiation doses of 66 Gy were applied concomitantly with an administration with gemcitabine (600 mg/m², four times) and cisplatin (80 mg/m² twice) during 6 weeks. The optimum regimen for safe administration of gemcitabine with therapeutic doses of radiation has not yet been determined in all tumour types.

Non-concurrent (given >7 days apart) - Analysis of the data does not indicate any enhanced toxicity when gemcitabine is administered more than 7 days before or after radiation, other than radiation recall. Data suggest that gemcitabine can be started after the acute effects of radiation have resolved or at least one week after radiation.

Radiation injury has been reported on targeted tissues (e.g. oesophagitis, colitis, and pneumonitis) in association with both concurrent and non-concurrent use of gemcitabine.

Others

Yellow fever and other live attenuated vaccines are not recommended due to the risk of systemic, possibly fatal, disease, particularly in immunosuppressed patients.

4.6 Fertility, pregnancy and lactation

Women of childbearing age/contraception in men and women

Due to the genotoxic potential of gemcitabine, women of childbearing potential should use effective methods of contraception during treatment with gemcitabine and for 6 months after stopping treatment.

Men should be advised to use effective methods of contraception and not to father children during treatment with gemcitabine and for 3 months after stopping it.

Pregnancy

There are no adequate data from the use of gemcitabine in pregnant women. Studies in animals have shown reproductive toxicity. Based on results from animal studies and the mechanism of action of gemcitabine, this substance should not be used during pregnancy unless clearly necessary. Women should be advised not to become pregnant during treatment with gemcitabine and to warn their attending physician immediately, should this occur after all.

Breast-feeding

It is not known whether gemcitabine is excreted in human milk and adverse effects on the suckling child cannot be excluded. Breast-feeding must be discontinued during gemcitabine therapy.

Fertility

In fertility studies gemcitabine caused hypospermatogenesis in male mice. Therefore, men being treated with gemcitabine are advised not to father a child during and up to 6 months after treatment and to seek further advice regarding cryoconservation of sperm prior to treatment because of the possibility of infertility due to therapy with gemcitabine.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. However, gemcitabine has been reported to cause mild to moderate somnolence, especially in combination with alcohol consumption. Patients should be cautioned against driving or operating machinery until it is established that they do not become somnolent.

4.8 Adverse Effects

The most commonly reported adverse drug reactions associated with Gemcitabine treatment include: nausea with or without vomiting, raised liver transaminases (AST/ALT) and alkaline phosphatase, reported in approximately 60% of patients; proteinuria and haematuria reported in approximately 50% patients; dyspnoea reported in 10-40% of patients (highest incidence in lung cancer patients); allergic skin rashes occur in approximately 25% of patients and are associated with itching in 10% of patients.

Frequencies are defined as: Very common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1000$), Very Rare ($< 1/10,000$), Not known (cannot be estimated from the available data).

The following table of undesirable effects and frequencies is based on data from clinical trials. Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

System Organ Class	Frequency	Adverse Event
Infections and infestations	Common	Infections
	Not known	Sepsis
Blood and lymphatic system disorders	Very common	Leucopaenia (Neutropaenia Grade 3 = 19.3 %; Grade 4 = 6 %). Bone-marrow suppression is usually mild to moderate and mostly affects the granulocyte count, Thrombocytopaenia, Anaemia
	Common	Febrile neutropaenia
	Very rare	Thrombocytosis, Thrombotic microangiopathy
Immune system disorders	Very Rare	Anaphylactoid reaction
Metabolism and nutrition disorders	Common	Anorexia
Nervous system disorders	Common	Headache, Insomnia, Somnolence
	Uncommon	Cerebrovascular accident
	Very rare	Posterior reversible encephalopathy syndrome
Cardiac disorders	Uncommon	Arrhythmias, predominantly supraventricular in nature Heart failure
	Rare	Myocardial infarct
Vascular disorders	Rare	Clinical signs of peripheral vasculitis and gangrene, Hypotension

	Very rare	Capillary leak syndrome
Respiratory, thoracic and mediastinal disorders	Very common	Dyspnoea –usually mild and passes rapidly without treatment
	Common	Cough, Rhinitis
	Uncommon	Interstitial pneumonitis, Bronchospasm – usually mild and transient but may require parenteral treatment
	Rare	Pulmonary oedema, Adult respiratory distress syndrome
	Not known	Pulmonary eosinophilia
Gastrointestinal disorders	Very common	Vomiting, Nausea
	Common	Diarrhoea, Stomatitis and ulceration of the mouth, Constipation
	Very rare	Ischaemic colitis
Hepatobiliary disorders	Very common	Elevation of liver transaminases (AST and ALT) and alkaline phosphatase
	Common	Increased bilirubin
	Uncommon	Serious hepatotoxicity, including liver failure and death
	Rare	Increased gamma-glutamyl transferase (GGT)
Skin and subcutaneous tissue disorders	Very common	Allergic skin rash frequently associated with pruritus, Alopecia
	Common	Itching, Sweating
	Rare	Severe skin reactions, including desquamation and bullous skin eruptions, Ulceration, Vesicle and sore formation, Scaling
	Very rare	Toxic epidermal necrolysis, Stevens-Johnson Syndrome
	Not known	Pseudocellulitis
Musculoskeletal and connective tissue disorders	Common	Back pain, Myalgia

Renal and urinary disorders	Very Common	Haematuria, Mild proteinuria
	Uncommon	Renal failure, Haemolytic uraemic syndrome
General disorders and administration site conditions	Very common	Influenza-like symptoms - the most common symptoms are fever, headache, chills, myalgia, asthenia and anorexia. Cough, rhinitis, malaise, perspiration and sleeping difficulties have also been reported, Oedema/peripheral oedema, including facial oedema. Oedema is usually reversible after stopping treatment.
	Common	Fever, Asthenia, Chills
	Rare	Injection site reactions - mainly mild in nature
Injury, poisoning, and procedural complications	Rare	Radiation toxicity, Radiation recall

Combination use in breast cancer

The frequency of grade 3 and 4 haematological toxicities, particularly neutropaenia, increases when gemcitabine is used in combination with paclitaxel. However, the increase in these adverse reactions is not associated with an increased incidence of infections or haemorrhagic events. Fatigue and febrile neutropaenia occur more frequently when gemcitabine is used in combination with paclitaxel. Fatigue, which is not associated with anaemia, usually resolves after the first cycle.

Grade 3 and 4 Adverse Events Paclitaxel versus gemcitabine plus paclitaxel				
	Number (%) of Patients			
	Paclitaxel arm (N=259)		Gemcitabine plus Paclitaxel arm (N=262)	
	Grade 3	Grade 4	Grade 3	Grade 4
Laboratory				
Anaemia	5 (1.9)	1 (0.4)	15 (5.7)	3 (1.1)
Thrombocytopaenia	0	0	14 (5.3)	1 (0.4)
Neutropaenia	11 (4.2)	17 (6.6)*	82 (31.3)	45 (17.2)*
Non-laboratory				
Febrile neutropaenia	3 (1.2)	0	12 (4.6)	1(0.4)

Fatigue	3 (1.2)	1 (0.4)	15 (5.7)	2 (0.8)
Diarrhoea	5 (1.9)	0	8 (3.1)	0
Motor neuropathy	2 (0.8)	0	6 (2.3)	1 (0.4)
Sensory neuropathy	9 (3.5)	0	14 (5.3)	1 (0.4)

*Grade 4 neutropaenia lasting for more than 7 days occurred in 12.6% of patients in the combination arm and 5.0% of patients in the paclitaxel arm.

Combination use in bladder cancer

Grade 3 and 4 Adverse Events MVAC versus Gemcitabine plus cisplatin				
	Number (%) of Patients			
	MVAC* arm (N=196)		Gemcitabine plus cisplatin arm (N=200)	
	Grade 3	Grade 4	Grade 3	Grade 4
Laboratory				
Anaemia	30 (16)	4 (2)	47 (24)	7 (4)
Thrombocytopenia	15 (8)	25 (13)	57 (29)	57 (29)
Non-laboratory				
Nausea and vomiting	37 (19)	3 (2)	44 (22)	0 (0)
Diarrhoea	15 (8)	1 (1)	6 (3)	0 (0)
Infection	19 (10)	10 (5)	4 (2)	1 (1)
Stomatitis	34 (18)	8 (4)	2 (1)	0 (0)

*Methotrexate, Vinblastine, Doxorubicin and Cisplatin

Combination use in ovarian cancer

Grade 3 and 4 Adverse Events Carboplatin versus Gemcitabine plus carboplatin				
	Number (%) of Patients			
	Carboplatin arm (N=174)		Gemcitabine plus carboplatin arm (N=175)	
	Grade 3	Grade 4	Grade 3	Grade 4
Laboratory				
Anaemia	10(5.7)	4 (2.3)	39 (22.3)	9 (5.1)
Neutropaenia	19(10.9)	2(1.1)	73(41.7)	50 (28.6)
Thrombocytopenia	18(10.3)	2(1.1)	53(30.3)	8 (4.6)
Leucopaenia	11(6.3)	1(0.6)	84(48.0)	9 (5.1)
Non-laboratory				

Haemorrhage	0(0.0)	0(0.0)	3(1.8)	(0.0)
Febrile neutropaenia	0(0.0)	0(0.0)	2(1.1)	(0.0)
Infection without neutropaenia	0(0)	0(0.0)	(0.0)	1(0.6)

Sensory neuropathy was also more frequent in the combination arm than with single agent Carboplatin

4.9 Overdose

There is no known antidote for overdose of gemcitabine. Doses as high as 5700 mg/m² have been administered by intravenous infusion over 30-minutes every 2 weeks with clinically acceptable toxicity. In the event of suspected overdose, the patient should be monitored with appropriate blood counts and receive supportive therapy, as necessary.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: pyrimidine analogues ATC code: L01BC05

Cytotoxic activity in cell cultures

Gemcitabine shows significant cytotoxic effects against a variety of cultured murine and human tumour cells. Its action is phase-specific such that gemcitabine primarily kills cells that are undergoing DNA synthesis (S-phase) and, under certain circumstances, blocks the progression of cells at the junction of the G1/S phase boundary. *In vitro*, the cytotoxic effect of gemcitabine is dependent on both concentration and time.

Mechanism of action

Cellular metabolism and mechanism of action: Gemcitabine (dFdC), which is a pyrimidine antimetabolite, is metabolised intracellularly by nucleoside kinase to the active diphosphate (dFdCDP) and triphosphate (dFdCTP) nucleosides. The cytotoxic effect of gemcitabine is due to inhibition of DNA synthesis by two mechanisms of action by dFdCDP and dFdCTP. First, dFdCDP inhibits ribonucleotide reductase, which is uniquely responsible for catalysing the reactions that produce deoxynucleoside triphosphates (dCTP) for DNA synthesis. Inhibition of this enzyme by dFdCDP reduces the concentration of deoxynucleosides in general and, in particular, dCTP. Second, dFdCTP competes with dCTP for incorporation into DNA (self-potentialiation).

Likewise, a small amount of gemcitabine may also be incorporated into RNA. Thus, the reduced intracellular concentration of dCTP potentiates the incorporation of dFdCTP into DNA. DNA polymerase epsilon lacks the ability to eliminate gemcitabine and to repair the growing DNA strands. After gemcitabine is incorporated into DNA, one additional nucleotide is added to the growing DNA strands. After this addition there is essentially a complete inhibition in further DNA synthesis (masked chain termination). After incorporation into DNA, gemcitabine appears to induce the programmed cell death process known as apoptosis.

5.2 Pharmacokinetic properties

Absorption

Peak plasma concentrations (obtained within 5 minutes of the end of the infusion) were 3.2 to 45.5 µg/ml. Plasma concentrations of the parent compound following a dose of 1,000 mg/m²/30-minutes are greater than

5 µg/ml for approximately 30-minutes after the end of the infusion, and greater than 0.4 µg/ml for an additional hour.

Distribution

The volume of distribution of the central compartment was 12.4 l/m² for women and 17.5 l/m² for men (inter-individual variability was 91.9%). The volume of distribution of the peripheral compartment was 47.4 l/m². The volume of the peripheral compartment was not sensitive to gender. The plasma protein binding was considered to be negligible.

Half-life: This ranged from 42 to 94 minutes depending on age and gender. For the recommended dosing schedule, gemcitabine elimination should be virtually complete within 5 to 11 hours of the start of the infusion. Gemcitabine does not accumulate when administered once weekly.

Biotransformation

Gemcitabine is rapidly metabolised by cytidine deaminase in the liver, kidney, blood and other tissues. Intracellular metabolism of gemcitabine produces the gemcitabine mono, di and triphosphates (dFdCMP, dFdCDP and dFdCTP) of which dFdCDP and dFdCTP are considered active. These intracellular metabolites have not been detected in plasma or urine. The primary metabolite, 2'-deoxy-2', 2'-difluorouridine (dFdU), is not active and is found in plasma and urine.

Elimination

Systemic clearance ranged from 29.2 l/hr/m² to 92.2 l/hr/m² depending on gender and age (inter-individual variability was 52.2%). Clearance for women is approximately 25% lower than the values for men. Although rapid, clearance for both men and women appears to decrease with age. For the recommended gemcitabine dose of 1000 mg/m² given as a 30-minute infusion, lower clearance values for women and men should not necessitate a decrease in the gemcitabine dose. Urinary excretion: Less than 10% is excreted as unchanged drug.

Renal clearance was 2 to 7 l/hr/m².

During the week following administration, 92 to 98% of the dose of gemcitabine administered is recovered, 99% in the urine, mainly in the form of dFdU and 1% of the dose is excreted in faeces.

dFdCTP kinetics

This metabolite can be found in peripheral blood mononuclear cells and the information below refers to these cells. Intracellular concentrations increase in proportion to gemcitabine doses of 35-350 mg/m²/30-minutes, which give steady state concentrations of 0.4-5 µg/ml. At gemcitabine plasma concentrations above 5 µg/ml, dFdCTP levels do not increase, suggesting that the formation is saturable in these cells.

Half-life of terminal elimination: 0.7-12 hours.

dFdU kinetics

Peak plasma concentrations (3-15 minutes after end of 30-minute infusion, 1000 mg/m²): 28-52 µg/ml. Trough concentration following once weekly dosing: 0.07-1.12 µg/ml, with no apparent accumulation. Triphasic plasma concentration versus time curve, mean half-life of terminal phase 65 hours (range 33-84 hr).

Formation of dFdU from parent compound: 91%-98%.

Mean volume of distribution of central compartment: 18 l/m² (range 11-22 l/m²).

Mean steady state volume of distribution (V_{ss}): 150 l/m² (range 96-228 l/m²).

Tissue distribution: Extensive.

Mean apparent clearance: 2.5 l/hr/m² (range 1-4 l/hr/m²).

Urinary excretion: All.

Gemcitabine and paclitaxel combination therapy

Combination therapy did not alter the pharmacokinetics of either gemcitabine or paclitaxel.

Gemcitabine and carboplatin combination therapy

When given in combination with carboplatin the pharmacokinetics of gemcitabine were not altered.

Renal impairment

Mild to moderate renal insufficiency (GFR from 30 ml/min to 80 ml/min) has no consistent, significant effect on gemcitabine pharmacokinetics.

5.3 Preclinical safety data

In repeat-dose studies of up to 6 months in duration in mice and dogs, the principal finding was schedule and dose-dependent haematopoietic suppression which was reversible.

Gemcitabine is mutagenic in an *in vitro* mutation test and an *in vivo* bone marrow micronucleus test. Long term animal studies evaluating the carcinogenic potential have not been performed.

In fertility studies, gemcitabine caused reversible hypospermatogenesis in male mice. No effect on the fertility of females has been detected.

Evaluation of experimental animal studies has shown reproductive toxicity e.g. birth defects and other effects on the development of the embryo or foetus, the course of gestation or peri- and postnatal development.

6. Pharmaceutical particulars

6.1 List of excipients

Mannitol USP

Sodium Acetate Trihydrate USP

Sodium Hydroxide USP

Water for Injection USP

6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products, except those mentioned.

6.3 Shelf life

24 months

6.4 Special precautions for storage

Store below 30°C. Protect from light.
Do not refrigerate after reconstitution.

6.5 Nature and contents of container

Gemcitabine for Injection USP is available in a vial containing Gemcitabine hydrochloride USP equivalent to Gemcitabine 200mg, 1g and 1.4g.

6.6 Special precautions for disposal and other handling

Handling

The normal safety precautions for cytostatic agents must be observed when preparing and disposing of the infusion solution. Handling of the concentrate should be done in a safety box and protective coats and gloves should be used. If no safety box is available, the equipment should be supplemented with a mask and protective glasses.

If the preparation comes into contact with the eyes, this may cause serious irritation. The eyes should be rinsed immediately and thoroughly with water. If there is lasting irritation, a doctor should be consulted. If the solution is spilled on the skin, rinse thoroughly with water.

7. Marketing authorisation holder

<< marketing authorisation holder details >>

8. Marketing authorisation number(s)

<< marketing authorisation number details >>

9. Date of first authorisation/renewal of the authorisation

<< date of authorisation/renewal >>

10. Date of revision of the text

15-Nov-2023

Company Contact Details

Naprod Life Sciences Pvt. Ltd.
Factory: G-17/1, MIDC, Tarapur, Boisar,
Dist. – Palghar – 401 506,
Maharashtra, INDIA.
Tel.: +91 2525 661110/ 661111