



**1.3 Product Information**

**1.3.1 Summary of Product Characteristics (SmPC)**

**SUMMARY OF PRODUCT CHARACTERISTICS**

**1. Name of the Medicinal Product**

**SWIVASC 10**

(Amlodipine Besylate Tablets USP 10 mg)

**1.2. Strength**

10 mg

**1.3. Pharmaceutical Form**

Solid Dosage form (Tablet)

**2. Quality and Quantitative Composition**

**Qualitative Composition:**

Each film coated Tablet contains:

Amlodipine Besylate USP

E.q. to Amlodipine                    10 mg

Colour:      Titanium Dioxide BP

Sr. No.	Ingredients	Standard
1	Calcium Hydrogen Phosphate (Dibasic Calcium Phosphate)	BP



# VAPI CARE PHARMA PVT. LTD.

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

2	Microcrystalline Cellulose pH 101 (Chemixil)	BP
3	Methyl Hydroxybenzoate (Methyl Paraben)	BP
4	Propyl Hydroxybenzoate (Propyl Paraben)	BP
5	Povidone (PVP K-30)	BP
6	Isopropyl Alcohol	BP
7	Magnesium Stearate	BP
8	Purified Talc	BP
9	Colloidal Anhydrous Silica	BP
10	Sodium Lauryl Sulphate	BP
11	Croscarmellose Sodium (Ac-Di-Sol)	BP
12	Hydroxypropylcellulose (Low Substituted)	BP
13	Insta Coat ICS223 White	IH
14	Isopropyl Alcohol	BP
15	Dichloromethane	BP

### Quantitative Composition:

Each film coated Tablet contains:

Amlodipine Besylate USP

E.q. to Amlodipine                    10 mg

Colour:            Titanium Dioxide BP

Sr. No.	Ingredients	Standard	Quantity / Tablet
1	Calcium Hydrogen Phosphate (Dibasic Calcium Phosphate)	BP	50.000 mg
2	Microcrystalline Cellulose pH 101	BP	30.000 mg



**VAPI CARE PHARMA PVT. LTD.**

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

	(Chemixil)		
3	Methyl Hydroxybenzoate (Methyl Paraben)	BP	0.148 mg
4	Propyl Hydroxybenzoate (Propyl Paraben)	BP	0.017 mg
5	Povidone (PVP K-30)	BP	5.000 mg
6	Isopropyl Alcohol	BP	0.025 ml
7	Magnesium Stearate	BP	2.000 mg
8	Purified Talc	BP	3.000 mg
9	Colloidal Anhydrous Silica	BP	3.000 mg
10	Sodium Lauryl Sulphate	BP	2.000 mg
11	Croscarmellose Sodium (Ac-Di-Sol)	BP	10.000 mg
12	Hydroxypropylcellulose (Low Substituted)	BP	6.000 mg
13	Insta Coat ICS223 White	IH	4.000 mg
14	Isopropyl Alcohol	BP	0.0313 ml
15	Dichloromethane	BP	0.0470 ml

### 3. Pharmaceutical Form

Solid Dosage form (Tablets)

White to off white coloured circular biconvex film coated tablet, with "A" embossed on one side & "10" on other side.

### 4. Clinical Particulars

#### 4.1 Therapeutic indications

SWIVASC 10 is indicated in the following condition.



**VAPI CARE PHARMA PVT. LTD.**

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

- Hypertension
- Chronic stable angina pectoris
- Vasospastic (Prinzmetal's) angina

Amlodipine is used with or without other medications to treat high blood pressure. Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Amlodipine belongs to a class of drugs known as calcium channel blockers. It works by relaxing blood vessels so blood can flow more easily.

Amlodipine is also used to prevent certain types of chest pain (angina). It may help to increase ability to exercise and decrease the frequency of angina attacks. It should not be used to treat attacks of chest pain when they occur. Use other medications (such as sublingual nitroglycerin) to relieve attacks of chest pain as directed by doctor.

#### **4.2 Posology and method of administration**

##### Posology

##### Adults

For both hypertension and angina the usual initial dose is 5 mg Amlodipine once daily which may be increased to a maximum dose of 10 mg depending on the individual patient's response.

In hypertensive patients, SWIVASC 10 has been used in combination with a thiazide diuretic, alpha blocker, beta blocker, or an angiotensin converting enzyme inhibitor. For angina, SWIVASC 10 may be used as monotherapy or in combination with other antianginal medicinal products in patients with angina that is refractory to nitrates and/or to adequate doses of beta blockers.

No dose adjustment of SWIVASC 10 is required upon concomitant administration of thiazide diuretics, beta blockers, and angiotensin-converting enzyme inhibitors.



**VAPI CARE PHARMA PVT. LTD.**

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

#### Special populations

##### Elderly patients

SWIVASC 10 used at similar doses in elderly or younger patients is equally well tolerated. Normal dosage regimens are recommended in the elderly, but increase of the dosage should take place with care.

##### Patients with hepatic impairment

Dosage recommendations have not been established in patients with mild to moderate hepatic impairment; therefore dose selection should be cautious and should start at the lower end of the dosing range. The pharmacokinetics of amlodipine have not been studied in severe hepatic impairment. Amlodipine should be initiated at the lowest dose and titrated slowly in patients with severe hepatic impairment.

##### Patients with renal impairment

Changes in amlodipine plasma concentrations are not correlated with degree of renal impairment, therefore the normal dosage is recommended. Amlodipine is not dialysable.

##### Paediatric population

Children and adolescents with hypertension from 6 years to 17 years of age.

The recommended antihypertensive oral dose in paediatric patients ages 6-17 years is 2.5 mg once daily as a starting dose, up-titrated to 5 mg once daily if blood pressure goal is not achieved after 4 weeks. Doses in excess of 5 mg daily have not been studied in paediatric patients.

##### Children under 6 years old

No data are available.

##### Method of administration

Tablet for oral administration.



#### **4.3 Contraindications**

Amlodipine is contraindicated in patients with hypersensitivity to dihydropyridine derivatives, amlodipine or to any of the excipients used in the formulation.

Severe hypotension.

Shock (including cardiogenic shock).

Obstruction of the outflow tract of the left ventricle (e.g., high grade aortic stenosis).

Haemodynamically unstable heart failure after acute myocardial infarction.

#### **4.4 Special warnings and precautions for use**

The safety and efficacy of amlodipine in hypertensive crisis has not been established.

Patients with cardiac failure

Patients with heart failure should be treated with caution. In a long-term, placebo controlled study in patients with severe heart failure (NYHA class III and IV) the reported incidence of pulmonary oedema was higher in the amlodipine treated group than in the placebo group. Calcium channel blockers, including amlodipine, should be used with caution in patients with congestive heart failure, as they may increase the risk of future cardiovascular events and mortality.

Patients with hepatic impairment

The half-life of amlodipine is prolonged and AUC values are higher in patients with impaired liver function; dosage recommendations have not been established. Amlodipine should therefore be initiated at the lower end of the dosing range and caution should be used, both on initial treatment and when increasing the dose. Slow dose titration and careful monitoring may be required in patients with severe hepatic impairment.



#### Elderly patients

In the elderly increase of the dosage should take place with care.

#### Patients with renal impairment

Amlodipine may be used in such patients at normal doses. Changes in amlodipine plasma concentrations are not correlated with degree of renal impairment. Amlodipine is not dialysable.

### **4.5 Interaction with other medicinal products and other forms of interaction**

#### Effects of other medicinal products on amlodipine

##### CYP3A4 inhibitors

Concomitant use of amlodipine with strong or moderate CYP3A4 inhibitors (protease inhibitors,azole antifungals, macrolides like erythromycin or clarithromycin, verapamil or diltiazem) may give rise to significant increase in amlodipine exposure resulting in an increased risk of hypotension. The clinical translation of these PK variations may be more pronounced in the elderly. Clinical monitoring and dose adjustment may thus be required.

##### CYP3A4 inducers

There is no data available regarding the effect of CYP3A4 inducers on amlodipine. The concomitant use of CYP3A4 inducers (e.g., rifampicin, hypericum perforatum) may give a lower plasma concentration of amlodipine. Amlodipine should be used with caution together with CYP3A4 inducers.

Administration of amlodipine with grapefruit or grapefruit juice is not recommended as bioavailability may be increased in some patients resulting in increased blood pressure lowering effects.

##### Dantrolene (infusion)

In animals, lethal ventricular fibrillation and cardiovascular collapse are observed in association with hyperkalemia after administration of verapamil and intravenous dantrolene. Due to risk of hyperkalemia,



**VAPI CARE PHARMA PVT. LTD.**

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

it is recommended that the co-administration of calcium channel blockers such as amlodipine be avoided in patients susceptible to malignant hyperthermia and in the management of malignant hyperthermia.

#### Effects of amlodipine on other medicinal products

The blood pressure lowering effects of amlodipine adds to the blood pressure-lowering effects of other medicinal products with antihypertensive properties.

#### Tacrolimus

There is a risk of increased tacrolimus blood levels when co-administered with amlodipine but the pharmacokinetic mechanism of this interaction is not fully understood. In order to avoid toxicity of tacrolimus, administration of amlodipine in a patient treated with tacrolimus requires monitoring of tacrolimus blood levels and dose adjustment of tacrolimus when appropriate.

#### Cyclosporine

No drug interaction studies have been conducted with cyclosporine and amlodipine in healthy volunteers or other populations with the exception of renal transplant patients, where variable trough concentration increases (average 0% - 40%) of cyclosporine were observed. Consideration should be given for monitoring cyclosporine levels in renal transplant patients on amlodipine, and cyclosporine dose reductions should be made as necessary.

#### Simvastatin

Co-administration of multiple doses of 10 mg of amlodipine with 80 mg simvastatin resulted in a 77% increase in exposure to simvastatin compared to simvastatin alone. Limit the dose of simvastatin in patients on amlodipine to 20 mg daily.

In clinical interaction studies, amlodipine did not affect the pharmacokinetics of atorvastatin, digoxin or warfarin.





#### **4.6 Fertility, pregnancy and lactation**

##### **Pregnancy**

The safety of amlodipine in human pregnancy has not been established.

In animal studies, reproductive toxicity was observed at high doses.

Use in pregnancy is only recommended when there is no safer alternative and when the disease itself carries greater risk for the mother and foetus.

##### **Breast-feeding**

It is not known whether amlodipine is excreted in breast milk. A decision on whether to continue/discontinue breast-feeding or to continue/discontinue therapy with amlodipine should be made taking into account the benefit of breast-feeding to the child and the benefit of amlodipine therapy to the mother.

##### **Fertility**

Reversible biochemical changes in the head of spermatozoa have been reported in some patients treated by calcium channel blockers. Clinical data are insufficient regarding the potential effect of amlodipine on fertility. In one rat study, adverse effects were found on male fertility.

#### **4.7 Effects on ability to drive and use machines**

Amlodipine can have minor or moderate influence on the ability to drive and use machines. If patients taking amlodipine suffer from dizziness, headache, fatigue or nausea the ability to react may be impaired. Caution is recommended especially at the start of treatment.

#### **4.8 Undesirable effects**

##### **Summary of the safety profile**

The most commonly reported adverse reactions during treatment are somnolence, dizziness, headache, palpitations, flushing, abdominal pain, nausea, ankle swelling, oedema and fatigue.



## Tabulated list of adverse reactions

The following adverse reactions have been observed and reported during treatment with amlodipine with the following frequencies: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

System organ class	Frequency	Adverse reactions
<b>Blood and lymphatic system disorders</b>	Very rare	Leukocytopenia, thrombocytopenia
<b>Immune system disorders</b>	Very rare	Allergic reactions
<b>Metabolism and nutrition disorders</b>	Very rare	Hyperglycaemia
<b>Psychiatric disorders</b>	Uncommon	Depression, mood changes (including anxiety), insomnia
	Rare	Confusion
<b>Nervous system disorders</b>	Common	Somnolence, dizziness, headache (especially at the beginning of the treatment)
	Uncommon	Tremor, dysgeusia, syncope, hypoaesthesia, paraesthesia
	Very rare	Hypertonia, peripheral neuropathy
<b>Eye disorders</b>	Common	Visual disturbance (including diplopia)
<b>Ear and labyrinth disorders</b>	Uncommon	Tinnitus
<b>Cardiac disorders</b>	Common	Palpitations
	Uncommon	Arrhythmia (including bradycardia,



# VAPI CARE PHARMA PVT. LTD.

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

		ventricular tachycardia and atrial fibrillation)
	Very rare	Myocardial infarction
<b>Vascular disorders</b>	Common	Flushing
	Uncommon	Hypotension
	Very rare	Vasculitis
<b>Respiratory, thoracic and mediastinal disorders</b>	Common	Dyspnoea
	Uncommon	Cough, rhinitis
<b>Gastrointestinal disorders</b>	Common	Abdominal pain, nausea, dyspepsia, altered bowel habits (including diarrhoea and constipation)
	Uncommon	Vomiting, dry mouth
	Very rare	Pancreatitis, gastritis, gingival hyperplasia
<b>Hepatobiliary disorders</b>	Very rare	Hepatitis, jaundice, hepatic enzyme increased*
<b>Skin and subcutaneous tissue disorders</b>	Uncommon	Alopecia, purpura, skin discolouration, hyperhidrosis, pruritus, rash, exanthema, urticaria
	Very rare	Angioedema, erythema multiforme, exfoliative dermatitis, Stevens-Johnson syndrome, Quincke oedema, photosensitivity
<b>Musculoskeletal and connective tissue disorders</b>	Common	Ankle swelling, muscle cramps
	Uncommon	Arthralgia, myalgia, back pain
<b>Renal and urinary disorders</b>	Uncommon	Micturition disorder, nocturia, increased urinary frequency



<b>Reproductive system and breast disorders</b>	Uncommon	Impotence, gynaecomastia
<b>General disorders and administration site conditions</b>	Very common	Oedema
	Common	Fatigue, asthenia
	Uncommon	Chest pain, pain, malaise
<b>Investigations</b>	Uncommon	Weight increased, weight decreased

#### **4.9 Overdose**

In humans experience with intentional overdose is limited.

##### Symptoms

Available data suggest that gross overdosage could result in excessive peripheral vasodilatation and possibly reflex tachycardia. Marked and probably prolonged systemic hypotension up to and including shock with fatal outcome have been reported.

##### Treatment

Clinically significant hypotension due to amlodipine overdosage calls for active cardiovascular support including frequent monitoring of cardiac and respiratory function, elevation of extremities and attention to circulating fluid volume and urine output.

A vasoconstrictor may be helpful in restoring vascular tone and blood pressure, provided that there is no contraindication to its use. Intravenous calcium gluconate may be beneficial in reversing the effects of calcium channel blockade.

Gastric lavage may be worthwhile in some cases. In healthy volunteers the use of charcoal up to 2 hours after administration of amlodipine 10 mg has been shown to reduce the absorption rate of amlodipine.



Since amlodipine is highly protein-bound, dialysis is not likely to be of benefit

## **5. Pharmacological Properties**

### **5.1 Pharmacodynamic Properties**

Pharmacotherapeutic group: Calcium channel blockers, selective calcium channel blockers with mainly vascular effects. ATC Code: C08CA01.

Amlodipine is a calcium ion influx inhibitor of the dihydropyridine group (slow channel blocker or calcium ion antagonist) and inhibits the transmembrane influx of calcium ions into cardiac and vascular smooth muscle.

The mechanism of the antihypertensive action of amlodipine is due to a direct relaxant effect on vascular smooth muscle. The precise mechanism by which amlodipine relieves angina has not been fully determined but amlodipine reduces total ischaemic burden by the following two actions:

- 1) Amlodipine dilates peripheral arterioles and thus, reduces the total peripheral resistance (afterload) against which the heart works. Since the heart rate remains stable, this unloading of the heart reduces myocardial energy consumption and oxygen requirements.
- 2) The mechanism of action of amlodipine also probably involves dilatation of the main coronary arteries and coronary arterioles, both in normal and ischaemic regions. This dilatation increases myocardial oxygen delivery in patients with coronary artery spasm (Prinzmetal's or variant angina).

In patients with hypertension, once daily dosing provides clinically significant reductions of blood pressure in both the supine and standing positions throughout the 24 hour interval. Due to the slow onset of action, acute hypotension is not a feature of amlodipine administration.



In patients with angina, once daily administration of amlodipine increases total exercise time, time to angina onset, and time to 1 mm ST segment depression, and decreases both angina attack frequency and glyceryl trinitrate tablet consumption.

Amlodipine has not been associated with any adverse metabolic effects or changes in plasma lipids and is suitable for use in patients with asthma, diabetes, and gout.

## **5.2 Pharmacokinetic properties**

Absorption, distribution, plasma protein binding: After oral administration of therapeutic doses, amlodipine is well absorbed with peak blood levels between 6-12 hours post dose. Absolute bioavailability has been estimated to be between 64 and 80%. The volume of distribution is approximately 21 l/kg. In vitro studies have shown that approximately 97.5% of circulating amlodipine is bound to plasma proteins.

The bioavailability of amlodipine is not affected by food intake.

### **Biotransformation/elimination**

The terminal plasma elimination half-life is about 35-50 hours and is consistent with once daily dosing. Amlodipine is extensively metabolised by the liver to inactive metabolites with 10% of the parent compound and 60% of metabolites excreted in the urine.

### **Hepatic impairment**

Very limited clinical data are available regarding amlodipine administration in patients with hepatic impairment. Patients with hepatic insufficiency have decreased clearance of amlodipine resulting in a longer half-life and an increase in AUC of approximately 40-60%.

### **Elderly population**



The time to reach peak plasma concentrations of amlodipine is similar in elderly and younger subjects. Amlodipine clearance tends to be decreased with resulting increases in AUC and elimination half-life in elderly patients. Increases in AUC and elimination half-life in patients with congestive heart failure were as expected for the patient age group studied.

#### Paediatric population

A population PK study has been conducted in 74 hypertensive children aged from 1 to 17 years (with 34 patients aged 6 to 12 years and 28 patients aged 13 to 17 years) receiving amlodipine between 1.25 and 20 mg given either once or twice daily. In children 6 to 12 years and in adolescents 13-17 years of age the typical oral clearance (CL/F) was 22.5 and 27.4 L/hr respectively in males and 16.4 and 21.3 L/hr respectively in females. Large variability in exposure between individuals was observed. Data reported in children below 6 years is limited.

### **5.3 Preclinical safety data**

#### Reproductive toxicology

Reproductive studies in rats and mice have shown delayed date of delivery, prolonged duration of labour and decreased pup survival at dosages approximately 50 times greater than the maximum recommended dosage for humans based on mg/kg.

#### Impairment of fertility

There was no effect on the fertility of rats treated with amlodipine (males for 64 days and females 14 days prior to mating) at doses up to 10 mg/kg/day (8 times\* the maximum recommended human dose of 10 mg on a mg/m<sup>2</sup> basis). In another rat study in which male rats were treated with Amlodipine Besylate for 30 days at a dose comparable with the human dose based on mg/kg, decreased plasma follicle-stimulating hormone and testosterone were found as well as decreases in sperm density and in the number of mature spermatids and Sertoli cells.

#### Carcinogenesis, mutagenesis



Rats and mice treated with amlodipine in the diet for two years, at concentrations calculated to provide daily dosage levels of 0.5, 1.25, and 2.5 mg/kg/day showed no evidence of carcinogenicity. The highest dose (for mice, similar to, and for rats twice\* the maximum recommended clinical dose of 10 mg on a mg/m<sup>2</sup> basis) was close to the maximum tolerated dose for mice but not for rats.

Mutagenicity studies revealed no drug related effects at either the gene or chromosome levels.

\*Based on patient weight of 50 kg

## 6.0 Pharmaceutical Particulars

### 6.1 List of excipients

Sr. No.	Ingredients	Standard
1	Calcium Hydrogen Phosphate (Dibasic Calcium Phosphate)	BP
2	Microcrystalline Cellulose pH 101 (Chemixil)	BP
3	Methyl Hydroxybenzoate (Methyl Paraben)	BP
4	Propyl Hydroxybenzoate (Propyl Paraben)	BP
5	Povidone (PVP K-30)	BP
6	Isopropyl Alcohol	BP
7	Magnesium Stearate	BP
8	Purified Talc	BP
9	Colloidal Anhydrous Silica	BP
10	Sodium Lauryl Sulphate	BP
11	Croscarmellose Sodium (Ac-Di-Sol)	BP
12	Hydroxypropylcellulose (Low Substituted)	BP





**VAPI CARE PHARMA PVT. LTD.**

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

<b>13</b>	Insta Coat ICS223 White	IH
<b>14</b>	Isopropyl Alcohol	BP
<b>15</b>	Dichloromethane	BP

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

3 years

## **6.4 Special precautions for storage**

Store below 30°C protected from light, Keep out of reach of the children.

## **6.5 Nature and contents of container**

Alu/Alu Blister pack of 1 x 10, 3 x 10, and 10 x 10 tablets.

## **6.6 Special precautions for disposal**

No special requirements

Any unused product or waste material should be disposed of in accordance with local requirements

## **7. Marketing Authorization Holder/ Registrant**

**Swiss Pharma Nig Ltd**

## **8. Marketing Authorization Numbers**

**A4-2279**



**VAPI CARE PHARMA PVT. LTD.**

*SWIVASC 10 (Amlodipine Besylate Tablets USP 10 mg)*

*MODULE 1 (Administrative & Prescribing Information)*

**9. Date of first authorization/renewal of the authorization**

**5/02/2019**

**10. Date of revision of the text**

-----