

**BIORAJ PHARMACEUTICALS LIMITED**

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

**SUMMARY OF PRODUCT CHARACTERISTICS**

**1. NAME OF THE MEDICINAL PRODUCT**

Bioraquine Syrup

**2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each 5ml contains Chloroquine Phosphate 80.0mg, equivalent to Chloroquine base 50mg see section 6.1.

**3. PHARMACEUTICAL FORM**

Oral Liquid.

**4. CLINICAL PARTICULARS**

**4.1 Therapeutic indications**

For the prophylaxis, suppression and treatment of malaria.

**4.2 Posology and method of administration**

As directed by a physician.

**4.3 Contraindications**

Bioraquine is contraindicated in severe hypersensitivity to chloroquine and in known allergy to anti-histamines or parabens.

**4.4 Special warnings and precautions for use**

When used as malaria prophylaxis official guidelines and local information on prevalence of resistance to anti-malarial drugs should be taken into consideration.

Irreversible retinal damage and corneal changes may develop during long term therapy and after the drug has been discontinued. Ophthalmic examination prior to, and at 3-6 monthly intervals during use is required if patients are receiving chloroquine:

- At continuous high doses for longer than 12 months
- As weekly treatment for longer than 3 years
- When total consumption exceeds 1.6g/kg (cumulative dose 100g)

Patients should be advised to stop taking the drug immediately and seek the advise of their doctor if any disturbances of vision occur.

Bone marrow suppression may occur rarely so full blood counts should be carried out during extended treatment. Caution is required if drugs known to induce blood disorders are used concurrently.

**BIORAJ PHARMACEUTICALS LIMITED**

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

**SUMMARY OF PRODUCT CHARACTERISTICS**

Use with caution in patients with impaired hepatic function, particularly cirrhosis.

Use with caution in patients with porphyria as the disease may be precipitated. This may be especially apparent in patients with a high alcohol intake.

Use with caution in patients with a renal impairment.

Use with caution in patients with a history of epilepsy, convulsions and other neurological disorders.

Use with caution in patients with psoriasis as chloroquine may precipitate a severe attack.

Use with caution in patients with severe gastro-intestinal disease.

Use with caution in patients with glucose-6-phosphate dehydrogenase deficiency, as there may be risk of haemolysis.

Chloroquine has been shown to cause severe hypoglycaemia including loss of consciousness that could be life threatening in patients treated with and without antidiabetic medications. Patients treated with chloroquine should be warned about the risk of hypoglycaemia and the associated clinical signs and symptoms. Patients presenting with clinical symptoms suggestive of hypoglycaemia during treatment with chloroquine should have their blood glucose level checked and treatment reviewed as necessary.

A small number of cases of diffuse parenchymal lung disease have been identified in patients taking chloroquine. A response after therapy with steroids has been observed in some of these cases.

Cases of drug rash with eosinophilia and systemic symptoms (DRESS) syndrome have been identified in patients taking chloroquine. Recovery after discontinuation of treatment and response after therapy with steroids has been observed.

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

**BIORAJ PHARMACEUTICALS LIMITED**

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

**SUMMARY OF PRODUCT CHARACTERISTICS**

**4.5 Interaction with other medicinal products and other forms of interaction**

If the patient is taking amiodarone then chloroquine may increase the risk of cardiac arrhythmias including ventricular arrhythmias, bradycardias and cardiac conduction defect.

Concurrent use is contraindicated. Co-administration with other drugs that have antiarrhythmogenic properties, e.g. moxifloxacin, droperidol, may increase the risk of cardiac arrhythmias.

Antacids and adsorbents (e.g. kaolin) may reduce the absorption of chloroquine, so should be administered at least four hours apart.

Concomitant use of drugs such as multidrug and toxin extrusion protein (MATE1) inhibitors (e.g., ciprofloxacin, cimetidine, omeprazole, pyrimethamine) may impact the renal clearance of chloroquine, which could theoretically lead to increased levels of chloroquine and potentially overdose (see section 4.9). In addition, care should be taken when alkalization of urine occurs as this may reduce chloroquine renal excretion.

Chloroquine increases risk of convulsions with mefloquine (anti-malarial drug).

Chloroquine antagonises the anticonvulsant effect of antiepileptics.

Chloroquine may possibly increase the plasma concentration of digoxin.

When co-administered with ciclosporin, chloroquine increases plasma ciclosporin concentration resulting in increased risk of toxicity.

Chloroquine has been reported to reduce the bioavailability of praziquantel. Caution is advised during co-administration.

Chloroquine has the potential to increase symptoms of myasthenia gravis and thus diminish effect of neostigmine and pyridostigmine.

Concomitant administration of chloroquine with rabies vaccine may affect the antibody response.

Concomitant administration of chloroquine inactivates oral typhoid vaccine, so the vaccine should be completed at least three days before the first dose of chloroquine.

## BIORAJ PHARMACEUTICALS LIMITED

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

### SUMMARY OF PRODUCT CHARACTERISTICS

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

Should not be used during pregnancy unless, in the judgement of the physician, potential benefit outweighs the risk. When given at high doses throughout pregnancy it has been reported to give rise to foetal abnormalities including visual loss, ototoxicity and cochlea-vestibular dysfunction.

Malaria in pregnant women increase the risk of maternal death, miscarriages, still-births and low birth weight infants with the associated risk of neonatal death. Travel to malarious areas should be avoided during pregnancy but if this is not possible women should receive effective prophylaxis.

##### Breast-feeding

Malarivon Syrup is excreted in breast milk, although amounts are probably too small to be harmful when used for malaria prophylaxis but as a consequence, they are insufficient to protect the infant.

#### **4.7 Effects on ability to drive and use machines**

At start of treatment chloroquine has a temporary effect on visual accommodation, causing blurred and/or double vision. Therefore, patients should be advised that the product may affect their ability to drive or operate machinery.

#### **4.8 Undesirable effects**

The following CIOMS frequency rating is used when applicable:

Very common  $\geq 10\%$ ; Common  $\geq 1$  and  $< 10\%$ ; Uncommon  $\geq 0.1$  and  $< 1\%$ ; Rare  $\geq 0.01$  and  $< 0.1\%$ ; Very rare  $< 0.01\%$ ; Not known (frequency cannot be estimated from available data)

##### Cardiac disorders

- Uncommon: cardiomyopathy has been reported during long term therapy at high doses, which may result in cardiac failure and in some cases a fatal outcome.
- Rare: cardiac arrhythmias, including QT prolongation, torsade de pointes, ventricular tachycardia and ventricular fibrillation have been reported with therapeutic doses of chloroquine as well as with overdose. The risk is greater if chloroquine is administered at high doses. Fatal cases have been reported.

**BIORAJ PHARMACEUTICALS LIMITED**

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

**SUMMARY OF PRODUCT CHARACTERISTICS**

- Not known - hypotension.

Nervous system disorders

- Very common: headache

- Common: convulsions have been reported rarely (these may result from cerebral malaria).

- Uncommon: neuropathy

- Rare: polyneuropathy

- Not known: acute extrapyramidal disorders (such as dystonia, dyskinesia, tongue protrusion, torticollis).

Psychiatric disorders

- Very common: insomnia

- Common: depression

- Rare: psychiatric disorders such as anxiety, agitation, confusion, hallucinations, delirium

- Not known: suicidal behaviour

Eye disorders

- Common: transient blurred vision

- Rare: reversible corneal opacity, cases of retinopathy as well as cases of irreversible retinal damage have been reported during long term, high dose therapy.

- Not known: maculopathy and macular degeneration have been reported and may be irreversible, macular defects of colour vision, optic atrophy, scotomas, field defects, blindness and pigmented deposits, difficult in focusing, diplopia.

Gastro-intestinal disorders

- Very common: gastrointestinal disturbances such as nausea, vomiting, diarrhoea.

- Not known: abdominal cramps

Blood and lymphatic system disorders

- Rare: bone marrow depression, including aplastic anaemia, agranulocytosis, pancytopenia, thrombocytopenia, neutropenia

Hepatobiliary disorders

- Rare: changes in liver function, including hepatitis and abnormal liver function tests

- Immune system disorders

- Common: allergic and anaphylactic reactions, including angioedema

## BIORAJ PHARMACEUTICALS LIMITED

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

### SUMMARY OF PRODUCT CHARACTERISTICS

#### Ear and labyrinth disorders

- Uncommon: ototoxicity such as tinnitus, hypoacusis, nerve deafness.

#### Musculoskeletal and connective tissue disorders

- Uncommon: myopathy

#### Skin and subcutaneous tissue disorders

- Very common: pruritis,
- Common: skin eruptions, urticaria
- Uncommon: alopecia, bluish-black pigmentation of the nails and mucosae (long term use).
- Rare: exacerbation of psoriasis, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis
- Very rare: exfoliative dermatitis and similar desquamation-type events.
- Not known: depigmentation, photosensitivity, Drug Rash with Eosinophilia and Systemic Symptoms (DRESS syndrome)

#### Metabolism and nutrition disorders

- Not known: hypoglycaemia

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

#### **4.9 Overdose**

Chloroquine is highly toxic in overdose and children are particularly susceptible. The chief symptoms of overdosage include circulatory collapse due to a potent cardiotoxic effect, respiratory arrest and coma. Symptoms may progress rapidly after initial headache, drowsiness, visual disturbances, nausea and vomiting. Cardiac complications may occur without progressively deepening coma.

Death may result from circulatory or respiratory failure or cardiac arrhythmia. If there is no demonstrable cardiac output due to arrhythmias, asystole or electromechanical dissociation, external chest compression should be persisted with for as long as necessary, or until adrenaline and diazepam can be given (see below).

**BIORAJ PHARMACEUTICALS LIMITED**

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

**SUMMARY OF PRODUCT CHARACTERISTICS**

Gastric lavage should be carried out urgently (as soon as possible within 2 hours of the overdose), first protecting the airway and instituting artificial ventilation where necessary. There is a risk of cardiac arrest following aspiration of gastric contents in more serious cases. Activated charcoal left in the stomach may reduce absorption of any remaining chloroquine from the gut (minimum 5 times the suspected maximum dose of chloroquine phosphate). Circulatory status (with central venous pressure measurement), respiration, plasma electrolytes and blood gases should be monitored, with correction of hypokalaemia and acidosis if indicated. Cardiac arrhythmias should not be treated unless life threatening; drugs with quinidine-like effects should be avoided. Intravenous sodium bicarbonate 1-2mmol/kg over 15 minutes may be effective in conduction disturbances, and DC shock is indicated for ventricular tachycardia and ventricular fibrillation.

Early administration of the following has been shown to improve survival in cases of serious poisoning:

1. Adrenaline infusion 0.25micrograms/kg/min initially, with increments of 0.25micrograms/kg/min until adequate systolic blood pressure (more than 100mg/Hg) is restored; adrenaline reduces the effects of chloroquine on the heart through its inotropic and vasoconstrictor effects.
2. Diazepam infusion (2mg/kg over 30 minutes as a loading dose, followed by 1-2mg/kg/day for up to 2-4 days). Diazepam may minimise cardiotoxicity.

Acidification of the urine, haemodialysis, peritoneal dialysis or exchange transfusion have not been shown to be of value in treating chloroquine poisoning. Chloroquine is excreted very slowly, therefore cases of overdosage require observation for several days.

## BIORAJ PHARMACEUTICALS LIMITED

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

### SUMMARY OF PRODUCT CHARACTERISTICS

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antimalarials, Aminoquinolines, ATC code: P01B A01  
Chloroquine Phosphate is an antimalarial, active against susceptible strains of *Plasmodium falciparum*, *P.ovale*, *P. Vivax* and *P. Malariae*.

### 5.2 Pharmacokinetic properties

Chloroquine Phosphate is rapidly and almost completely absorbed from the gastro intestinal tract following oral administration. It is then widely distributed in the body tissues with the highest concentrations being found in kidneys, lungs, liver and spleen. In addition, it is also concentrated in melanin containing cells such as in the eyes and skin. It both crosses the placenta and is found in breast milk. Chloroquine remains in the system for a long period after discontinuation of therapy. Metabolism is mainly in the liver with elimination being via the urine.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Sucrose

Sodium benzoate

Citric acid

Sorbitol

Strawberry flavor

Tartrazine yellow

### 6.2 Incompatibilities

None known

### 6.3 Shelf life

36 months

### 6.4 Special precautions for storage

Store below 30°C. Keep away from the reach of children.

### 6.5 Nature and contents of container

Amber 60ml bottle with monocartoon and leaflet.

### 6.6 Special precautions for disposal and other handling

None.



**BIORAJ PHARMACEUTICALS LIMITED**

**BRAND NAME: BIORAQUINE SYRUP**

**GENERIC NAME: CHLOROQUINE PHOSPHATE**

**SUMMARY OF PRODUCT CHARACTERISTICS**

**7.0 MANUFACTURER**

*Bioraj Pharmaceuticals Limite*

*No 405 Kaiama Road, Ilorin*

*biorajpharmaceuticalltd@gmail.com*