

## **1. NAME OF THE MEDICINAL PRODUCT**

GLUCOVANCE 1000 mg/5 mg, film-coated tablets

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each film-coated tablet contains 1000 mg metformin hydrochloride, equivalent to 780 mg metformin, and 5 mg glibenclamide.

Excipient(s) with known effect: lactose.

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Film-coated tablet.

White to off-white oval-shaped, biconvex film-coated tablet with '1000' engraved on one side and '5' engraved on the other side.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

Treatment of type 2 diabetes in adults, as replacement for previous combination therapy with metformin and glibenclamide in patients whose glycaemia is stable and well-controlled.

### **4.2 Posology and method of administration**

#### Posology

Oral route.

For use in adults only.

#### **General:**

As for all hypoglycaemic agents, the dosage should be adapted according to the individual metabolic response (glycaemia, HbA1c).

*Adults with normal renal function (GFR  $\geq$  90 mL/min)*

#### **Initiation of treatment:**

Treatment should be initiated with a dose of the combination product equivalent to previous individual doses of metformin and glibenclamide; the dose being gradually increased depending on results on glycaemic parameters.

**Dose titration:**

The dosage should be adjusted every 2 weeks or longer, by increments of 1 tablet of metformin hydrochloride/glibenclamide 500 mg/2.5 mg, depending on glycaemia results.

A gradual increase in the dosage may aid gastrointestinal tolerance and prevent the onset of hypoglycaemia.

For patients already treated with a combination of metformin and glibenclamide, two tablets of metformin hydrochloride/glibenclamide 500 mg/2.5 mg can be replaced by one tablet of Glucovance 1000 mg/5 mg.

**Maximum daily recommended dose:**

The maximum daily recommended dose is 3 tablets of Glucovance 1000 mg/5 mg.

•

**Combination with insulin therapy:**

No clinical data are available on the concomitant use of this product with insulin therapy.

*Renal impairment*

A GFR should be assessed before initiation of treatment with metformin containing products and at least annually thereafter. In patients at an increased risk of further progression of renal impairment and in the elderly, renal function should be assessed more frequently, e.g. every 3-6 months.

The maximum daily dose of metformin should preferably be divided into 2-3 daily doses. Factors that may increase the risk of lactic acidosis (see 4.4) should be reviewed before considering initiation of metformin in patients with GFR < 60 mL/min.

If no adequate strength of Glucovance is available, individual mono-components should be used instead of the fixed dose combination.

GFR mL/min	Metformin	Glibenclamide
60-89	Maximum daily dose is 3000 mg.  Dose reduction may be considered in relation to declining renal function.	No dose reduction required.
45-59	Maximum daily dose is 2000 mg.  The starting dose is at most half of the maximum dose	Maximum daily dose is 10.5 mg.
30-44	Maximum daily dose is 1000 mg.  The starting dose is at most half of the maximum dose.	Maximum daily dose is 10.5 mg.  Initiation of therapy is not recommended due to the risk of hypoglycemia.
< 30	Metformin / glibenclamide are contraindicated	

### Geriatric population:

The dosage of Glucovance should be adjusted depending on renal function parameters (start with 1 tablet of Glucovance 500 mg/2.5 mg); regular checks on the renal function are necessary (see section 4.4).

Patients aged 65 years and older: starting and maintenance doses of glibenclamide must be carefully adjusted to reduce the risk of hypoglycaemia. Treatment should be started with the lowest available dose and increased gradually if necessary (see section 4.4).

### Paediatric population:

Glucovance is not recommended for use in children (see section 5.1).

### Method of administration

The dosage regimen depends on the individual posology:

- Once a day, in the morning at breakfast, for a dosage of 1 tablet/day,
- Twice a day, morning and evening, for a dosage of 2 or 4 tablets/day,
- Three times a day, morning, noon and evening, for a dosage of 3 tablets/day.

The tablets should be taken with meals. The dosage regimen should be adjusted according to the individual eating habits. However, any intake must be followed by a meal with a sufficiently high carbohydrate content to prevent the onset of hypoglycaemic episodes.

When Glucovance is co-administered with a bile acid sequestrant, it is recommended that Glucovance should be administered at least 4 hours prior to the bile acid sequestrant in order to minimize the risk of reduced absorption (see section 4.5).

## **4.3 Contraindications**

- hypersensitivity to metformin, glibenclamide or other sulphonylurea(s) and sulphonamide(s) or to any of the excipients listed in section 6.1;
- type 1 diabetes (insulin-dependent diabetes), diabetic pre-coma;
- any type of acute metabolic acidosis (such as lactic acidosis, diabetic ketoacidosis);
- severe renal failure (GFR < 30 mL/min);
- acute conditions with the potential to alter renal function such as: dehydration, severe infection, shock;
- disease which may cause tissue hypoxia (especially acute disease, or worsening of chronic disease) such as decompensated heart failure, respiratory failure, recent myocardial infarction, shock;
- hepatic insufficiency, acute alcohol intoxication, alcoholism;
- porphyria;
- lactation;
- in association with miconazole (see section 4.5).

## **4.4 Special warnings and precautions for use**

### **Lactic acidosis**

Lactic acidosis, a very rare but serious metabolic complication, most often occurs at acute worsening of renal function or cardiorespiratory illness or sepsis. Metformin accumulation occurs at acute worsening of renal function and increases the risk of lactic acidosis.

In case of dehydration (severe diarrhoea or vomiting, fever or reduced fluid intake), metformin should be temporarily discontinued and contact with a health care professional is recommended.

Medicinal products that can acutely impair renal function (such as antihypertensives, diuretics and NSAIDs) should be initiated with caution in metformin-treated patients. Other risk factors for lactic acidosis are excessive alcohol intake, hepatic insufficiency, inadequately controlled diabetes, ketosis, prolonged fasting and any conditions associated with hypoxia, as well as concomitant use of medicinal products that may cause lactic acidosis (see sections 4.3 and 4.5).

Patients and/or care-givers should be informed of the risk of lactic acidosis. Lactic acidosis is characterised by acidotic dyspnoea, abdominal pain, muscle cramps, asthenia and hypothermia followed by coma. In case of suspected symptoms, the patient should stop taking metformin and seek immediate medical attention. Diagnostic laboratory findings are decreased blood pH (< 7.35), increased plasma lactate levels (> 5 mmol/L) and an increased anion gap and lactate/pyruvate ratio.

### **Hypoglycaemia**

As it contains a sulphonylurea, Glucovance exposes the patient to a risk of onset of hypoglycaemic episodes. After treatment initiation, a progressive dose titration may prevent the onset of hypoglycaemia. This treatment should only be prescribed if the patient adheres to a regular meal schedule (including breakfast). It is important that carbohydrate intake is regular since the risk of hypoglycaemia is increased by a late meal, insufficient or unbalanced carbohydrate intakes. Hypoglycaemia is more likely to occur in case of energy-restricted diet, after intensive or prolonged exercise, when alcohol intake or during the administration of a combination of hypoglycaemic agents.

#### *Diagnosis:*

The symptoms of hypoglycaemia are: headache, hunger, nausea, vomiting, extreme tiredness, sleep disorder, restlessness, aggression, impaired concentration and reactions, depression, confusion, speech impediment, visual disturbances, trembling, paralysis and paraesthesia, dizziness, delirium, convulsions, somnolence, unconsciousness, superficial breathing and bradycardia. Due to a counterregulation caused by the hypoglycaemia sweating, fear, tachycardia, hypertension, palpitations, angina and arrhythmia can occur. These latter symptoms can be absent when the hypoglycaemia is developed slowly, in case of autonomic neuropathy or when the patients take beta-blocking agents, clonidine, reserpine, guanethidine or sympathomimetics.

#### *Management of hypoglycaemia:*

Moderate hypoglycaemic symptoms without loss of consciousness or neurological manifestations should be corrected by the immediate intake of sugar. An adjustment to the dosage and/or changes to meal patterns should be ensured. Severe hypoglycaemic reactions with coma, seizures or other neurological signs are also possible and constitute a medical emergency requiring immediate treatment with intravenous glucose once the cause is diagnosed or suspected, prior to prompt hospitalisation of the patient.

The careful selection of patients and dosage and adequate instructions for the patient are important to reduce the risk of hypoglycaemic episodes. If the patient encounters repeated episodes of hypoglycaemia, which are either severe or associated with unawareness of the situation, antidiabetic treatment options other than Glucovance should be taken into consideration.

#### Factors favouring hypoglycaemia:

- concomitant administration of alcohol, especially combined with fasting,
- refusal or (more particularly in elderly patients) inability of the patient to co-operate,
- malnutrition, irregular meals, missed meals, fasting or changes to diet,
- poor balance between physical exercise and carbohydrate intake,

- renal failure,
- severe liver failure,
- overdose of Glucovance,
- certain endocrine disturbances: thyroid insufficiency, pituitary and adrenal gland insufficiency,
- concomitant administration of certain other drugs (see section 4.5).

*Elderly patients:*

Age 65 years and older has been identified as a risk factor for hypoglycemia in patients treated with sulfonylureas. Hypoglycemia can be difficult to recognize in the elderly. Starting and maintenance doses of glibenclamide must be carefully adjusted to reduce the risk of hypoglycaemia (see section 4.2).

*Renal and hepatic failure:*

The pharmacokinetics and/or pharmacodynamics of Glucovance may be modified in patients with hepatic failure or severe renal failure. If hypoglycaemia occurs in such patients, it may be prolonged, and appropriate treatment must be initiated.

*Patient information:*

The risks of hypoglycaemia, its symptoms and its treatment, as well as its predisposing conditions, must be explained to the patient and his or her family. Similarly, the risk of lactic acidosis must be considered in the event of non-specific signs such as muscle cramps accompanied by digestive disorders, abdominal pain and severe asthenia, dyspnoea attributed to acidose, hypothermia and coma.

In particular, the patient should be informed of the importance of adhering to a diet, following a programme of regular physical exercise and making regular checks on glycaemia.

**Blood sugar imbalance**

In case of surgery or any other cause of diabetic decompensation, temporary insulin therapy should be envisaged instead of this treatment.

The symptoms of hyperglycaemia are: increased urinating, raging thirst and a dry skin.

**Cardiac function**

Patients with heart failure are more at risk of hypoxia and renal insufficiency. In patients with stable chronic heart failure, Glucovance may be used with a regular monitoring of cardiac and renal function.

For patients with acute and unstable heart failure, Glucovance is contraindicated (see section 4.3).

**Renal function**

GFR should be assessed before treatment initiation and regularly thereafter, see section 4.2. Metformin is contraindicated in patients with GFR < 30 mL/min and should be temporarily discontinued in the presence of conditions that alter renal function, see section 4.3.

**Administration of iodinated contrast agents**

Intravascular administration of iodinated contrast agents may lead to contrast induced nephropathy, resulting in metformin accumulation and an increased risk of lactic acidosis. Glucovance should be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours

after, provided that renal function has been re-evaluated and found to be stable, see sections 4.2 and 4.5.

### **Concomitant use of glibenclamide with other medicinal products**

The concomitant use of glibenclamide with alcohol, phenylbutazone or danazol is not recommended (see section 4.5).

### **Surgery**

Glucovance must be discontinued at the time of surgery under general, spinal or epidural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and provided that renal function has been re-evaluated and found to be stable.

### **Other precautions**

All patients should continue their diet, with a regular distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet.

Regular physical exercise is as necessary as taking Glucovance.

The usual laboratory tests for diabetes monitoring (glycaemia, HbA1c) should be performed regularly,

Metformin may reduce vitamin B12 serum levels. The risk of low vitamin B12 levels increases with increasing metformin dose, treatment duration, and/or in patients with risk factors known to cause vitamin B12 deficiency. In case of suspicion of vitamin B12 deficiency (such as anemia or neuropathy), vitamin B12 serum levels should be monitored. Periodic vitamin B12 monitoring could be necessary in patients with risk factors for vitamin B12 deficiency. Metformin therapy should be continued for as long as it is tolerated and not contra-indicated and appropriate corrective treatment for vitamin B12 deficiency provided in line with current clinical guidelines.

Treatment of patients with G6PD-deficiency with sulphonylurea agents can lead to haemolytic anaemia. Since glibenclamide belongs to the chemical class of sulphonylurea drugs, caution is recommended when using Glucovance in patients with G6PD-deficiency and a non-sulphonylurea alternative may be considered.

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, i.e. essentially 'sodium-free'.

## 4.5 Interaction with other medicinal products and other forms of interaction

### Contraindicated combination

#### Related to glibenclamide

*Miconazole (systemic route, oromucosal gel)*: Increase in the hypoglycaemic effect with possible onset of hypoglycaemic manifestations, or even coma (see section 4.3).

### Combinations not recommended

#### Related to sulphonylurea(s)

##### *Alcohol:*

Antabuse effect (intolerance to alcohol), notably for chlorpropamide, glibenclamide, glipizide, tolbutamide.

Increase of the hypoglycaemic reaction (inhibition of compensation reactions), which may facilitate the onset of a hypoglycaemic coma (see section 4.4).

Avoid consumption of alcohol and alcohol-containing medications.

##### *Phenylbutazone (systemic route):*

Increase in the hypoglycaemic effect of sulphonylurea(s) (displacement of sulphonylurea(s) from protein-binding sites and/or decrease in their elimination). Preferably use another anti-inflammatory agent exhibiting fewer interactions, or else warn the patient and step up self-monitoring; if necessary, adjust the dosage during treatment with the anti-inflammatory agent and after its withdrawal.

#### Related to all antidiabetic agents

##### *Danazol:*

If the combination cannot be avoided, warn the patient and step up self-monitoring of blood glucose. Possibly adjust the dosage of the antidiabetic treatment during treatment with danazol and after its withdrawal.

#### Related to metformin

##### *Alcohol:*

Alcohol intoxication is associated with an increased risk of lactic acidosis, particularly in cases of fasting, malnutrition and or hepatic impairment.

##### *Iodinated contrast agents:*

Glucovance must be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable, see sections 4.2 and 4.4.

### Combinations requiring precautions

#### Related to all antidiabetic agents

##### *Chlorpromazine:*

At high dosages (100 mg per day of chlorpromazine), elevation in blood glucose (reduction in release of insulin).

Precaution for use: warn the patient and step up self-monitoring of blood glucose. Possibly adjust the dosage of the antidiabetic treatment during treatment with the neuroleptic and after its withdrawal.

*Corticosteroids (glucocorticoids) and tetracosactides (systemic and local routes):*

Elevation in blood glucose, sometimes accompanied by ketosis (decreased carbohydrate tolerance with corticosteroids).

Precaution for use: warn the patient and step up self-monitoring of blood glucose. Possibly adjust the dosage of the antidiabetic during treatment with corticosteroids and after their withdrawal.

*β<sub>2</sub>-agonists:*

Elevation in blood glucose due to the β<sub>2</sub>-agonists.

Precaution for use: warn the patient, step up blood glucose monitoring and possibly transfer to insulin therapy.

### Related to metformin

Some medicinal products can adversely affect renal function which may increase the risk of lactic acidosis, e.g. NSAIDs, including selective cyclo-oxygenase (COX) II inhibitors, ACE inhibitors, angiotensin II receptor antagonists and diuretics, especially loop diuretics. When starting or using such products in combination with metformin, close monitoring of renal function is necessary.

*Organic cation transporters (OCT):*

Metformin is a substrate of both transporters OCT1 and OCT2.

Co-administration of metformin with

- Inhibitors of OCT1 (such as verapamil) may reduce efficacy.
- Inducers of OCT1 (such as rifampicin) may increase gastrointestinal absorption and efficacy of metformin.
- Inhibitors of OCT2 (such as cimetidine, dolutegravir, ranolazine, trimethoprim, vandetanib, isavuconazole) may decrease the renal elimination of metformin and thus lead to an increase in metformin plasma concentration.
- Inhibitors of both OCT1 and OCT2 (such as crizotinib, olaparib) may alter efficacy and renal elimination of metformin.

Caution is therefore advised, especially in patients with renal impairment, when these drugs are co-administered with metformin, as metformin plasma concentration may increase. If needed, dose adjustment of metformin may be considered as OCT inhibitors/inducers may alter the efficacy of metformin.

### Related to glibenclamide

*Beta-blockers:*

All beta-blockers mask some of the symptoms of hypoglycaemia: palpitations and tachycardia; Most non-cardioselective beta-blockers increase the incidence and severity of hypoglycaemia.

Warn the patient and step up blood glucose self-monitoring, especially at the start of treatment.

*Angiotensin converting enzyme inhibitors (e.g. captopril, enalapril):*

ACE inhibitors may decrease the blood glucose levels. If necessary, adjust the dosage of Glucovance during therapy with an ACE inhibitor and upon its discontinuation.

*Fluconazole:*

Increase in the half-life of sulphonylurea with possible onset of hypoglycaemic manifestations.

Warn the patient and step up blood glucose self-monitoring, and possibly adjust the dosage of the antidiabetic treatment during treatment with fluconazole and after its withdrawal.

*Bosentan:*

Risk of decreased hypoglycaemic effect of glibenclamide because bosentan reduces the plasma concentration of glibenclamide. An increased risk of liver enzyme elevations was reported in patients receiving glibenclamide concomitantly with bosentan.

Warn the patient, set-up monitoring of glycaemia and liver enzymes and adjust the dosage of the antidiabetic treatment if necessary.

*Bile acid sequestrants:*

When co-administered simultaneously the plasma concentration of glibenclamide is reduced which may lead to a reduced hypoglycaemic effect. This effect was not observed if glibenclamide is given in a certain period of time before taking the other medicine. It is recommended that Glucovance should be administered at least 4 hours prior a bile acid sequestrant.

**Other interaction: combination to be taken into account:**

Related to glibenclamide

*Desmopressin:*

Reduction in antidiuretic activity.

**4.6 Fertility, pregnancy and lactation**

**Pregnancy**

No preclinical and clinical data on exposed pregnancies are available for Glucovance.

Risk related to diabetes

When uncontrolled, diabetes (gestational or permanent) gives rise to an increase in congenital abnormalities and perinatal mortality. Diabetes must be controlled as far as possible during the period of conception in order to reduce the risk of congenital abnormalities.

Risk related to metformin (see section 5.3)

Animal studies do not indicate harmful effects with respect to pregnancy, embryonic or fetal development, parturition or postnatal development.

A limited amount of data from the use of metformin in pregnant women does not indicate an increased risk of congenital abnormalities.

Risk related to glibenclamide (see section 5.3)

Studies in animals have shown no evidence of teratogenic activity. In the absence of a teratogenic effect in animals, foetal malformation in humans is not to be expected since to date, substances known to cause malformation in humans have proved to be teratogenic in well-conducted animal studies in two species.

In clinical practice, there are currently no relevant data on which to base an evaluation of potential malformation or fetotoxicity due to glibenclamide when administered during pregnancy.

Management

Adequate blood glucose control allows pregnancy to proceed normally in this category of patients. Glucovance must not be used for the treatment of diabetes during pregnancy.

It is imperative that insulin be used to achieve adequate blood glucose control. It is recommended that the patient be transferred from oral antidiabetic therapy to insulin as soon as she plans to become pregnant or if pregnancy is exposed to this medicinal product. Neonatal blood glucose monitoring is recommended.

## **Breast-feeding**

Metformin is excreted into human breast milk. No adverse effects were observed in breastfed newborns/infants of mothers treated with metformin alone. However, in humans, in the absence of data concerning passage of glibenclamide into breast milk, and in view of the risk of neonatal hypoglycaemia, this medicinal product is contraindicated in the event of breast-feeding.

## **Fertility**

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose based on body surface area comparisons.

Fertility of male or female rats was unaffected by glibenclamide when administered orally at dose of 100 and 300 mg/kg/day.

### **4.7 Effects on ability to drive and use machines**

Patients should be alerted to the symptoms of hypoglycaemia and should be advised to exercise caution when driving or using machines.

### **4.8 Undesirable effects**

During treatment initiation, the most common adverse reactions are nausea, vomiting, diarrhoea, abdominal pain and loss of appetite which resolve spontaneously in most cases. To prevent them, it is recommended to take Glucovance in 2 or 3 daily doses and to increase slowly the doses. Transient visual disturbances may occur at the start of treatment due to a decrease in glycaemia levels.

The following adverse reactions may occur under treatment with Glucovance. Frequencies are defined as follows: very common:  $\geq 1/10$ ; common  $\geq 1/100$ ,  $< 1/10$ ; uncommon:  $\geq 1/1,000$ ,  $< 1/100$ ; rare  $\geq 1/10,000$ ,  $< 1/1,000$ ; very rare  $< 1/10,000$ .

#### **Blood and lymphatic system disorders:**

These are reversible upon treatment discontinuation.

*Rare:* Leukopenia, thrombocytopenia.

*Very rare:* Agranulocytosis, haemolytic anaemia, bone marrow aplasia and pancytopenia.

#### **Metabolism and nutrition disorders:**

Hypoglycaemia (see section 4.4).

*Uncommon:* Crises of hepatic porphyria and porphyria cutanea.

*Common:* Vitamin B12 decrease/deficiency  
(see section 4.4).

*Very rare:* Lactic acidosis (see section 4.4).  
Disulfiram-like reaction with alcohol intake.

#### **Nervous system disorders:**

*Common:* Taste disturbance.

#### **Eye disorders:**

Transient visual disturbances may occur at the start of treatment due to a decrease in glycaemia levels.

#### **Gastrointestinal disorders:**

*Very common:* Gastrointestinal disorders such as nausea, vomiting, diarrhoea, abdominal pain and loss of appetite. These undesirable effects occur more frequently during treatment initiation and resolve spontaneously in most cases. To prevent them, it is recommended that Glucovance be taken in 2 or 3 daily doses. A slow increase of the dose may also improve gastrointestinal tolerability.

#### **Skin and subcutaneous tissue disorders:**

A cross reactivity to sulphonamide(s) and their derivatives may occur.

*Rare:* Skin reactions such as pruritus, urticaria, maculopapular rash.

*Very rare:* Cutaneous or visceral allergic angiitis, erythema multiforme, exfoliative dermatitis, photosensitization, urticaria evolving to shock.

#### **Hepatobiliary disorders:**

*Very rare:* Liver function test abnormalities or hepatitis requiring treatment discontinuation

#### **Investigations:**

*Uncommon:* Average to moderate elevations in serum urea and creatinine concentrations.

*Very rare:* Hyponatremia.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via **the national reporting system**.

### **4.9 Overdose**

Overdose may precipitate hypoglycaemia due to the presence of the sulphonylurea (see section 4.4).

High overdose or the existence of concomitant risk factors may lead to lactic acidosis due to the presence of metformin (see section 4.4). Lactic acidosis is a medical emergency and must be treated in hospital. The most effective treatment is to remove lactate and metformin by haemodialysis.

The plasma clearance of glibenclamide may be prolonged in patients suffering from liver disease. Since glibenclamide is extensively bound to proteins, it is not eliminated by dialysis.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Biguanides and sulphonamide(s) in combination. ATC code: A10BD02

Metformin is a biguanide with antihyperglycaemic effects, lowering both basal and postprandial plasma glucose. It does not stimulate insulin secretion and therefore does not produce hypoglycaemia.

Metformin may act via 3 mechanisms:

- (1) by reducing hepatic glucose production by inhibiting gluconeogenesis and glycogenolysis
- (2) in muscle, by increasing insulin sensitivity, improving peripheral glucose uptake and utilisation
- (3) and by delaying intestinal glucose absorption.

Metformin stimulates intracellular glycogen synthesis by acting on glycogen synthase. Metformin increases the transport capacity of all types of membrane glucose transporters (GLUT).

In humans, independently of its action on glycaemia, metformin has favourable effects on lipid metabolism. This has been shown at therapeutic doses in controlled, medium-term or long-term clinical studies: metformin reduces total cholesterol, LDL-cholesterol and triglyceride levels. In clinical trials conducted so far with combination therapy with metformin and glibenclamide, these favourable effects on lipid metabolism have not been shown.

Glibenclamide is a second generation sulphonylurea with a medium half-life: it causes acute lowering of blood glucose by stimulating the release of insulin by the pancreas, this effect being dependent on the presence of functioning beta cells in the islets of Langerhans.

The stimulation of insulin secretion by glibenclamide in response to a meal is of major importance.

The administration of glibenclamide to diabetics induces an increase in the postprandial insulin-stimulating response. The increased postprandial responses in insulin and C-peptide secretion persist after at least 6 months of treatment.

Metformin and glibenclamide have different mechanisms and sites of action, but their action is complementary. Glibenclamide stimulates the pancreas to secrete insulin, while metformin reduces cell resistance to insulin by acting on peripheral (skeletal muscle) and hepatic sensitivity to insulin.

Results from controlled, double blind clinical trials versus reference products in the treatment of type 2 diabetes inadequately controlled by monotherapy with metformin or glibenclamide combined with diet and exercise, have demonstrated that the combination had an additive effect on glucose regulation.

#### **Paediatric population:**

In a 26-week, active controlled, double-blind, clinical study performed in 167 paediatric patients aged 9 to 16 years with type 2 diabetes not adequately controlled with diet and exercise, with or without an oral antidiabetic treatment, a fixed combination of metformin hydrochloride 250 mg and glibenclamide 1.25 mg was not shown more effective to either metformin hydrochloride or glibenclamide in reducing HbA1c from baseline. Therefore, Glucovance should not be used in paediatric patients.

## **5.2 Pharmacokinetic properties**

### **Related to the combination**

The bioavailability of metformin and glibenclamide in the combination is similar to that noted when one tablet of metformin and one tablet of glibenclamide are taken simultaneously. The bioavailability of metformin in the combination is unaffected by the ingestion of food. The bioavailability of glibenclamide in the combination is unaffected by the ingestion of food, but the absorption speed of glibenclamide is increased by eating.

### **Related to metformin**

#### **Absorption:**

After an oral dose of metformin tablet, maximum plasma concentration ( $C_{max}$ ) is reached in approximately 2.5 hours ( $t_{max}$ ). Absolute bioavailability of a 500 mg or 850 mg metformin tablet is approximately 50-60% in healthy subjects. After an oral dose, the non-absorbed fraction recovered in faeces was 20-30%.

After oral administration, metformin absorption is saturable and incomplete. It is assumed that the pharmacokinetics of metformin absorption is non-linear. At the usual metformin doses and dosing schedules, steady state plasma concentrations are reached within 24 to 48 hours and are generally less than 1 µg/ml. In controlled clinical trials, maximum metformin plasma levels ( $C_{max}$ ) did not exceed 5 µg/ml, even at maximum doses.

Distribution:

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears at approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean volume of distribution  $V_d$  ranged from 63 to 276 l.

Biotransformation:

Metformin is excreted unchanged in the urine. No metabolites have been identified in humans.

Elimination:

Renal clearance of metformin is  $> 400$  ml/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours.

When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

**Related to glibenclamide**

Absorption:

Glibenclamide is very readily absorbed ( $> 95\%$ ) following oral administration. The peak plasma concentration is reached in about 4 hours.

Distribution:

Glibenclamide is extensively bound to plasma albumin (99%), which may account for certain drug interactions.

Biotransformation:

Glibenclamide is completely metabolised in the liver to two metabolites. Hepatocellular failure decreases glibenclamide metabolism and appreciably slows down its excretion.

Elimination:

Glibenclamide is excreted in the form of metabolites via biliary route (60%) and urine (40%), elimination being complete within 45 to 72 hours. Its terminal elimination half-life is 4 to 11 hours.

Biliary excretion of the metabolites increases in cases of renal insufficiency, according to the severity of renal impairment until a creatinine clearance at 30 ml/min. Thus, glibenclamide elimination is unaffected by renal insufficiency as long as the creatinine clearance remains above 30 ml/min.

## **Paediatric population**

There were no differences in pharmacokinetics of glibenclamide and metformin between paediatric patients and weight-and gender-matched healthy adults.

### **5.3 Preclinical safety data**

No preclinical studies have been performed on the combination product. Preclinical evaluation of the constituents metformin and glibenclamide revealed no special hazard for humans based on conventional studies of repeated dose toxicity, genotoxicity and carcinogenic potential.

Animal studies on metformin and glibenclamide do not indicate direct or indirect harmful effects with respect to pregnancy, embryonic/ foetal development, parturition or postnatal development. (see section 4.6).

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Microcrystalline cellulose  
Sodium croscarmellose  
Povidone K30  
Magnesium stearate

#### Film-coating

Opadry II OY-L-28900 white (lactose monohydrate, hypromellose, titanium dioxide (E171), macrogol 4000).

### **6.2 Incompatibilities**

Not applicable

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

30, 60 or 90 tablets in clear blister (PVC/Aluminium).  
Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

Any unused product or waste material should be disposed of in accordance with local requirements.

**7. DATE OF REVISION OF THE TEXT**

June-2022

