

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

Gentamicin Eye/Ear Drops 0.3%, 10ml

Submitted by

**PHARMAX INDIA PVT. LTD.,
9, KURLA INDUSTRIAL ESTATE, NETAJI PALKAR ROAD,
GHATKOPAR (WEST), MAHARASHTRA, MUMBAI- 400 086, INDIA.**

1.3 Product Information

1.3.1 Summary of Product Characteristics (SmPC)

1. Name of the medicinal product

Gentamicin Eye/Ear Drops 0.3% 10ml

2. Qualitative and quantitative composition

Each 10ml contains: Gentamicin Sulfate Equivalent to 30mg gentamicin (0.3% w/v).
For the full list of excipients, see section 6.1.

3. Pharmaceutical form

Eye/ear drops

A clear, colorless to slightly yellow solution.

4. Clinical particulars

4.1 Therapeutic indications

Gentamicin eye/ear drops are indicated in adults and children:

1. For the treatment of superficial eye and ear infections caused by organisms sensitive to gentamicin.
2. For prophylaxis against infection in trauma of the eye or ear

4.2 Posology and method of administration

Posology

Adults, the elderly and the paediatric population

Eyes: 1 or 2 drops should be instilled in the affected eye up to six times a day, or more frequently if required. (Severe infections may require 1 or 2 drops every fifteen to twenty minutes initially, reducing the frequency of instillation gradually as the infection is controlled).

Ears: The area should be cleaned and 2 - 3 drops instilled in the affected ear three to four times a day and at night, or more frequently if required.

Method of administration

Auricular and ocular use.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Myasthenia gravis.

Known or suspected perforation of the ear drum.

4.4 Special warnings and precautions for use

Long-term continuous topical therapy should be avoided. Prolonged use may lead to

skin sensitisation and the emergence of resistant organisms. Cross sensitivity with other aminoglycoside antibiotics may occur.

In severe infections, topical use of gentamicin should be supplemented with appropriate systemic antibiotic treatment.

Gentamicin may cause irreversible partial or total deafness when given systemically or when applied topically to open wounds or damaged skin. This effect is dose-related and is enhanced by renal and/or hepatic impairment and is more likely in the elderly. The condition of the ear drum must always be checked before this medicinal product is prescribed. The medicinal product must not be used if the integrity of the ear drum cannot be guaranteed.

Irreversible toxic effects may result from direct contact of gentamicin with the middle and inner ear. The benefits of gentamicin therapy should be considered against the risk of infection itself causing hearing loss.

Contact lenses should be removed during the period of treatment of ocular infections. Serious adverse reactions including neurotoxicity, ototoxicity and nephrotoxicity have occurred in patients receiving systemic gentamicin therapy. Although these effects have not been reported following topical otic use of gentamicin, caution is advised when used concomitantly with systemic aminoglycosides.

Patients should be monitored in case of prolonged use.

4.5 Interaction with other medicinal products and other forms of interaction

Potent diuretics such as ethacrynic acid and furosemide are believed to enhance any risk of ototoxicity whilst amphotericin B, cisplatin and cyclosporine and cephalosporins are potential enhancers of nephrotoxicity.

Concurrent use with other potentially nephrotoxic or ototoxic drugs should be avoided unless considered essential by the physician.

Neuromuscular blockade and respiratory paralysis have been reported in patients from the administration of aminoglycosides to patients who have received curare-type muscle relaxants during anaesthesia.

4.6 Pregnancy and lactation

Pregnancy

Safety for use in pregnancy has not been established. Gentamicin should only be used in pregnancy when considered essential by the physician, after careful assessment of the potential risks and benefits.

Breast-feeding

Safety for use in lactation has not been established. In the absence of gastrointestinal inflammation the amount of gentamicin ingested from the milk is unlikely to result in significant blood levels in breast-fed infants. Gentamicin should only be used in lactation when considered essential by the physician, after careful assessment of the potential risks and benefits.

4.7 Effects on ability to drive and use machines

Patients should be advised that the use of Gentamicin eye/ear drops in the eye may

cause transient blurring of vision. If affected, patients should not drive or operate machinery until vision has cleared.

4.8 Undesirable effects

The following adverse reactions identified from post-marketing experience are included in the table below. Adverse drug reactions are classified by system organ classes using the following frequency:

Not known: frequency cannot be estimated from the available data

System organ class	Frequency	Adverse reactions
Eye disorders	Not known	- local sensitivity: - blurred vision - eye irritation - burning sensation - stinging sensation - itching (eye pruritus)
Ear and labyrinth disorders	Not known	- local sensitivity - ototoxicity - vestibular disorder: - hearing loss
Skin and subcutaneous tissue disorders	Not known	- irritation - burning sensation - stinging - itching (pruritus): - dermatitis.
Renal and urinary disorders	Not known	- nephrotoxicity* - acute renal failure

*Gentamicin may cause nephrotoxicity when given systemically. However, it is likely that systemic absorption following topical administration does not constitute a comparable risk.

In the event of irritation, sensitisation or super-infection, treatment should be discontinued and appropriate therapy instituted.

Very rarely, cases of dizziness, tinnitus, hypersensitivity and allergic reaction have been reported.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions.

4.9 Overdose

Symptoms

The oral ingestion of the contents of one bottle is unlikely to cause any significant adverse effect.

Management

Haemodialysis and peritoneal dialysis will aid the removal from blood but the former is probably more efficient. Calcium salts given intravenously have been used to counter the neuromuscular blockade cause by gentamicin.

5. Pharmacological properties

5.1 Pharmacodynamic properties

ATC code: S03AA06

Pharmacotherapeutic group: Antiinfectives

Mechanism of action

Gentamicin is a mixture of antibiotic substances produced by the growth of micromonospora purpurea. It is a bactericidal antibiotic which acts by inhibiting protein synthesis.

It has greater antibacterial activity than streptomycin, neomycin or kanamycin.

Gentamicin exerts a number of effects on cells of susceptible bacteria. It affects the integrity of the plasma membrane and the metabolism of RNA, but it's most important effect is inhibition of protein synthesis at the level of the 30s ribosomal subunit.

5.2 Pharmacokinetic properties

Absorption

Topical application of gentamicin can result in some systemic absorption. Treatment of large areas can result in plasma concentrations of up to 1 microgram/ml.

Gentamicin is 70-85% bound to plasma albumin following administration.

Effective plasma concentration is 4 - 8 µg/ml

The volume of distribution (V_D) is 0.31/kg

Elimination

> 90% Gentamicin is excreted unchanged in the urine by glomerular filtration.

$T_{1/2}$ = 2 - 3 hours in individuals with normal kidney function, but can be increased in cases of renal insufficiency.

The elimination rate constant is;

0.02 Hr⁻¹ for anuric patients*

0.30 Hr⁻¹ normal

*Therefore in those with anuria care must be exercised.

5.3 Preclinical safety data

Not relevant.

6. Pharmaceutical particulars

6.1 List of excipients

Sodium sulfite
Sodium chloride
Sodium dihydrogen phosphate
Disodium hydrogen phosphate
Ethyl parahydroxybenzoate
Water for injection

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Unopened: 3 years.

After the container is opened for the first time: 4 weeks.

6.4 Special precautions for storage

Store below 30°C. Do not freeze. Protect from light.

6.5 Nature and contents of container

10ml opaque LDPE bottles with closures comprising polythene plug droppers and HDPE tamper evident caps.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. Manufactured by

PHARMAX INDIA PVT. LTD.,
9, KURLA INDUSTRIAL ESTATE,
NETAJI PALKAR ROAD,
GHATKOPAR
(WEST), MAHARASHTRA,
MUMBAI- 400 086, INDIA

8. Marketing authorisation holder

KATARI PHARMACEUTICAL
CO. LTD., NO. G.G. 5, WARRI
STREET, KADUNA, KADUNA
STATE, NIGERIA.

9. Date of first Authorization /renewal of the authorization: ---

10. Date of revision of text: July 2023