



<b>Brand Name : SPEXACIN-OZ</b>	
<b>Generic Name : Ofloxacin USP 200 mg and Ornidazole 500 mg Tablets</b>	<b>2018</b>
<b>Module 1 : Administrative Information</b>	
<b>1.3 : Product information</b>	<b>Confidential</b>
<b>1.3.1 : Summary of Product Characteristics (SmPC)</b>	

## 1.3.1 Summary of Product Characteristics (SmPC)

### 1- Name of the Medicinal Product:

#### 1.1 Product Name

**-Generic Name or International Non-Proprietary Name (INN)**

OFLOXACIN USP 200 MG AND ORNIDAZOLE 500 MG TABLETS

**-Brand Name**

SPEXACIN-OZ

#### 1.2 Dosage Strength

Each Film coated tablet contains:

Ofloxacin USP.....200 mg

Ornidazole.....500 mg

Excipients.....q.s.

Colour: Sunset yellow FCF, Tartrazine, Titanium Dioxide BP

#### 1.3 Dosage Form

Film coated Tablets

### 2- Quality and Quantitative Composition:

#### 2.1 Qualitative Declaration

Each Film coated tablet contains:

Ofloxacin USP.....200 mg

Ornidazole.....500 mg

Excipients.....q.s.

Colour: Sunset yellow FCF, Tartrazine, Titanium Dioxide BP

RA EXECUTIVE

Prepared By

46 of 77

Q.A.MANAGER

Approved By

**2.2 Quantitative Declaration****Composition:**

Batch Size - 1, 00,000 Tablets.

Sr. No	Ingredients	Labeled Claim (mg)	Specification	mg/Tab	Qty/Batch 1 lac in Kg	Function
<b>SIFTING/MIXING</b>						
1	Ofloxacin	200.00	USP	200.000	20.000	Active
2	Ornidazole	500.00	IH	500.000	50.000	Active
3	Maize Starch	---	BP	101.000	10.100	Diluent
4	Dibasic Calcium Phosphate	---	BP	60.000	6.000	Diluent
5	Cross Carmellose Sodium sodium (vivasole)	---	BP	6.000	0.600	Diluent
<b>PASTE PREPRATION</b>						
6	Maize Starch	---	BP	33.000	3.300	Diluent
7	Gelatine	---	BP	5.000	0.500	Preservative
8	Purified Water	---	IH	q.s	q.s	Binder Solvent
<b>LUBRICATON</b>						
9	Magnesium Stearate	---	BP	10.000	1.000	Lubricant
10	Talcum	---	BP	10.000	1.000	Lubricant
11	Cross Carmellose Sodium (vivasole)	---	BP	15.000	1.500	Lubricant
12	Sodium Starch Glycolate	---	BP	10.000	1.000	Disintegrant
<b>Avg. Weight of Uncoated tablet</b>				<b>950.00 mg</b>	<b>Limit: 950 mg<math>\pm</math> 5.0%</b>	
13	H.P.M.C 15	---	BP	13.500	1.350	Coating Polymer
14	PEG 6000	---	BP	1.000	0.100	Coating Polymer
15	Talcum	---	BP	3.000	0.300	Lubricant
16	Titanium Dioxide	---	BP	0.200	0.020	Opacifier
17	Lake of Tartrazine	---	IH	0.750	0.075	Colour
18	Methylene Chloride (DCM)*	---	BP	162.000	16.200	Coating Solvent

RA EXECUTIVE

Prepared By

Q.A.MANAGER

Approved By



19	Iso Propyl Alcohol*	---	BP	108.000	10.800	Coating Solvent
20	Sunset Yellow Lake	---	IH	1.750	0.175	Colour
Avg. Weight of Film coated tablet				970.20 mg	Limit: 970.20 $\pm$ 5.0%	

**NOTE:** Active material is to be calculated on Assay / Potency basis.

USP = United States Pharmacopoeia,

BP = British Pharmacopoeia.

\* Does not appear in the finished Product.

### 3- Pharmaceutical Form:

Orange coloured capsule shaped film coated tablet having break line on one side and other side plain.

### 4- Clinical Particulars:

#### 4.1 Therapeutic indications.

Ofloxacin Ornidazole Tablet is a medicine that is used for the treatment of Bacterial infections, Urinary tract infections, Bacterial infections by inflammation of the peritoneum, Inflammatory discharge from the urethra or vagina, Eye and ear infection, Bacterial infection and other conditions.

The complete list of uses and indications for Ofloxacin Ornidazole Tablet is as follows:

- Bacterial infections
- Urinary tract infections
- Bacterial infections by inflammation of the peritoneum
- Inflammatory discharge from the urethra or vagina
- Eye and ear infection
- Bacterial infection
- Typhoid fever
- Skin infections
- Sexually transmitted infections
- Protozoan infections
- Respiratory infections
- Typhoid
- Infections of the skin

RA EXECUTIVE

Prepared By

48 of 77

Q.A.MANAGER

Approved By



- Vaginal infections
- Soft tissue infections
- Infections during surgical procedures
- Vaginal infection
- Tuberculosis
- Skin infection
- Infections of the urinary tract
- Infections of vagina
- Infection of respiratory tract
- Phthisis
- Vagina infection
- Sexually transmitted infection
- Infection of the vagina
- Respiratory infection
- Urinary tract infection

Ofloxacin Ornidazole Tablet may also be used for purposes not listed here.

#### 4.2 Posology and method of administration

Ofloxacin and ornidazole tablet dosage and duration of treatment are depended on bacteria sensitivity, and infection kind and severity. The average dose for adults is 1-2 tablets two times per day during 7-10 days. The treatment should be prolonged not less than 3 days after the disappearing of clinical symptoms.

In general doctors prefer the combination of Ofloxacin & ornidazole in the dosage of one tablet twice daily for 5-10 days.

- In 1. Diarrhea& Dysentery,
2. Gastroenteritis,
3. Lower respiratory & Urinary tract infection & pelvic inflammatory diseases,
4. Septic abortion,
5. Puerperal sepsis
6. Post-surgical infection,
7. Skin & soft tissue infection,
8. ENT infections,

RA EXECUTIVE

Prepared By

49 of 77

Q.A.MANAGER

Approved By



## 9. Oro-dental Infections.

### Mixed Amoebiasis

Adults: 1 tab twice daily for 5-7 days; Children: ½ tab once daily for 5 to 10 days

### Mixed Amoebic dysentery

Adults: 3 tablets once daily for 3 days; Children: 1 tablet once daily for 3 days

### Mixed Giardiasis:

Adults: 3 tablets once daily for 1-2 days; Children: 1 tablet for 2 days

### Trichomoniasis

Adults: 3 tablets once or 1 tablet twice daily for 5 days. Sexual partner should be simultaneously treated.

### Bacterial vaginosis and STD

Adults: 3 tablets once or 1 tablet once daily for 5-7 days

### Dental Infections

Initiate oral therapy as soon as possible after I.V. infusion in surgical conditions;

Adults: 1 tablet twice daily for 5 to 10 days; Children: ½ tablet twice daily

## 4.3 **Contraindications**

The drug is contraindicated in patients with known hypersensitivity to this product or any of its ingredients. It is not advocated during the first trimester of pregnancy and in those with history of tendinitis or tendon rupture following use of quinolones. Because of the potential for serious adverse reactions in the nursing infant, the drug must be either stopped or discontinued during lactation for at least 3 days, depending on the importance of the drug to the lactating mother.

## 4.4 **Special warning and precautions for use**

If you use other drugs or over the counter products at the same time, the effects of Ofloxacin Ornidazole Tablet may change. This may increase your risk for side-effects or cause your drug not to work properly. Tell your doctor about all the drugs, vitamins, and herbal supplements you are using, so that you doctor can help you prevent or manage drug interactions. Ofloxacin Ornidazole Tablet may interact with the following drugs and products:

Amiodarone

Anti-psychotics

RA EXECUTIVE

Prepared By

50 of 77

Q.A.MANAGER

Approved By



Azithromycin

Disopyramide

Dofetilide

Hydroquinidine

Ibutilide

Quinidine

Sotalol

Tricyclic antidepressants

Vecuronium bromide

Warfarin

Before using Ofloxacin Ornidazole Tablet, inform your doctor about your current list of medications, over the counter products (e.g. vitamins, herbal supplements, etc.), allergies, pre-existing diseases, and current health conditions (e.g. pregnancy, upcoming surgery, etc.). Some health conditions may make you more susceptible to the side-effects of the drug. Take as directed by your doctor or follow the direction printed on the product insert. Dosage is based on your condition. Tell your doctor if your condition persists or worsens. Important counseling points are listed below.

- Avoid consuming milk and dairy products
- Consult the doctor in case of pregnancy or breastfeeding
- Consult your doctor before taking this medicine if having epilepsy and multiple sclerosis
- Do not consume if you have any disease condition like epilepsy or kidney problems
- Do not consume ofloxacin if allergic to it
- Do not drive a vehicle or operate heavy machinery after consuming the medicine
- Do not drive or operate heavy machinery
- Do not take the medicine on empty stomach
- Swallow the tablet whole with water

RA EXECUTIVE

Prepared By

51 of 77

Q.A.MANAGER

Approved By



## 4.5 Interaction with other medicinal products and other forms of interaction

### *Antacids, Sucralfate, Metal Cations*

Co-administered magnesium/aluminum antacids, sucralfate, zinc or iron preparations and didanosine chewable/buffered tablets can reduce absorption of ofloxacin tablets. Therefore, ofloxacin should be taken 2 hours before such preparations.

### *Theophylline, fenbufen or similar non-steroidal anti-inflammatory drugs*

No pharmacokinetic interactions of ofloxacin were found with theophylline in a clinical study. However, a pronounced lowering of the cerebral seizure threshold may occur when quinolones are given concurrently with theophylline, nonsteroidal anti-inflammatory drugs, or other agents, which lower the seizure threshold.

### *Probenecid, cimetidine, furosemide, and methotrexate*

Probenecid decreased the total clearance of ofloxacin by 24%, and increased AUC by 16%. The proposed mechanism is a competition or inhibition for active transport at the renal tubular excretion. Caution should be exercised when ofloxacin is coadministered with drugs that affect the tubular renal secretion such as probenecid, cimetidine, furosemide and methotrexate.

### *Drugs known to prolong QT interval*

Ofloxacin, like other fluoroquinolones, should be used with caution in patients receiving drugs known to prolong the QT interval (e.g. Class IA and III antiarrhythmics, tricyclic antidepressants, macrolides, and antipsychotics)

### *Vitamin K antagonists*

Increased coagulation tests (PT/INR) and/or bleeding, which may be severe, have been reported in patients treated with ofloxacin in combination with a vitamin K antagonist (e.g. warfarin). Coagulation tests should, therefore, be monitored in patients treated with vitamin K antagonists because of a possible increase in the effect of coumarin derivatives.

### *Glibenclamide*

Ofloxacin may cause a slight increase in plasma glibenclamide levels when administered concurrently, it is therefore recommended that patients treated concomitantly with ofloxacin and glibenclamide be monitored particularly closely.

RA EXECUTIVE

Prepared By

52 of 77

Q.A.MANAGER

Approved By



Since hypoglycaemia is then more likely to occur, close monitoring of blood sugar levels is recommended in such cases.

## 4.6 Fertility, Pregnancy and lactation

### *Pregnancy*

Based on a limited amount of human data, the use of fluoroquinolones in the first trimester of pregnancy has not been associated with an increased risk of major malformations or other adverse effects on pregnancy outcome. Animal studies have shown damage to the joint cartilage in immature animals but no teratogenic effects. Therefore ofloxacin must not be used during pregnancy

### *Breast-feeding*

Ofloxacin is excreted into human breast milk in small amounts. Because of the potential for arthropathy and other serious toxicity in the nursing infant, breast-feeding should be discontinued during treatment with ofloxacin (see section 4.3).

## 4.7 Effects on ability to drive and use machine

Since there have been occasional reports of drowsiness/somnolence, impairment of skills, dizziness/vertigo and visual disturbances, which may impair the patient's ability to concentrate and react, and therefore may constitute a risk in situations where these abilities are of special importance (e.g. driving a car or operating machinery), patients should know how they react to ofloxacin before they drive or operate machinery. These effects may be enhanced by alcohol.

## 4.8 Undesirable effects

The information given below is based on data from clinical studies and on extensive post marketing experience.

System organ class	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Very rare (< 1/10,000)	Not known (cannot be estimated from available data)*
Infections and infestations	Fungal infection, Pathogen resistance			
Blood and lymphatic system disorders			Anaemia, Haemolyticanaemia, Leucopenia, Eosinophilia, Thrombocytopenia	Agranulocytosis, Bone marrow failure, Pancytopenia
Immune system disorders		Anaphylactic reaction *, Anaphylactoid	Anaphylactic shock *, Anaphylactoid shock *	

RA EXECUTIVE

Prepared By

53 of 77

Q.A.MANAGER

Approved By





		reaction *, Angioedema *		
Metabolism and Nutrition disorders		Anorexia		Hypoglycaemia in diabetics treated with hypoglycaemic agents (see section 4.4), Hyperglycaemia, Hypoglycaemic coma
Psychiatric disorders	Agitation, Sleep disorder, Insomnia	Psychotic disorder (for e.g. hallucination), Anxiety, Confusional state, Nightmares, Depression		Psychotic disorder and depression with self-endangering behaviour including suicidal ideation or suicide attempt (see Section 4.4), Nervousness
Nervous system disorders	Dizziness, Headache	Somnolence, Paraesthesia, Dysgeusia, Parosmia	Peripheral sensory neuropathy *, Peripheral sensory motor neuropathy *, Convulsion *, Extra-pyramidal symptoms or other disorders of muscular coordination	Tremor, Dyckinesia, Ageusia, Syncope
Eye disorders	Eye irritation	Visual disturbance		Uveitis
Ear and labyrinth disorders	Vertigo		Tinnitus, Hearing loss	Hearing impaired
Cardiac disorders		Tachycardia		Ventricular arrhythmias and torsades de pointes (reported predominantly in patients with risk factors for QT prolongation), ECG QT prolonged (see section 4.4 and 4.9)
Vascular disorders		Hypotension		
Respiratory, thoracic and mediastinal disorders	Cough, Nasopharyngitis	Dyspnoea, Bronchospasm		Allergic pneumonitis, Severe dyspnoea
Gastrointestinal disorders	Abdominal pain, Diarrhoea, Nausea, Vomiting	Enterocolitis, sometimes haemorrhagic	Pseudomembranous colitis *	Dyspepsia, Flatulence, Constipation, Pancreatitis
Hepatobiliary disorders		Hepatic enzymes increased (ALAT, ASAT, LDH, gamma- GT and/or alkaline phosphatase), Blood bilirubin increased	Jaundice cholestatic	Hepatitis, which may be severe * Severe liver injury, including cases with acute liver failure, sometimes fatal, have been reported with ofloxacin, primarily in patients with underlying liver disorders (see section 4.4).
Skin and subcutaneous tissue disorders	Pruritus, Rash	Urticaria, Hot flushes, Hyperhidrosis Pustular rash	Erythema multiforme, Toxic epidermal necrolysis, Photo-sensitivity reaction *,	Stevens-Johnson syndrome, Acute generalised exanthematous pustulosis, Drug rash, Stomatitis

RA EXECUTIVE

Prepared By

54 of 77

Q.A.MANAGER

Approved By



			Drug eruption , Vascular purpura, Vasculitis, which can lead in exceptional cases to skin necrosis	Exfoliative dermatitis
Musculoskeletal and connective tissue disorders		Tendonitis	Arthralgia, Myalgia, Tendon rupture (e.g. Achilles tendon) which may occur within 48 hours of treatment start and may be bilateral	Rhabdomyolysis and/or Myopathy, Muscular weakness, Muscle tear, Muscle rupture, Ligament rupture, Arthritis
Renal and urinary disorders		Serum creatinine increased	Acute renal failure	Acute interstitial nephritis
Congenital, familial and genetic disorders				Attacks of porphyria in patients with porphyria
General disorders and administration site conditions				Asthenia, Pyrexia, Pain (including pain in back, chest and extremities)

\* postmarketing experience

## Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard).

## 4.9 Overdose and treatment

### *Symptoms*

The most important signs to be expected following acute overdose are CNS symptoms such as confusion, dizziness, impairment of consciousness and convulsive seizures increases in QT interval as well as gastrointestinal reactions such as nausea and mucosal erosions.

CNS effects including confusional state, convulsion, hallucination, and tremor have been observed in post marketing experience.

### *Management*

In the case of overdose steps to remove any unabsorbed ofloxacin e.g. gastric lavage, administration of adsorbants and sodium sulphate, if possible during the first 30 minutes, are recommended; antacids are recommended for protection of the gastric mucosa.

RA EXECUTIVE

Prepared By

55 of 77

Q.A.MANAGER

Approved By



In the event of overdose, symptomatic treatment should be implemented. ECG monitoring should be undertaken, because of the possibility of QT interval prolongation. Antacids may be used for protection of gastric mucosa. A fraction of ofloxacin may be removed from the body with haemodialysis. Peritoneal dialysis and CAPD are not effective in removing ofloxacin from the body. No specific antidote exists. Elimination of ofloxacin may be increased by forced diuresis.

## 5- Pharmacological Properties:

### 5.1 Pharmacodynamic Properties

**Pharmacotherapeutic group:** Quinolone Antibacterials, Fluoroquinolones

**ATC code:** J01 MA 01

#### Mechanism of action

Ofloxacin inhibits bacterial DNA replication by inhibiting bacterial topoisomerases, particularly DNA gyrase and topoisomerase IV. It is active after oral administration.

Therapeutic doses of ofloxacin are devoid of pharmacological effects on the voluntary or autonomic nervous system.

The NCCLS MIC breakpoint recommendations are as follows:

$S \leq 2 \text{ mg/l}$  and  $R \geq 1 \text{ mg/l}$

*Haemophilus influenzae* and *Neisseria gonorrhoea* are exceptions with breakpoints at

$S \leq 0.25 \text{ mg/l}$  and  $R \geq 1 \text{ mg/l}$

The BSAC general recommendations are  $S \leq 2 \text{ mg/l}$  and  $R \geq 4 \text{ mg/l}$

According to DIN 58 940, the following limits apply for ofloxacin:

$S \leq 1 \text{ mg/L}$ ,  $I = 2 \text{ mg/L}$ ,  $R \geq 4 \text{ mg/L}$ .

The prevalence of resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. This information gives only an approximate guidance on probabilities whether micro-organisms will be susceptible to ofloxacin or not.

Only those pathogens relevant to the indications are listed.

	<i>European range of acquired bacterial resistance to ofloxacin</i>
<i>Normally susceptible</i>	
<b>Aerobic Gram-positive micro organisms</b>	
<i>S. aureus</i> - methicillin-sensitive	0.3-12.6%
<i>S. pyogenes</i>	2-5%

RA EXECUTIVE

Prepared By

56 of 77

Q.A.MANAGER

Approved By



<b>Aerobic Gram-negative micro organisms</b>	
<i>Acinetobacter spp</i>	0.3-7.3%
<i>Citrobacter spp.</i>	3-15%
<i>Enterobacter spp.</i>	2-13%
<i>E. coli</i>	1-8%
<i>H. influenzae</i>	1%
<i>Klebsiella spp.</i>	1-10%
<i>Moraxella spp.</i>	0-0.2%
<i>Morganellamorganii</i>	0-6.9%
<i>N. gonorrhoeae</i>	25%
<i>Proteus spp.</i>	1-15%
<i>Serratiamarcescens</i>	2-2.4%
<b>Others</b>	
<i>Chlamydia spp</i>	
<i>L. pneumophila</i>	
<b>Intermediately susceptible</b>	
<b>Aerobic Gram-positive micro organisms</b>	
<i>S. pneumoniae</i>	70%
<i>Providentia</i>	17.1%
<b>Aerobic Gram-negative micro organisms</b>	
<i>E. faecalis</i>	50%
<i>P. aeruginosa</i>	20-30%
<i>Serratia spp.</i>	20-40%
<i>Stenotrophomonasmaltophilia</i>	5.1-11%
<b>Others</b>	
<i>Mycoplasma spp.</i>	0-5.3%
<i>Ureaplasma spp.</i>	0-2.1%
<b>Resistant</b>	
<b>Anaerobic bacteria</b>	
<i>S. aureus</i> - methicillin-resistant	69.2-85.7%
<i>T. pallidum</i>	

## Resistance

The main mechanism of bacterial resistance to ofloxacin involves one or more mutations in the target enzymes, which generally confer resistance to other active substances in the class. Efflux pump and impermeability mechanisms of resistance have also been described and may confer variable resistance to active substances in other classes.

RA EXECUTIVE

Prepared By

57 of 77

Q.A.MANAGER

Approved By



## 5.2 Pharmacokinetic Properties

### *Absorption*

The administration of oral doses to fasting volunteers was followed by a rapid and almost complete absorption of ofloxacin. The peak plasma concentration after a single oral dose of 200mg averaged 2.6 µg/ml and was reached within one hour. The plasma elimination half-life was 5.7 to 7.0 hours and was not dose related.

### *Distribution*

The apparent distribution volume was 120 litres. The plasma concentration did not materially rise with repeat doses (accumulation factor for twice daily dosage: 1.5). The plasma protein binding was approx. 25%.

### *Biotransformation*

The biotransformation of ofloxacin was below 5%. The two main metabolites found in the urine were N-desmethyl-ofloxacin and ofloxacin-N-oxide.

### *Elimination*

Excretion is primarily renal. Between 80 and 90% of the dose were recovered from the urine as unchanged substance.

Ofloxacin was present in the bile in glucuronidised form. The pharmacokinetics of ofloxacin after intravenous infusion are very similar to those after oral doses. The plasma half-life is prolonged in persons with renal insufficiency; total and renal clearance decrease in accordance with the creatinine clearance. In renal insufficiency the dose should be reduced.

No clinically relevant interactions were seen with food and no interaction was found between ofloxacin and theophylline.

## 5.3 Preclinical safety Data

Preclinical effects in conventional studies of safety pharmacology, acute toxicity, repeated dose toxicity, reproductive studies were observed only at exposures considered sufficiently in excess of the maximum human exposure indicating little relevance to clinical use. Joint toxicity was observed at exposure in the human therapeutic range in juvenile rats and dogs. Ofloxacin exhibits a neurotoxic potential and causes reversible testicular alterations at high doses.

RA EXECUTIVE

Prepared By

58 of 77

Q.A.MANAGER

Approved By



Mutagenicity studies showed no evidence for mutagenicity of ofloxacin. However, like some other quinolones Ofloxacin is phototoxic in animals at exposure in the human therapeutic range. The phototoxic, photomutagenic and photocarcinogenic potential of ofloxacin is comparable with that of other gyrase inhibitors.

Preclinical data from conventional genotoxicity studies reveal no special hazard to humans, carcinogen potential has not be investigated.

## **6- Pharmaceutical Particulars :**

### **6.1 List of Excipients**

Dibasic Calcium Phosphate  
Maize Starch  
Gelatin  
Purified water  
Cross Carmellose sodium (Vivasole)  
Magnesium Stearate  
Talcum  
Titanium Dioxide  
Tartrazine Yellow Lake  
Sodium Starch Glycolate  
H.P.M.C. E 15  
Lack of Tartrazine  
Sunset Yellow Lake  
Iso Propyl Alcohol  
Methylene Chloride DCM  
PEG 6000

### **6.2 Incompatibilities**

None known

### **6.3 Shelf life**

36 months from the date of manufacture.

### **6.4 Special precautions for storage**

Store in a cool and dry place, protected from light

### **6.5 Nature and contents of container**

10 tablets packed in one blister. Such 1 blister packed in unit printed duplex board carton along with its package insert. Such 10 unit printed carton packed in box and such box add in export worthy shipper.

Note: All pack style may not be marketed.

RA EXECUTIVE

Prepared By

59 of 77

Q.A.MANAGER

Approved By



**7- Marketing Authorization Holder:**

**APHANTEE FORMULATION NIG. LTD.**

Suit FF11, First Floor,  
Pacific Complex No. 9,  
Awka Road, Onitsha,  
Anambra State, Nigeria

**8- Marketing Authorization Number (s) :**

**-Product license / registration Number (s)**

-----

**9- Manufacturer Name:**

**- Name : GLOBELA PHARMA PVT. LTD.**

**- Address :** Plot No. 357, G.I.D.C.,  
Sachin,  
Surat – 394 230,  
Gujarat,  
India.

**- Phone : +91– 261–2398058**

**- Fax : +91– 261–2398058**

**- E-mail : info@globelapharma.com**

**10- Date of first authorization/renewal of the authorization:**

-----

**11- Date of revision of the text:**

-----

**RA EXECUTIVE**

**Prepared By**

**Q.A.MANAGER**

**Approved By**