



UNICURE PHARMACEUTICAL LTD.

**SUMMARY OF PRODUCT CHARACTERISTICS**  
**FOR**  
**FORINT**  
**METFORMIN 500mg TABLETS**

## TABLE OF CONTENT

1. Name of the medicinal product
2. Qualitative and quantitative composition
3. Pharmaceutical form
4. Clinical particulars
  - 4.1 Therapeutic indications
  - 4.2 Posology and method of administration
  - 4.3 Contraindications
  - 4.4 Special warnings and precautions for use
  - 4.5 Interaction with other medicinal products and other forms of interaction
  - 4.6 Fertility, pregnancy and lactation
  - 4.7 Effects on ability to drive and use machines
  - 4.8 Undesirable effects
  - 4.9 Overdose
5. Pharmacological properties
  - 5.1 Pharmacodynamic properties
  - 5.2 Pharmacokinetic properties
  - 5.3 Preclinical safety data
6. Pharmaceutical particulars
  - 6.1 List of excipients
  - 6.2 Incompatibilities
  - 6.3 Shelf life
  - 6.4 Special precautions for storage
  - 6.5 Nature and contents of container

For

(Unicure Pharmaceuticals Limited)

## 6.6 Special precautions for disposal and other handling

---

### 1. Name of the medicinal product

FORINT Metformin 500mg Tablets

---

### 2. Qualitative and quantitative composition

Each film-coated tablet contains metformin hydrochloride 500mg.

For a full list of excipients, see section 6.1.

---

### 3. Pharmaceutical form

White coloured, film-coated, round biconvex tablets with inscription ' ' on one side.

---

### 4. Clinical particulars

#### 4.1 Therapeutic indications

Treatment of type 2 diabetes mellitus, particularly in overweight patients, when dietary management and exercise alone does not result in adequate glycaemic control.

- In adults, FORINT Metformin tablets may be used as monotherapy or in combination with other oral anti-diabetic agents, or with insulin.
- In children from 10 years of age and adolescents, FORINT Metformin tablets may be used as monotherapy or in combination with insulin.

A reduction of diabetic complications has been shown in overweight type 2 diabetic patients treated with metformin as first-line therapy after diet failure (see section 5.1).

#### 4.2 Posology and method of administration

##### Posology

For the different dose regimens FORINT is available as 500 mg, 850 mg or 1000 mg film-coated tablets. In patients receiving a high FORINT metformin hydrochloride dose (2 to 3 grams per day), it is possible to replace two 500 mg film-coated tablets with one 1000 mg film-coated tablet.

Adults with normal renal function ( $GFR \geq 90$  mL/min)

Monotherapy and combination with other oral antidiabetic agents

- The usual starting dose is 500 mg or 850 mg FORINT metformin hydrochloride 2 or 3 times daily given during or after meals. After 10 to 15 days the dose should be adjusted on the basis of blood glucose measurements. A slow increase of dose may improve gastrointestinal tolerability.

The maximum recommended dose of FORINT metformin hydrochloride is 3 g daily, taken as 3 divided doses.

- If transfer from another oral antidiabetic agent is intended, discontinue the other agent and initiate metformin at the dose indicated above.

Combination with insulin

FORINT Metformin and insulin may be used in combination therapy to achieve better blood glucose control. Metformin hydrochloride is given at the usual starting dose of 500 mg or 850 mg 2 or 3 times daily, while insulin dosage is adjusted on the basis of blood glucose measurements.

Due to potential for decreased renal function in elderly subjects, the metformin dosage should be adjusted based on renal function. Regular assessment of renal function is necessary (see section 4.4)

#### Renal impairment

A GFR should be assessed before initiation of treatment with FORINT metformin containing products and at least annually thereafter. In patients at an increased risk of further progression of renal impairment and in the elderly, renal function should be assessed more frequently, e.g. every 3-6 months.

GFR (mL/min)	Total maximum daily dose (to be divided into 2-3 daily doses)	Additional considerations
60-89	3000 mg	Dose reduction may be considered in relation to declining renal function
45-59	2000 mg	Factors that may increase the risk of lactic acidosis (see section 4.4) should be reviewed before considering initiation of metformin.
30-44	1000 mg	The starting dose is at most half of the maximum dose.
<30	-	Metformin is contraindicated

#### Paediatric population

#### Monotherapy and combination with insulin

- FORINT Metformin tablets can be used in children from 10 years of age and adolescents.
- The usual starting dose is one tablet of 500 mg or 850 mg metformin hydrochloride once daily, given during or after meals.

After 10 to 15 days the dose should be adjusted on the basis of blood glucose measurements. A slow increase of dose may improve gastrointestinal tolerability. The maximum recommended dose of metformin hydrochloride is 2 g daily, taken as 2 or 3 divided doses.

#### **Pre-diabetic state: impaired fasting glycaemia (IFG) and/or impaired glucose tolerance (IGT)**

The usual starting dose is 500 mg (1 film-coated tablet Formetic 500 mg or ½ film-coated tablet Formetic 1000 mg) daily. Dose should be adjusted according to the clinical effect up to 1700 mg daily, taken as divided doses.

#### **Polycystic Ovary Syndrome (PCOS)**

The usual dose is 500 mg (1 film-coated tablet FORINT 500 mg or ½ film-coated tablet Formetic 1000 mg) three times a day.

#### **4.3 Contraindications**

- ✓ Hypersensitivity to FORINT metformin or to any of the Excipients listed in section 6.1.
- ✓ Any type of acute metabolic acidosis (such as lactic acidosis, diabetic ketoacidosis)
- ✓ Diabetic pre-coma
- ✓ Severe renal failure (GFR <30 mL/min)
- ✓ Acute conditions with the potential to alter renal function such as: dehydration, severe infection, shock

For

(Unicure Pharmaceuticals Limited)

- ✓ Disease which may cause tissue hypoxia (especially acute disease, or worsening of chronic disease) such as: decompensated heart failure, respiratory failure, recent myocardial infarction, shock
- ✓ Hepatic insufficiency, acute alcohol intoxication, alcoholism

#### 4.4 Special warnings and precautions for use

##### **Lactic acidosis:**

Lactic acidosis, a very rare but serious metabolic complication, most often occurs at acute worsening of renal function or cardiorespiratory illness or sepsis. FORINT Metformin accumulation occurs at acute worsening of renal function and increases the risk of lactic acidosis.

In case of dehydration (severe diarrhoea or vomiting, fever or reduced fluid intake), FORINT metformin should be temporarily discontinued and contact with a health care professional is recommended.

Medicinal products that can acutely impair renal function (such as antihypertensives, diuretics and NSAIDs) should be initiated with caution in metformin-treated patients. Other risk factors for lactic acidosis are inadequately controlled diabetes, ketosis, prolonged fasting, excessive alcohol intake, hepatic insufficiency and any condition associated with hypoxia, as well as concomitant use of medicinal products that may cause lactic acidosis (see section 4.3 and 4.5)

Patients and/or care-givers should be informed of the risk of lactic acidosis.

Lactic acidosis is characterised by acidotic dyspnoea, abdominal pain, muscle cramps, asthenia and hypothermia followed by coma. In case of suspected symptoms, the patient should stop taking metformin and seek immediate medical attention. Diagnostic laboratory findings are decreased blood pH (<7.35), increased plasma lactate levels (>5 mmol/L) and an increased anion gap and lactate/pyruvate ratio.

##### Renal Function

GFR should be assessed before treatment initiation and regularly thereafter, see section 4.2. FORINT Metformin is contraindicated in patients with GFR <30 mL/min and should be temporarily discontinued in the presence of conditions that alter renal function, see section 4.3.

##### Cardiac function

Patients with heart failure are more at risk of hypoxia and renal insufficiency. In patients with stable chronic heart failure, metformin may be used with a regular monitoring of cardiac and renal function.

For patients with acute and unstable heart failure, metformin is contraindicated (see section 4.3).

##### Administration of iodinated contrast agent

Intravascular administration of iodinated contrast agents may lead to contrast induced nephropathy resulting in FORINT metformin accumulation and an increased risk of lactic acidosis. FORINT Metformin should be discontinued prior to, or at the time of the imaging procedure and not restarted until at least 48 hours after, provided that renal function has been re-evaluated and found to be stable (see sections 4.2 and 4.5)

##### Surgery

FORINT Metformin Hydrochloride must be discontinued at the time of surgery under general, spinal or epidural anaesthesia. Therapy may be restarted no earlier than 48 hours following surgery or resumption of oral nutrition and provided that renal function has been re-evaluated and found to be stable.

### Children and Adolescents

The diagnosis of type 2 diabetes mellitus should be confirmed before treatment with metformin is initiated.

No effect of metformin on growth and puberty has been detected during controlled clinical studies of one-year duration but no long-term on these specific points are available.

Therefore, a careful follow-up of the effect of metformin on these parameters in metformin-treated children, especially prepubescent children, is recommended

### **Children aged between 10 and 12 years**

Only 15 subjects aged between 10 and 12 years were included in the controlled clinical studies conducted in children and adolescents. Although efficacy and safety of FORINT metformin in these children did not differ from efficacy and safety in older children and adolescents, particular caution is recommended when prescribing to children aged between 10 and 12 years.

### **Other precautions:**

- All patients should continue their diet with a regular distribution of carbohydrate intake during the day. Overweight patients should continue their energy-restricted diet.
- The usual laboratory tests for diabetes monitoring should be performed regularly.
- FORINT Metformin alone does not cause hypoglycaemia, but caution is advised when it is used in combination with insulin or other oral antidiabetics (e.g. sulfonylureas or meglitinides).

### **Metformin tablets contains Sodium:**

FORINT Metformin tablets contains less than 1 mmol (23 mg) of sodium per tablet, that is to say it is essentially 'sodium-free'.

## **4.5 Interaction with other medicinal products and other forms of interaction**

### Concomitant use not recommended

#### *Alcohol*

Alcohol intoxication is associated with an increased risk of lactic acidosis, particularly in case of fasting, malnutrition or hepatic impairment.

#### *Iodinated contrast agents*

FORINT Metformin must be discontinued prior to or at the time of the imaging procedure and not restarted until at least 48 hours, provided that renal function has been re-evaluated and found to be stable, see sections 4.2 and 4.4.

### Combinations requiring precautions for use

Some medicinal products can adversely affect renal function which may increase the risk of lactic acidosis, e.g. NSAIDs, including selective cyclooxygenase (COX) II inhibitors, ACE inhibitors, angiotensin II receptor antagonists and diuretics, especially loop diuretics. When starting or using such products in combination with metformin, close monitoring of renal function is necessary.

*Medicinal products with intrinsic hyperglycaemic activity (e.g. glucocorticoids (systemic and local routes) and sympathomimetics).*

More frequent blood glucose monitoring may be required, especially at the beginning of the treatment. If necessary, adjust the metformin dosage of the antidiabetic drug during therapy with the respective medicinal product and upon its discontinuation.

#### *Organic cation transporters (OCT)*

Metformin is a substrate of both transporters OCT1 and OCT2.

Co-administration of metformin with

- Inhibitors of OCT1 (such as verapamil) may reduce efficacy of metformin.
- Inducers of OCT1 (such as rifampicin) may increase gastrointestinal absorption and efficacy of metformin.
- Inhibitors of OCT2 (such as cimetidine, dolutegravir, ranolazine, trimethoprim, vandetanib, isavuconazole) may decrease the renal elimination of metformin and thus lead to an increase in metformin plasma concentration.
- Inhibitors of both OCT1 and OCT2 (such as crizotinib, olaparib) may alter efficacy and renal elimination of metformin.

Caution is therefore advised, especially in patients with renal impairment, when these drugs are co-administered with metformin, as metformin plasma concentration may increase. If needed, dose adjustment of FORINT metformin may be considered as OCT inhibitors/inducers may alter the efficacy of metformin.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

Uncontrolled diabetes during pregnancy (gestational or permanent) is associated with increased risk of congenital abnormalities and perinatal mortality.

A limited amount of data from the use of FORINT metformin in pregnant women does not indicate an increased risk of congenital abnormalities. Animal studies do not indicate harmful effects with respect to pregnancy, embryonic or foetal development, parturition or postnatal development (see section 5.3).

When the patient plans to become pregnant and during pregnancy, it is recommended that diabetes is not treated with FORINT metformin but insulin be used to maintain blood glucose levels as close to normal as possible, to reduce the risk of foetal malformations of the foetus.

### Breast-feeding

FORINT Metformin is excreted into human breast milk. No adverse effects were observed in breastfed newborns/infants. However, as only limited data are available, breast-feeding is not recommended during FORINT metformin treatment. A decision on whether to discontinue breast-feeding should be made, taking into account the benefit of breast-feeding and the potential risk to adverse effects on the child.

### Fertility

Fertility of male or female rats was unaffected by FORINT metformin when administered at doses as high as 600 mg/kg/day, which is approximately three times the maximum recommended human daily dose based on body surface area comparisons.

#### 4.7 Effects on ability to drive and use machines

FORINT Metformin monotherapy does not cause hypoglycaemia and therefore has no effect on the ability to drive or to use machines.

However, patients should be alerted to the risk of hypoglycaemia when metformin is used in combination with other antidiabetic agents (e.g. sulfonylureas, insulin, or meglitinides).

#### 4.8 Undesirable effects

During treatment initiation, the most common adverse reactions are nausea, vomiting, diarrhoea, abdominal pain and loss of appetite which resolve spontaneously in most cases. To prevent them, it is recommended to take metformin in 2 or 3 daily doses and to increase slowly the doses.

The following adverse reactions may occur under treatment with FORINT metformin. Frequencies are defined as follows: very common:  $\geq 1/10$ ; common  $> 1/100$ ,  $< 1/10$ ; uncommon  $> 1/1,000$ ,  $< 1/100$ ; rare  $> 1/10,000$ ,  $< 1/1,000$ ; very rare  $< 1/10,000$ , not known (cannot be estimated from the available data).

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

##### Metabolism and nutrition disorders:

###### *Very rare*

- Lactic acidosis (see section 4.4).
- Decrease of vitamin B12 absorption with decrease of serum levels during long term use of metformin hydrochloride. Consideration of such aetiology is recommended if a patient present with megaloblastic anaemia.

##### Nervous system disorders:

###### *Common*

- Taste disturbance

##### Gastrointestinal disorders:

###### *Very common*

- Gastrointestinal disorders such as nausea, vomiting, diarrhoea, abdominal pain and loss of appetite. These undesirable effects occur most frequently during initiation of therapy and resolve spontaneously in most cases. To prevent them, it is recommended that Metformin Hydrochloride be taken in 2 or 3 daily doses during or after meals. A slow increase of the dose may also improve gastrointestinal tolerability.

##### Hepatobiliary disorders:

###### *Very rare*

- Isolated reports of liver function tests abnormalities or hepatitis resolving upon metformin hydrochloride discontinuation.

##### Skin and subcutaneous tissue disorders:

###### *Very rare*

- Skin reactions such as erythema, pruritus, urticaria

##### Paediatric population



In published and post marketing data and in controlled clinical studies in a limited paediatric population aged 10-16 years treated during 1 year, adverse event reporting was similar in nature and severity to that reported in adults.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal products is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

### **4.9 Overdose**

Hypoglycaemia has not been seen with FORINT metformin hydrochloride doses of up to 85 g, although lactic acidosis has occurred in such circumstances. High overdose of metformin or concomitant risks may lead to lactic acidosis. Lactic acidosis is a medical emergency and must be treated in hospital. The most effective method to remove lactate and FORINT metformin is haemodialysis.

---

## **5. Pharmacological properties**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic Group: Blood glucose lowering drugs. Biguanide oral hypoglycaemic agents-ATC code: A10BA02

#### Mechanism of action

FORINT Metformin is a biguanide with antihyperglycaemic effects, lowering both basal and post-prandial plasma glucose. It does not stimulate insulin secretion and therefore does not produce hypoglycaemia.

FORINT Metformin may act via 3 mechanisms:

1. Reduction of hepatic glucose production by inhibiting gluconeogenesis and glycogenolysis.
2. In muscle, by increasing insulin sensitivity, improving peripheral glucose uptake and utilisation.
3. And delay of intestinal glucose absorption.

FORINT Metformin stimulates intracellular glycogen synthesis by acting on glycogen synthase.

FORINT Metformin increases the transport capacity of all types of membrane glucose transporters (GLUTs) known to date.

#### Pharmacodynamic effects

In clinical studies, use of metformin was associated with either a stable body weight or modest weight loss. In humans, independently of its action on glycaemia, metformin has favourable effects on lipid metabolism. This has been shown at therapeutic doses in controlled, medium-term or long-term clinical studies: Metformin reduces total cholesterol, LDL, cholesterol and triglyceride levels.

Clinical Efficacy:

The prospective randomised study (UKPDS) has established the long-term benefit of intensive blood glucose control in adult patients with type 2 diabetes.

Analysis of the results for overweight patients treated with metformin after failure of diet alone showed:

- A significant reduction of the absolute risk of any diabetes-related complication in the metformin group (29.8 events/1000 patients-years) versus diet alone (43.3 events/1000 patient-years),  $p=0.0023$ , and versus the combined sulphonylurea and insulin monotherapy groups (40.1 events/1000 patients-years),  $p=0.0034$ .
- A significant reduction of the absolute risk of the diabetes-related mortality: Metformin 7.5 events/1000 patient-years, diet alone 12.7 events/1000 patient-year,  $p=0.017$
- A significant reduction of the absolute risk of overall mortality: metformin hydrochloride 13.5 events/1000 patient-years versus diet alone 20.6 events/1000 patient-years ( $p=0.011$ ), and versus the combined sulphonylurea and insulin monotherapy groups 18.9 events/1000 patient-years ( $p=0.021$ )
- A significant reduction in the absolute risk of myocardial infarction: metformin 11 events/1000 patients-years, diet alone 18 events/1000 patient-years ( $p=0.01$ )

Benefit regarding clinical outcome has not been shown for metformin used as second-line therapy, in combination with a sulphonylurea.

In type I diabetes, the combination of metformin and insulin has been used in selected patients, but the clinical benefit of this combination has not been formally established.

#### Paediatric population

Controlled clinical studies in a limited paediatric population aged 10-16 years treated during 1 year demonstrated a similar response in glycaemic control to that seen in adults.

## **5.2 Pharmacokinetic properties**

### Absorption

After an oral dose of metformin hydrochloride tablet, maximum plasma concentration ( $C_{max}$ ) is reached in 2.5 hours ( $t_{max}$ ). Absolute bioavailability of a 500mg or 850mg metformin hydrochloride tablet is approximately 50-60% in healthy subjects. After an oral dose, the non-absorbed fraction recovered in faeces was 20-30%.

After oral administration, metformin absorption is saturable and incomplete. It is assumed that the pharmacokinetics of metformin absorption is non-linear. At the recommended metformin doses and the dosing schedules, steady state plasma concentrations are reached within 24 to 48 hours and are generally less than  $1\mu\text{g/ml}$ . In controlled clinical trials, maximum metformin plasma levels ( $C_{max}$ ) did not exceed  $5\mu\text{g/ml}$ , even at maximum doses.

Food decreases the extent and slightly delays the absorption of metformin.

Following oral administration of a 850 mg, a 40% lower plasma peak concentration a 25% decrease in AUC (area under the curve) and a 35 minute prolongation of the time to peak plasma concentration were observed. The clinical relevance of these findings is unknown.

### Distribution

Plasma protein binding is negligible. Metformin partitions into erythrocytes. The blood peak is lower than the plasma peak and appears approximately the same time. The red blood cells most likely represent a secondary compartment of distribution. The mean volume of distribution ( $V_d$ ) ranged between 63-276 L.

### Metabolism

FORINT Metformin is excreted unchanged in the urine. No metabolites have been identified in humans.

### Elimination

Renal clearance of metformin is >400ml/min, indicating that metformin is eliminated by glomerular filtration and tubular secretion. Following an oral dose, the apparent terminal elimination half-life is approximately 6.5 hours.

When renal function is impaired, renal clearance is decreased in proportion to that of creatinine and thus the elimination half-life is prolonged, leading to increased levels of metformin in plasma.

### Characteristics in specific groups of patients

#### Renal impairment

The available data in subjects with moderate renal insufficiency are scarce and no reliable estimation of the systemic exposure to metformin in this subgroup as compared to subjects with normal renal function could be made. Therefore, the dose adaptation should be made upon clinical efficacy/tolerability considerations (see section 4.2).

#### Paediatric population

Single dose study: After single doses of metformin hydrochloride 500 mg, paediatric patients have shown similar pharmacokinetic profile to that observed in healthy adults.

Multiple dose study: Data are restricted to one study. After repeated doses of 500 mg twice daily for 7 days in paediatric patients the peak plasma concentration (C<sub>max</sub>) and systemic exposure (AUC<sub>0-t</sub>) were reduced by approximately 33% and 40%, respectively compared to diabetic adults who received repeated doses of 500 mg twice daily for 14 days. As the dose is individually titrated based on glycaemic control, this is of limited clinical relevance.

### **5.3 Preclinical safety data**

Preclinical data reveal no special hazard for humans based on conventional studies on safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and reproductive toxicity.

---

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Starch

Calcium sulfate

7.5% Hypromellose in 50% Ethanol

15% starch syrup

talcum powder

Sodium Carboxymethyl Starch

## **6.2 Incompatibilities**

Not applicable

## **6.3 Shelf life**

36 Months

## **6.4 Special precautions for storage**

Do not store above 25°C. Store in the original package. Keep the containers tightly closed.

## **6.5 Nature and contents of container**

Blister: Transparent PVC /PVdC/Aluminium foil. Packs of 6 x 14 Tablets

## **6.6 Special precautions for disposal and other handling**

Not applicable

---

## **7. Marketing authorisation holder**

Unicure Pharmaceutical Ltd

Ikofa Village, Lagos - Benin Expressway, Ikofa

Ijebu- Ode

Email: [unicurepharms@163.com](mailto:unicurepharms@163.com)

---

## **8. Marketing authorisation number(s)**

---

## **9. Date of first authorisation/renewal of the authorisation**

---

## **10. Date of revision of the text**