



National Agency for Food & Drug Administration & Control (NAFDAC)

Registration & Regulatory Affairs (R & R) Directorate

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

1. Name of the medicinal product

Dr. Botsford Ciprofloxacin 500 mg film-coated tablets

2. Qualitative and quantitative composition

Each film-coated tablet contains 500 mg ciprofloxacin (as hydrochloride).

For the full list of excipients, see section 6.1.

3. Pharmaceutical form

Film-coated tablet.

White to off white, capsule shaped, film coated tablets, with a score line on one side and debossed with 'EGMP' on the other side. The tablet can be divided into equal doses.

4. Clinical particulars

4.1 Therapeutic indications

Because of the risk of prolonged, disabling and potentially irreversible serious adverse drug reactions (see section 4.4 and section 4.8) this product must only be prescribed when other antibiotics that are commonly recommended for the infection are inappropriate. This applies to all indications listed below. Situations where other antibiotics are considered to be inappropriate are where:

- there is resistance to other first-line antibiotics recommended for the infection;
- other first-line antibiotics are contraindicated in an individual patient;
- other first-line antibiotics have caused side effects requiring treatment to be stopped;
- treatment with other first-line antibiotics has failed.

Ciprofloxacin film-coated tablets are indicated for the treatment of the following infections (see sections 4.4 and 5.1). Special attention should be paid to available information on resistance to ciprofloxacin before commencing therapy.

Adults

- Lower respiratory tract infections due to Gram-negative bacteria
 - pneumonia
 - broncho-pulmonary infections in cystic fibrosis or in bronchiectasis
- Chronic suppurative otitis media
- Acute exacerbation of chronic sinusitis especially if these are caused by Gram-negative bacteria
- Acute pyelonephritis
- Complicated urinary tract infections
- Bacterial prostatitis
- Genital tract infections
 - gonococcal urethritis and cervicitis due to susceptible *Neisseria gonorrhoeae*
 - epididymo-orchitis including cases due to susceptible *Neisseria gonorrhoeae*
 - pelvic inflammatory disease including cases due to susceptible *Neisseria gonorrhoeae*
- Intra-abdominal infections
- Infections of the skin and soft tissue caused by Gram-negative bacteria
- Infections of the bones and joints
- Inhalation anthrax (post-exposure prophylaxis and curative treatment)

Ciprofloxacin may be used in the management of neutropenic patients with fever that is suspected to be due to a bacterial infection.

In **exacerbations of chronic obstructive pulmonary disease** Ciprofloxacin should be used only when it is considered inappropriate to use other antibacterial agents that are commonly recommended for the treatment of these infections.

In **uncomplicated acute cystitis** Ciprofloxacin should be used only when it is considered inappropriate to use other antibacterial agents that are commonly recommended for the treatment of these infections.

Children and adolescents

- Broncho-pulmonary infections due to *Pseudomonas aeruginosa* in patients with cystic fibrosis
- Complicated urinary tract infections and acute pyelonephritis
- Inhalation anthrax (post-exposure prophylaxis and curative treatment)

Ciprofloxacin may also be used to treat severe infections in children and adolescents when this is considered to be necessary.

Treatment should be initiated only by physicians who are experienced in the treatment of cystic fibrosis and/or severe infections in children and adolescents (see sections 4.4 and 5.1).

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

The dosage is determined by the indication, the severity and the site of the infection, the susceptibility to ciprofloxacin of the causative organism(s), the renal function of the patient and, in children and adolescents the body weight.

The duration of treatment depends on the severity of the illness and on the clinical and bacteriological course.

Treatment of infections due to certain bacteria (e.g. *Pseudomonas aeruginosa*, *Acinetobacter* or *Staphylococci*) may require higher ciprofloxacin doses and co-administration with other appropriate antibacterial agents.

Treatment of some infections (e.g. pelvic inflammatory disease, intra-abdominal infections, infections in neutropenic patients and infections of bones and joints) may require co-administration with other appropriate antibacterial agents depending on the pathogens involved.

Elderly patients

Elderly patients should receive a dose selected according to the severity of the infection and the patient's creatinine clearance.

Patients with renal and hepatic impairment

Recommended starting and maintenance doses for patients with impaired renal function:

Creatinine [mL/min/1.73 m²]	Clearance Serum [mol/L]	Creatinine [μ Oral Dose [mg]
> 60	< 124	See Usual Dosage.
30-60	124 to 168	250-500 mg every 12 h
<30	> 169	250-500 mg every 24 h
Patients on haemodialysis	> 169	250-500 mg every 24 h (after dialysis)
Patients on peritoneal dialysis	> 169	250-500 mg every 24 h

In patients with impaired liver function no dose adjustment is required.

Dosing in children with impaired renal and/or hepatic function has not been studied.

Method of administration

Tablets are to be swallowed unchewed with fluid. They can be taken independent of mealtimes. If taken on an empty stomach, the active substance is absorbed more rapidly. Ciprofloxacin tablets should not be taken with dairy products (e.g. milk, yoghurt) or mineral-fortified fruit-juice (e.g. calcium-fortified orange juice) (see section 4.5).

In severe cases or if the patient is unable to take tablets (e.g. patients on enteral nutrition), it is recommended to commence therapy with intravenous ciprofloxacin until a switch to oral administration is possible.

4.3 Contraindications

- Hypersensitivity to the active substance, to other quinolones or to any of the excipients listed in section 6.1.
- Concomitant administration of ciprofloxacin and tizanidine (see section 4.5).

4.4 Special warnings and precautions for use

The use of ciprofloxacin should be avoided in patients who have experienced serious adverse reactions in the past when using quinolone or fluoroquinolone containing products (see section 4.8). Treatment of these patients with ciprofloxacin should only be initiated in the absence of alternative treatment options and after careful benefit/risk assessment (see also section 4.3).

Prolonged, disabling and potentially irreversible serious adverse drug reactions

Cases of prolonged (continuing for months or years), disabling and potentially irreversible serious adverse drug reactions affecting different, sometimes multiple, body systems (including musculoskeletal, nervous, psychiatric and senses) have been reported in patients receiving quinolones and fluoroquinolones irrespective of their age and pre-existing risk factors.

There are no pharmacological treatments established to be effective treatments of the symptoms of long lasting or disabling side effects associated with fluoroquinolones. Ciprofloxacin should be discontinued immediately at the first signs or symptoms of any serious adverse reaction and patients should be advised to contact their prescriber for advice, so that symptoms can be appropriately investigated and to avoid further exposure which could potentially worsen adverse reactions.

Severe infections and mixed infections with Gram-positive and anaerobic pathogens

Ciprofloxacin monotherapy is not suited for treatment of severe infections and infections that might be due to Gram-positive or anaerobic pathogens. In such infections ciprofloxacin must be co-administered with other appropriate antibacterial agents.

Streptococcal Infections (including *Streptococcus pneumoniae*)

Ciprofloxacin is not recommended for the treatment of streptococcal infections due to inadequate efficacy.

Genital tract infections

Gonococcal urethritis, cervicitis, epididymo-orchitis and pelvic inflammatory diseases may be caused by fluoroquinolone-resistant *Neisseria gonorrhoeae* isolates.

Therefore, ciprofloxacin should be administered for the treatment of gonococcal urethritis or cervicitis only if ciprofloxacin-resistant *Neisseria gonorrhoeae* can be excluded.

For epididymo-orchitis and pelvic inflammatory diseases, empirical ciprofloxacin should only be considered in combination with another appropriate antibacterial agent (e.g. a cephalosporin) unless ciprofloxacin resistant *Neisseria gonorrhoeae* can be excluded. If clinical improvement is not achieved after 3 days of treatment, the therapy should be reconsidered.

Urinary tract infections

Resistance to fluoroquinolones of *Escherichia coli* – the most common pathogen involved in urinary tract infections – varies across the European Union. Prescribers are advised to take into account the local prevalence of resistance in *Escherichia coli* to quinolones.

The single dose of ciprofloxacin that may be used in uncomplicated cystitis in premenopausal women is expected to be associated with lower efficacy than the longer treatment duration. This is all the more to be taken into account as regards the increasing resistance level of *Escherichia coli* to quinolones.

Intra-abdominal infections

There are limited data on the efficacy of ciprofloxacin in the treatment of post-surgical intra-abdominal infections.

Travellers' diarrhoea

The choice of ciprofloxacin should take into account information on resistance to ciprofloxacin in relevant pathogens in the countries visited.

Infections of the bones and joints

Ciprofloxacin should be used in combination with other antimicrobial agents depending on the results of the microbiological documentation.

Inhalational anthrax

Use in humans is based on *in-vitro* susceptibility data and on animal experimental data together with limited human data. Treating physicians should refer to national and/or international consensus documents regarding the treatment of anthrax.

Paediatric population

The use of ciprofloxacin in children and adolescents should follow available official guidance. Ciprofloxacin treatment should be initiated only by physicians who are experienced in the treatment of cystic fibrosis and/or severe infections in children and adolescents.

Ciprofloxacin has been shown to cause arthropathy in weight-bearing joints of immature animals. Safety data from a randomised double-blind study on ciprofloxacin use in children (ciprofloxacin: n=335, mean age = 6.3 years; comparators: n=349, mean age = 6.2 years; age range = 1 to 17 years) revealed an incidence of suspected drug-related arthropathy (discerned from joint-related clinical signs and symptoms) by Day +42 of 7.2% and 4.6%. Respectively, an incidence of drug-related arthropathy by 1-year follow-up was 9.0% and 5.7%. The increase of suspected drug-related arthropathy cases over time was not statistically significant between groups. Treatment should be initiated only after a careful benefit/risk evaluation, due to possible adverse events related to joints and/or surrounding tissue (see section 4.8)..

Broncho-pulmonary infections in cystic fibrosis

Clinical trials have included children and adolescents aged 5-17 years. More limited experience is available in treating children between 1 and 5 years of age.

Complicated urinary tract infections and pyelonephritis

Ciprofloxacin treatment of urinary tract infections should be considered when other treatments cannot be used, and should be based on the results of the microbiological documentation.

Clinical trials have included children and adolescents aged 1-17 years.

Other specific severe infections

Other severe infections in accordance with official guidance, or after careful benefit-risk evaluation when other treatments cannot be used, or after failure to conventional therapy and when the microbiological documentation can justify a ciprofloxacin use.

The use of ciprofloxacin for specific severe infections other than those mentioned above has not been evaluated in clinical trials and the clinical experience is limited. Consequently, caution is advised when treating patients with these infections.

Hypersensitivity

Hypersensitivity and allergic reactions, including anaphylaxis and anaphylactoid reactions, may occur following a single dose (see section 4.8) and may be life-threatening. If such reaction occurs, ciprofloxacin should be discontinued and an adequate medical treatment is required.

Musculoskeletal System

Ciprofloxacin should generally not be used in patients with a history of tendon disease/disorder related to quinolone treatment. Nevertheless, in very rare instances, after microbiological documentation of the causative organism and evaluation of the risk/benefit balance, ciprofloxacin may be prescribed to these patients for the treatment of certain severe infections, particularly in the event of failure of the standard therapy or bacterial resistance, where the microbiological data may justify the use of ciprofloxacin.

Tendinitis and tendon rupture

Tendinitis and tendon rupture (especially but not limited to Achilles tendon), sometimes bilateral, may occur as early as within 48 hours of starting treatment with quinolones and fluoroquinolones and have been reported to occur even up to several months after discontinuation of treatment. The risk of tendinitis and tendon rupture is increased in older patients, patients with renal impairment, patients with solid organ transplants, and those treated concurrently with corticosteroids. Therefore, concomitant use of corticosteroids should be avoided.

At the first sign of tendinitis (e.g. painful swelling, inflammation) the treatment with ciprofloxacin should be discontinued and alternative treatment should be considered. The affected limb(s) should be appropriately treated (e.g. immobilisation). Corticosteroids should not be used if signs of tendinopathy occur.

Ciprofloxacin should be used with caution in patients with myasthenia gravis, because symptoms can be exacerbated (see section 4.8).

Vision disorders

If vision becomes impaired or any effects on the eyes are experienced, an eye specialist should be consulted immediately.

Photosensitivity

Ciprofloxacin has been shown to cause photosensitivity reactions. Patients taking ciprofloxacin should be advised to avoid direct exposure to either extensive sunlight or UV irradiation during treatment (see section 4.8).

Central Nervous System

Ciprofloxacin like other quinolones are known to trigger seizures or lower the seizure threshold. Cases of status epilepticus have been reported. Ciprofloxacin should be used with caution in patients with CNS disorders which may be predisposed to seizure. If seizures occur ciprofloxacin should be discontinued (see section 4.8). Psychiatric reactions may occur even after the first administration of ciprofloxacin. In rare cases, depression or psychosis can progress to suicidal ideations/thoughts culminating in attempted suicide or completed suicide. In the occurrence of such cases, ciprofloxacin should be discontinued.

Peripheral neuropathy

Cases of sensory or sensorimotor polyneuropathy resulting in paraesthesia, hypaesthesia, dysesthesia, or weakness have been reported in patients receiving quinolones and fluoroquinolones. Patients under treatment with ciprofloxacin should be advised to inform their doctor prior to continuing treatment if symptoms of neuropathy such as pain, burning, tingling, numbness, or weakness develop in order to prevent the development of potentially irreversible condition (see section 4.8).

Cardiac disorders

Caution should be taken when using fluoroquinolones, including ciprofloxacin, in patients with known risk factors for prolongation of the QT interval such as, for example:

- congenital long QT syndrome
- concomitant use of drugs that are known to prolong the QT interval (e.g. Class IA and III antiarrhythmics, tricyclic antidepressants, macrolides, antipsychotics)

- uncorrected electrolyte imbalance (e.g. hypokalaemia, hypomagnesaemia)
- cardiac disease (e.g. heart failure, myocardial infarction, bradycardia)

Elderly patients and women may be more sensitive to QTc-prolonging medications. Therefore, caution should be taken when using fluoroquinolones, including ciprofloxacin, in these populations. (See section 4.2 Elderly patients, section 4.5, section 4.8, section 4.9).

Dysglycaemia

As with all quinolones, disturbances in blood glucose, including both hypoglycaemia and hyperglycaemia have been reported, usually in diabetic patients receiving concomitant treatment with an oral hypoglycaemic agent (e.g., glibenclamide) or with insulin. Cases of hypoglycaemic coma have been reported. In diabetic patients, careful monitoring of blood glucose is recommended (see section 4.8).

Gastrointestinal System

The occurrence of severe and persistent diarrhoea during or after treatment (including several weeks after treatment) may indicate an antibiotic-associated colitis (life-threatening with possible fatal outcome), requiring immediate treatment. In such cases, ciprofloxacin should immediately be discontinued, and an appropriate therapy initiated. Anti-peristaltic drugs are contraindicated in this situation.

Renal and urinary system

Crystalluria related to the use of ciprofloxacin has been reported (see section 4.8). Patients receiving ciprofloxacin should be well hydrated and excessive alkalinity of the urine should be avoided.

Impaired renal function

Since ciprofloxacin is largely excreted unchanged via renal pathway dose adjustment is needed in patients with impaired renal function as described in section 4.2 to avoid an increase in adverse drug reactions due to accumulation of ciprofloxacin.

- for heart valve regurgitation/incompetence (e.g. infective endocarditis).

The risk of aortic aneurysm and dissection, and their rupture may also be increased in patients treated concurrently with systemic corticosteroids.

In case of sudden abdominal, chest or back pain, patients should be advised to immediately consult a physician in an emergency department.

Patients should be advised to seek immediate medical attention in case of acute dyspnoea, new onset of heart palpitations, or development of oedema of the abdomen or lower extremities.

Sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Effects of other products on ciprofloxacin:

Drugs known to prolong QT interval

Ciprofloxacin, like other fluoroquinolones, should be used with caution in patients receiving drugs known to prolong QT interval (e.g. Class IA and III antiarrhythmics, tricyclic antidepressants, macrolides, antipsychotics) (see section 4.4).

Chelation Complex Formation

The simultaneous administration of ciprofloxacin (oral) and multivalent cation-containing drugs and mineral supplements (e.g. calcium, magnesium, aluminium, iron), polymeric phosphate binders (e.g. sevelamer or lanthanum carbonate), sucralfate or antacids, and highly buffered drugs (e.g. didanosine tablets) containing magnesium, aluminium, or calcium reduces the absorption of ciprofloxacin. Consequently, ciprofloxacin should be administered either 1-2 hours before or at least 4 hours after these preparations. The restriction does not apply to antacids belonging to the class of H₂ receptor blockers.

Food and Dairy Products

Dietary calcium as part of a meal does not significantly affect absorption. However, the concurrent administration of dairy products or mineral-fortified drinks alone (e.g. milk, yoghurt, calcium-fortified orange juice) with ciprofloxacin should be avoided because absorption of ciprofloxacin may be reduced.

Probenecid

Probenecid interferes with renal secretion of ciprofloxacin. Co-administration of probenecid and ciprofloxacin increases ciprofloxacin serum concentrations.

Metoclopramide

Metoclopramide accelerates the absorption of ciprofloxacin (oral) resulting in a shorter time to reach maximum plasma concentrations. No effect was seen on the bioavailability of ciprofloxacin.

Omeprazole

Concomitant administration of ciprofloxacin and omeprazole containing medicinal products results in a slight reduction of C_{max} and AUC of ciprofloxacin.

Effects of ciprofloxacin on other medicinal products:

Agomelatine

In clinical studies, it was demonstrated that fluvoxamine, as a strong inhibitor of the CYP450 1A2 isoenzyme, markedly inhibits the metabolism of agomelatine resulting in a 60-fold increase of agomelatine exposure. Although no clinical data are available for a possible interaction with ciprofloxacin, a moderate inhibitor of CYP450 1A2, similar effects can be expected upon concomitant administration (see 'Cytochrome P450' in section 4.4).

Tizanidine

Tizanidine must not be administered together with ciprofloxacin (see section 4.3). In a clinical study with healthy subjects, there was an increase in serum tizanidine concentration (C_{max} increase: 7-fold, range: 4 to 21-fold; AUC increase: 10-fold, range: 6 to 24-fold) when given concomitantly with ciprofloxacin. Increased serum tizanidine concentration is associated with a potentiated hypotensive and sedative effect.

Methotrexate

Renal tubular transport of methotrexate may be inhibited by concomitant administration of ciprofloxacin, potentially leading to increased plasma levels of methotrexate and increased risk of methotrexate-associated toxic reactions. The concomitant use is not recommended (see section 4.4).

Theophylline

Concurrent administration of ciprofloxacin and theophylline can cause an undesirable increase in serum theophylline concentration. This can lead to theophylline-induced side effects that may rarely be life threatening or fatal. During the combination, serum theophylline concentrations should be checked and the theophylline dose reduced as necessary (see section 4.4).

Other xanthine derivatives

On concurrent administration of ciprofloxacin and caffeine or pentoxifylline (oxpentifylline), raised serum concentrations of these xanthine derivatives were reported.

Phenytoin

Simultaneous administration of ciprofloxacin and phenytoin may result in increased or reduced serum levels of phenytoin such that monitoring of drug levels is recommended.

Cyclosporin

A transient rise in the concentration of serum creatinine was observed when ciprofloxacin and cyclosporin containing medicinal products were administered simultaneously. Therefore, it is frequently (twice a week) necessary to control the serum creatinine concentrations in these patients.

Vitamin K antagonists

Simultaneous administration of ciprofloxacin **with a vitamin K antagonist** may augment its anti-coagulant effects. The risk may vary with the underlying infection, age and general status

of the patient so that the contribution of ciprofloxacin to the increase in INR (international normalised ratio) is difficult to assess. The INR should be monitored frequently during and shortly after co-administration of ciprofloxacin with a vitamin K antagonist (e.g., warfarin, acenocoumarol, phenprocoumon, or fluindione).

Duloxetine

In clinical studies, it was demonstrated that concomitant use of duloxetine with strong inhibitors of the CYP450 1A2 isozyme such as fluvoxamine, may result in an increase of AUC and C_{max} of duloxetine. Although no clinical data are available on a possible interaction with ciprofloxacin, similar effects can be expected upon concomitant administration (see section 4.4).

Ropinirole

It was shown in a clinical study that concomitant use of ropinirole with ciprofloxacin, a moderate inhibitor of the CYP450 1A2 isozyme, results in an increase of C_{max} and AUC of ropinirole by 60% and 84%, respectively. Monitoring of ropinirole-related side effects and dose adjustment as appropriate is recommended during and shortly after co-administration with ciprofloxacin (see section 4.4).

Lidocaine

It was demonstrated in healthy subjects that concomitant use of lidocaine containing medicinal products with ciprofloxacin, a moderate inhibitor of CYP450 1A2 isozyme, reduces clearance of intravenous lidocaine by 22%. Although lidocaine treatment was well tolerated, a possible interaction with ciprofloxacin associated with side effects may occur upon concomitant administration.

Clozapine

Following concomitant administration of 250 mg ciprofloxacin with clozapine for 7 days, serum concentrations of clozapine and N-desmethylclozapine were increased by 29% and 31%, respectively. Clinical surveillance and appropriate adjustment of clozapine dosage during and shortly after coadministration with ciprofloxacin are advised (see section 4.4).

Sildenafil

C_{max} and AUC of sildenafil were increased approximately twofold in healthy subjects after an oral dose of 50 mg given concomitantly with 500 mg ciprofloxacin. Therefore, caution should be used prescribing ciprofloxacin concomitantly with sildenafil taking into consideration the risks and the benefits.

Zolpidem

Co-administration of ciprofloxacin may increase blood levels of zolpidem, concurrent use is not recommended.

4.6 Pregnancy and lactation

Pregnancy

The data that are available on administration of ciprofloxacin to pregnant women indicates no malformative or fetoneonatal toxicity of ciprofloxacin. Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity. In juvenile and prenatal animals exposed to quinolones, effects on immature cartilage have been observed, thus, it cannot be excluded that the drug could cause damage to articular cartilage in the human immature organism / foetus (see section 5.3).

As a precautionary measure, it is preferable to avoid the use of ciprofloxacin during pregnancy.

Breast-feeding

Ciprofloxacin is excreted in breast milk. Due to the potential risk of articular damage, ciprofloxacin should not be used during breast-feeding.

4.7 Effects on ability to drive and use machines

Due to its neurological effects, ciprofloxacin may affect reaction time. Thus, the ability to drive or to operate machinery may be impaired.

* Cases of prolonged (up to months or years), disabling and potentially irreversible serious drug reactions affecting several, sometimes multiple, system organ classes and senses (including reactions such as tendonitis, tendon rupture, arthralgia, pain in extremities, gait disturbance, neuropathies associated with paraesthesia, , fatigue, psychiatric symptoms, memory impairment, and impairment of hearing, vision, taste and smell) have been reported in association with the use of quinolones and fluoroquinolones in some cases irrespective of pre-existing risk factors (see Section 4.4). A range of psychiatric symptoms may occur as part of these side effects, which may include, but are not necessarily limited to, sleep disorders, anxiety, panic attacks, confusion, or depression. There are no pharmacological treatments established to be effective treatments of the symptoms of long lasting or disabling side effects associated with fluoroquinolones. The frequency of these prolonged, disabling and potentially irreversible serious drug reactions cannot be estimated with precision using available data, but the reporting incidence from adverse drug reaction reports indicates the frequency is at minimum between 1/1,000 and 1/10,000 (corresponding to the Rare frequency category).

** Cases of aortic aneurysm and dissection, sometimes complicated by rupture (including fatal ones), and of regurgitation/incompetence of any of the heart valves have been reported in patients receiving fluoroquinolones (see section 4.4).

Paediatric population

The incidence of arthropathy (arthralgia, arthritis), mentioned above, is referring to data collected in studies with adults. In children, arthropathy is reported to occur commonly (see section 4.4).

4.8 Overdose

An overdose of 12 g has been reported to lead to mild symptoms of toxicity. An acute overdose of 16 g has been reported to cause acute renal failure.

Symptoms in overdose consist of dizziness, tremor, headache, tiredness, seizures, hallucinations, confusion, abdominal discomfort, renal and hepatic impairment as well as crystalluria and haematuria. Reversible renal toxicity has been reported.

Apart from routine emergency measures, e.g. ventricular emptying followed by medical carbon it is recommended to monitor renal function, including urinary pH and acidify, if required, to prevent crystalluria. Patients should be kept well hydrated. Calcium or magnesium containing antacids may theoretically reduce the absorption of ciprofloxacin in overdoses.

Only a small quantity of ciprofloxacin (<10%) is eliminated by haemodialysis or peritoneal dialysis.

In the event of overdose, symptomatic treatment should be implemented. ECG monitoring should be undertaken, because of the possibility of QT interval prolongation.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Fluoroquinolones, ATC code: J01MA02

Mechanism of action:

As a fluoroquinolone antibacterial agent, the bactericidal action of ciprofloxacin results from the inhibition of both type II topoisomerase (DNA-gyrase) and topoisomerase IV, required for bacterial DNA replication, transcription, repair and recombination.

Pharmacokinetic /Pharmacodynamic relationship:

Efficacy mainly depends on the relation between the maximum concentration in serum (C_{max}) and the minimum inhibitory concentration (MIC) of ciprofloxacin for a bacterial pathogen and the relation between the area under the curve (AUC) and the MIC.

Mechanism of resistance:

In-vitro resistance to ciprofloxacin can be acquired through a stepwise process by target site mutations in both DNA gyrase and topoisomerase IV. The degree of cross-resistance between ciprofloxacin and other fluoroquinolones that results is variable. Single mutations may not result in clinical resistance, but multiple mutations generally result in clinical resistance to many or all active substances within the class.

Impermeability and/or active substance efflux pump mechanisms of resistance may have a variable effect on susceptibility to fluoroquinolones, which depends on the physiochemical properties of the various active substances within the class and the affinity of transport systems for each active substance. All *in-vitro* mechanisms of resistance are commonly observed in clinical isolates. Resistance mechanisms that inactivate other antibiotics such as permeation barriers (common in *Pseudomonas aeruginosa*) and efflux mechanisms may affect susceptibility to ciprofloxacin. Plasmid-mediated resistance encoded by *qnr*-genes has been reported.

5.2 Pharmacokinetic properties

Absorption

Following oral administration of single doses of 250 mg, 500 mg, and 750 mg of ciprofloxacin tablets, ciprofloxacin is absorbed rapidly and extensively, mainly from the small intestine, reaching maximum serum concentrations 1-2 hours later.

Single doses of 100-750 mg produced dose-dependent maximum serum concentrations (C_{max}) between 0.56 and 3.7 mg/L. Serum concentrations increase proportionately with doses up to 1000 mg.

The absolute bioavailability is approximately 70-80%.

A 500 mg oral dose given every 12 hours has been shown to produce an area under the serum concentration-time curve (AUC) equivalent to that produced by an intravenous infusion of 400 mg ciprofloxacin given over 60 minutes every 12 hours.

Distribution

Protein binding of ciprofloxacin is low (20-30%). Ciprofloxacin is present in plasma largely in a non-ionised form and has a large steady state distribution volume of 2-3 L/kg body weight. Ciprofloxacin reaches high concentrations in a variety of tissues such as lung (epithelial fluid, alveolar macrophages, biopsy tissue), sinuses, inflamed lesions (cantharides blister fluid), and the urogenital tract (urine, prostate, endometrium) where total concentrations exceeding those of plasma concentrations are reached.

Biotransformation

Low concentrations of four metabolites have been reported, which were identified as: desethyleneciprofloxacin (M 1), sulphociprofloxacin (M 2), oxociprofloxacin (M 3) and formylciprofloxacin (M 4). The metabolites display *in-vitro* antimicrobial activity but to a lower degree than the parent compound.

Ciprofloxacin is known to be a moderate inhibitor of the CYP 450 1A2 iso-enzymes.

Elimination

Ciprofloxacin is largely excreted unchanged both renally and, to a smaller extent, faecally. The serum elimination half-life in subjects with normal renal function is approximately 4-7 hours.

Excretion of ciprofloxacin (% of dose)

	Oral Administration	
	Urine	Faeces
Ciprofloxacin	44.7	25.0
Metabolites (M1-M4)	11.3	7.5

Renal clearance is between 180-300 mL/kg/h and the total body clearance is between 480-600 mL/kg/h. Ciprofloxacin undergoes both glomerular filtration and tubular secretion. Severely impaired renal function leads to increased half lives of ciprofloxacin of up to 12 h. Non-renal clearance of ciprofloxacin is mainly due to active trans-intestinal secretion and metabolism. 1% of the dose is excreted via the biliary route. Ciprofloxacin is present in the bile in high concentrations.

Paediatric patients

The pharmacokinetic data in paediatric patients are limited.

Based on population pharmacokinetic analysis of paediatric patients with various infections, the predicted mean half-life in children is approx. 4-5 hours and the bioavailability of the oral suspension ranges from 50 to 80%.

5.3 Preclinical safety data

Non-clinical data reveal no special hazards for humans based on conventional studies of single dose toxicity, repeated dose toxicity, carcinogenic potential, or toxicity to reproduction. Like a number of other quinolones, ciprofloxacin is phototoxic in animals at clinically relevant exposure levels. Data on photomutagenicity/photocarcinogenicity show a weak photomutagenic or phototumorigenic effect of ciprofloxacin *in-vitro* and in animal experiments. This effect was comparable to that of other gyrase inhibitors.

Articular tolerability:

As reported for other gyrase inhibitors, ciprofloxacin causes damage to the large weight-bearing joints in immature animals. The extent of the cartilage damage varies according to age, species and dose; the damage can be reduced by taking the weight off the joints. Studies with mature animals (rat, dog) revealed no evidence of cartilage lesions. In a study in young beagle dogs, ciprofloxacin caused severe articular changes at therapeutic doses after two weeks of treatment, which were still observed after 5 months.

6. Pharmaceutical particulars

6.1 List of excipients

Tablet core:

Cellulose microcrystalline
Sodium starch glycolate (Type A)
Povidone (K 30)
Silica, colloidal anhydrous
Magnesium stearate

Film coating:

Hypromellose
Titanium dioxide (E 171)
Macrogol 400

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

4 years.

6.5 Nature and contents of container

Ciprofloxacin film-coated tablets are available in PVC/PVdC-Aluminum foil blister pack.

6.6 Special precautions for disposal and other handling

Any unused product or waste material should be disposed of in accordance with local requirements.

6.6 Special precautions for the disposal of unused veterinary medicinal products or waste materials

None.

7.0 APPLICANT/MANUFACTURER

7.1 Name and Address

EVERGREEN MORNING STAR PHARMACEUTICALS INDUSTRY LIMITED

NDIOWERRE ORLU L.G.A

ORLU, IMO STATE

7.2 Tel number

+234 703 872 9291

+234 813 693 3443

+234 803 364 9955

7.3 Email

egmpsecdept@gmail.com