

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

**1.3 PRODUCT INFORMATION**

**1.3.1 Summary of Product Characteristics (SmPC)**

**1. 1.Name of the medicinal product:**

**1.1 Product Name : ARTELUMEX TABLET**

**1.2 Strength : --**

**1.3 Pharmaceutical Dosage Form : Tablet**

**2. 2. Qualitative and Quantitative Composition**

**3.**

**Composition:**

Each film coated tablet contains:

Artemether.....20mg

Lumefantrine.....120mg

Excipients.....q.s

Colour: Lake of Tartrazine & Titanium Dioxide U.S.P

**List of Excipients:**

1. Croscarmellose Sodium BP
2. Hydroxypropylmethylcellulose USP (Methocel E5 Premium)
3. Polysorbate 80 BP (Silvosorb 8020)
4. Isopropyl Alcohol BP
5. Purified Water BP
6. Microcrystalline Cellulose BP (PH 112)
7. Colloidal Anhydrous Silica BP
8. Purified Talc BP
9. Magnesium Stearate BP
10. IC-MS-218 IH
11. Methylene Chloride
12. Wincoat Ready mix, WT-1875 Yellow

**3. Pharmaceutical Form:**

Yellow coloured round, biconvex film coated tablets with monogram “Ω” on one side and plain on other side.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

**4. Clinical Particulars**

**4.1 Therapeutic indications**

Treatment of malaria caused by all forms of Plasmodium including severe malaria caused by multiple drug resistant strains of P. falciparum.

**4.2 Posology and method of administration:**

Weight in Kg	Total Tablets	Dosage Regimen					
		Day 1		Day 2		Day 3	
		0 Hour	8 Hours	24 Hours	36 Hours	48 Hours	60 Hours
5 – 14	6	1	1	1	1	1	1
15 - 24	12	2	2	2	2	2	2
25 - 34	18	3	3	3	3	3	3
35 and more (Adults)	24	4	4	4	4	4	4

Second dose to be taken after 8 hours of first dose

Better take with food especially fatty meals

**Or**

**As directed by physician.**

**Method of administration:** Tablets for oral administration.

**4.3 Contraindications:**

Artelumex Tablets is contraindicated in:

- Patients with known hypersensitivity to the active substances or to any of the excipients.
- Patients with severe malaria according to WHO definition.
- Patients who are taking any drug which is metabolised by the cytochrome enzyme CYP2D6 (e.g. metoprolol, imipramine, amitriptyline, clomipramine).
- Patients with a family history of sudden death or of congenital prolongation of the QTc interval on electrocardiograms, or with any other clinical condition known to prolong the QTc interval.
- Patients taking drugs that are known to prolong the QTc interval (proarrhythmic). These drugs include:
  - antiarrhythmics of classes IA and III,
  - neuroleptics, antidepressive agents,
  - certain antibiotics including some agents of the following classes: macrolides, fluoroquinolones, imidazole and triazole antifungal agents,

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

- certain non-sedating antihistamines (terfenadine, astemizole),
- cisapride.
- flecainide
- patients with a history of symptomatic cardiac arrhythmias or with clinically relevant bradycardia or with congestive cardiac failure accompanied by reduced left ventricle ejection fraction.
- patients with disturbances of electrolyte balance e.g. hypokalemia or hypomagnesemia.
- patients taking drugs that are strong inducers of CYP3A4 such as rifampin, carbamazepine, phenytoin, St. John's wort (*Hypericum perforatum*).

(\*Presence of one or more of the following clinical or laboratory features:

Clinical manifestation: Prostration; impaired consciousness or unarousable coma; failure to feed; deep breathing, respiratory distress (acidotic breathing); multiple convulsions; circulatory collapse or shock; pulmonary edema (radiological); abnormal bleeding; clinical jaundice; hemoglobinuria

Laboratory test: Severe normocytic anemia; hemoglobuniuria; hypoglycemia; metabolic acidosis; renal impairment; hyperlactatemia; hyperparasitemia)

**4.4 Special warnings and precautions for use:**

Artelumex must not be used in the first trimester of pregnancy in situations where other suitable and effective antimalarials are available.

Artelumex has not been evaluated for the treatment of severe malaria, including cases of cerebral malaria or other severe manifestations such as pulmonary oedema or renal failure.

Due to limited data on safety and efficacy, Artelumex should not be given concurrently with any other antimalarial agent unless there is no other treatment option.

If a patient deteriorates whilst taking Artelumex, alternative treatment for malaria should be started without delay. In such cases, monitoring of the ECG is recommended and steps should be taken to correct any electrolyte disturbances.

The long elimination half-life of lumefantrine must be taken into account when administering quinine in patients previously treated with Artelumex.

If quinine is given after Artelumex, close monitoring of the ECG is advised.

If Artelumex is given after mefloquine, close monitoring of food intake is advised.

In patients previously treated with halofantrine, Artelumex should not be administered earlier than one month after the last halofantrine dose.

Artelumex is not indicated and has not been evaluated for prophylaxis.

Artelumex should be used cautiously in patients on ARTs since decreased artemether, DHA, and/or lumefantrine concentrations may result in a decrease of antimalarial efficacy of Artelumex.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

Like other antimalarials (e.g. halofantrine, quinine and quinidine) Artelumex has the potential to cause QT prolongation.

Caution is recommended when combining Artelumex with drugs exhibiting variable patterns of inhibition, moderate induction or competition for CYP3A4 as the therapeutic effects of some drugs could be altered. Drugs that have a mixed inhibitory/induction effect on CYP3A4, especially anti-retroviral drugs such as HIV protease inhibitors and non-nucleoside reverse transcriptase inhibitors should be used with caution in patients taking Artelumex.

Caution is recommended when combining Artelumex with hormonal contraceptives. Artelumex may reduce the effectiveness of hormonal contraceptives. Therefore, patients using oral, transdermal patch, or other systemic hormonal contraceptives should be advised to use an additional non-hormonal method of birth control for about one month.

Patients who remain averse to food during treatment should be closely monitored as the risk of recrudescence may be greater.

In patients with severe hepatic impairment, a clinically relevant increase of exposure to artemether and lumefantrine and/or their metabolites cannot be ruled out. Therefore caution should be exercised in dosing patients with severe hepatic impairment.

**Renal impairment**

No specific studies have been carried out in this group of patients. There is no significant renal excretion of lumefantrine, artemether and dihydroartemisinin in studies conducted in healthy volunteers and clinical experience is limited. No dose adjustment for the use of Artelumex in patients with renal impairment is recommended. Caution is advised when administering Artelumex to patients with severe renal impairment. In these patients, ECG and blood potassium monitoring is advised.

**Hepatic impairment**

No specific studies have been carried out in this group of patients. No dose adjustment is recommended for patients with mild to moderate hepatic impairment. Caution is advised when administering Artelumex to patients with severe hepatic impairment. In these patients, ECG and blood potassium monitoring is advised.

**Elderly**

There is no information suggesting that the dosage in patients over 65 years of age should be different than in younger adults.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

**New infections**

Data for a limited number of patients in a malaria endemic area show that new infections can be treated with a second course of Artelumex. In the absence of carcinogenicity study data, and due to lack of clinical experience, more than two courses of Artelumex cannot be recommended.

**4.5 Interaction with other medicinal products and other forms of interaction**

**Contraindications of concomitant use**

**Interaction with drugs that are known to prolong the QTc interval**

Artelumex is contraindicated with concomitant use of drugs (they may cause prolonged QTc interval and Torsade de Pointes) such as: antiarrhythmics of classes IA and III, neuroleptics and antidepressant agents, certain antibiotics including some agents of the following classes: macrolides, fluoroquinolones, imidazole, and triazole antifungal agents, certain non-sedating antihistaminics (terfenadine, astemizole), cisapride, flecainide.

**Interaction with drugs metabolized by CYP2D6**

Lumefantrine was found to inhibit CYP2D6 in vitro. This may be of particular clinical relevance for compounds with a low therapeutic index. Co-administration of Artelumex with drugs that are metabolised by this iso-enzyme is contraindicated (e.g. neuroleptics, metoprolol, and tricyclic antidepressants such as imipramine, amitriptyline, clomipramine) is contraindicated.

**Interaction with strong inducers of CYP3A4 such as rifampin**

Oral administration of rifampin (600 mg daily), a strong CYP3A4 inducer, with Artelumex Tablets (6-dose regimen over 3 days) in six HIV-1 and tuberculosis co-infected adults without malaria resulted in significant decreases in exposure to artemether (89%), DHA (85%) and lumefantrine (68%) when compared to exposure values after Artelumex alone. Concomitant use of strong inducers of CYP3A4 such as rifampin, carbamazepine, phenytoin, St. John's Wort is contraindicated with Artelumex.

Inducers should not be administered at least one month after Artelumex administration, unless critical to use as judged by the prescriber.

**Concomitant use not recommended**

**Interaction with other antimalarial drugs**

Data on safety and efficacy are limited, and Artelumex should therefore not be given concurrently with other antimalarials unless there is no other treatment option.

If Artelumex is given following administration of mefloquine or quinine, close monitoring of food intake (for mefloquine) or of the ECG (for quinine) is advised. The long elimination half-life of lumefantrine must be taken into account when administering quinine in patients previously

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

treated with Artelumex. In patients previously treated with halofantrine, Artelumex should not be administered earlier than one month after the last halofantrine dose.

Mefloquine

A drug interaction study with Artelumex in man involved administration of a 6-dose regimen over 60 hours in healthy volunteers which was commenced at 12 hours after completion of a 3-dose regimen of mefloquine or placebo. Plasma mefloquine concentrations from the time of addition of Artelumex were not affected compared with a group which received mefloquine followed by placebo.

Pre-treatment with mefloquine had no effect on plasma concentrations of artemether or the artemether/dihydroartemisinin ratio but there was a significant reduction in plasma levels of lumefantrine, possibly due to lower absorption secondary to a mefloquine-induced decrease in bile production. Patients should be encouraged to eat at dosing times to compensate for the decrease in bioavailability.

Quinine

A drug interaction study in healthy male volunteers showed that the plasma concentrations of lumefantrine and quinine were not affected when i.v. quinine (10 mg/kg BW over 2 hours) was given sequentially 2 hours after the last (sixth) dose of Artelumex (so as to produce concurrent plasma peak levels of lumefantrine and quinine). Plasma concentrations of artemether and dihydroartemisinin (DHA) appeared to be lower. In this study, administration of Artelumex to 14 subjects had no effect on QTc interval. Infusion of quinine alone in 14 other subjects caused a transient prolongation of QTc interval, which was consistent with the known cardiotoxicity of quinine. This effect was slightly, but significantly, greater when quinine was infused after Artelumex in 14 additional subjects. It would thus appear that the inherent risk of QTc prolongation associated with i.v. quinine was enhanced by prior administration of Artelumex.

**Concomitant use requiring caution**

Interactions affecting the use of Artelumex

Interaction with CYP3A4 inhibitors

Both artemether and lumefantrine are metabolised predominantly by the cytochrome enzyme CYP3A4, but do not inhibit this enzyme at therapeutic concentrations.

Ketoconazole

The concurrent oral administration of ketoconazole with Artelumex led to a modest increase ( $\leq$  2-fold) in artemether, DHA, and lumefantrine exposure in healthy adult subjects. This increase in exposure to the antimalarial combination was not associated with increased side effects or changes in electrocardiographic parameters. Based on this study, dose adjustment of Artelumex

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

is considered unnecessary in falciparum malaria patients when administered in association with ketoconazole or other potent CYP3A4 inhibitors.

Artelumex should be used cautiously with drugs that inhibit CYP3A4 and are contraindicated with drugs which additionally are known to prolong QTc, due to potential for increased concentrations of lumefantrine which could lead to QT prolongation.

Grapefruit juice

Administration of artemether with grapefruit juice in healthy adult subjects resulted in an approximately two fold increase in systemic exposure to the parent drug. Grapefruit juice should be used cautiously during Artelumex treatment.

Interaction with weak to moderate inducers of CYP3A4

When Artelumex is co-administered with moderate inducers of CYP3A4, it may result in decreased concentrations of artemether and/or lumefantrine and loss of antimalarial efficacy.

Interaction with anti-retroviral drugs such as protease inhibitors and non-nucleoside reverse transcriptase inhibitors

Both artemether and lumefantrine are metabolized by CYP3A4. Anti-retroviral drugs (ARTs), such as protease inhibitors and non-nucleoside reverse transcriptase inhibitors, are known to have variable patterns of inhibition, induction or competition for CYP3A4. Artelumex should be used cautiously in patients on ARTs since decreased artemether, DHA, and/or lumefantrine concentrations may result in a decrease of antimalarial efficacy of Artelumex, and increased lumefantrine concentrations may cause QT prolongation.

Lopinavir/ ritonavir

In a clinical study in healthy volunteers, lopinavir/ritonavir decreased the systemic exposures to artemether and DHA by approximately 40% but increased the exposure to lumefantrine by approximately 2.3- fold. Exposures to lopinavir/ritonavir were not significantly affected by concomitant use of Artelumex.

Nevirapine

In a clinical study in HIV-infected adults, nevirapine significantly reduced the median Cmax and AUC of artemether by approximately 61% and 72%, respectively and reduced the median Cmax and AUC of dihydroartemisinin by approximately 45% and 37%, respectively. Lumefantrine Cmax and AUC were non-significantly reduced by nevirapine. Artemether/lumefantrine reduced the median Cmax and AUC of nevirapine by approximately 43% and 46% respectively.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

Efavirenz

Efavirenz decreased the exposures to artemether, DHA, and lumefantrine by approximately 50%, 45%, and 20%, respectively. Exposures to efavirenz were not significantly affected by concomitant use of Artelumex.

**Interactions resulting in effects of Artelumex on other drugs**

Interaction with drugs metabolized by CYP450 enzymes

When Artelumex is co-administered with substrates of CYP3A4 it may result in decreased concentrations of the substrate and potential loss of substrate efficacy. Studies in humans have demonstrated that Artemisinin has some capacity to induce CYP3A4 and CYP2C19 and inhibit CYP2D6 and CYP1A2. Although the magnitude of the changes was generally low it is possible that these effects could alter the therapeutic response of drugs that are predominantly metabolised by these enzymes.

Interaction with hormonal contraceptives

In vitro, the metabolism of ethinyl estradiol and levonorgestrel was not induced by artemether, DHA, or lumefantrine. However, artemether has been reported to weakly induce, in humans, the activity of CYP2C19, CYP2B6, and CYP3A. Therefore, Artelumex may potentially reduce the effectiveness of hormonal contraceptives. Patients using oral, transdermal patch, or other systemic hormonal contraceptives should be advised to use an additional nonhormonal method of birth control for about one month.

Drug-food/drink interactions

Artelumex should be taken with food or drinks rich in fat such as milk as the absorption of both artemether and lumefantrine is increased.

Grapefruit juice should be used cautiously during Artelumex treatment.

**4.6 Pregnancy and Lactation:**

Pregnancy

There is insufficient data from the use of artemether and lumefantrine in pregnant women. Based on animal data, Artelumex is suspected to cause serious birth defects when administered during the first trimester of pregnancy. Reproductive studies with artemether have shown evidence of post-implantation losses and teratogenicity in rats and rabbits. Other artemisinin derivatives have also demonstrated teratogenic potential with an increased risk during early gestation.



**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

Safety data from an observational pregnancy study of approximately 500 pregnant women who were exposed to Artelumex (including a third of patients who were exposed in the first trimester), and published data of another over 500 pregnant women who were exposed to artemether- lumefantrine (including over 50 patients who were exposed in the first trimester), as well as published data of over 1,000 pregnant women who were exposed to artemisinin derivatives, did not show an increase in adverse pregnancy outcomes or teratogenic effects over background rates.

Artelumex treatment must not be used during the first trimester of pregnancy in situations where other suitable and effective antimalarials are available. However, it should not be withheld in life-threatening situations, where no other effective antimalarials are available. During the second and third trimester, treatment should only be considered if the expected benefit to the mother outweighs the risk to the foetus.

Women of childbearing potential

Women using oral, transdermal patch, or other systemic hormonal contraceptives should be advised to use an additional non-hormonal method of birth control for about one month.

Lactation

Animal data suggest excretion into breast milk but no data are available in humans. Women taking Artelumex should not breast-feed during their treatment. Due to the long elimination half-life of lumefantrine (2 to 6 days), it is recommended that breast-feeding should not resume until at least one week after the last dose of Artelumex unless potential benefits to the mother and child outweigh the risks of Artelumex treatment.

Fertility

There is no information on the effects of Artelumex on human fertility.

**4.7 Effects on ability to drive and use machines**

Patients receiving Artelumex should be warned that dizziness or fatigue/asthenia may occur in which case they should not drive or use machines.

**4.8 Undesirable Effects:**

With Artemether virtually no side effects have been seen. Laboratory abnormalities such as slight rise in transaminases and a decrease in reticulocyte count are rare and transient. A lowering of sinus frequency without causing ECG changes has been noticed. At high doses transient abdominal pain, tinnitus and diarrhea have been described but a causal relationship is unclear.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

Some antimalarials such as Halofantrine and Quinine can influence the ECG pattern. Attention should be made to patient previously treated with those antimalarials. A reasonable period should be taken into account before starting treatment with Lumefantrine combinations. For such patients, mono therapy with Artemisinin derivatives is recommended.

Sometimes rash, trouble sleeping, nausea, vomiting, diarrhea, coughing may occur. They need medical attention when persisting.

**4.9 Overdosage:**

In cases of suspected overdosage symptomatic and supportive therapy should be given as appropriate, which should include ECG and blood potassium monitoring.

**5. Pharmacological properties:**

**5.1 Pharmacodynamic properties:**

Pharmacotherapeutic group: antimalarials, blood schizontocide,  
ATC code: P01 BF01.

Pharmacodynamic effects

Artelumex comprises a fixed ratio of 1:6 parts of artemether and lumefantrine, respectively. The site of antiparasitic action of both components is the food vacuole of the malarial parasite, where they are thought to interfere with the conversion of haem, a toxic intermediate produced during haemoglobin breakdown, to the nontoxic haemozoin, malaria pigment. Lumefantrine is thought to interfere with the polymerisation process, while artemether generates reactive metabolites as a result of the interaction between its peroxide bridge and haem iron. Both artemether and lumefantrine have a secondary action involving inhibition of nucleic acid- and protein synthesis within the malarial parasite.

Treatment of Acute Uncomplicated P. falciparum Malaria

The efficacy of Artelumex Tablets was evaluated for the treatment of acute, uncomplicated malaria (defined as symptomatic P. falciparum malaria without signs and symptoms of severe malaria or evidence of vital organ dysfunction) in five 6-dose regimen studies and one study comparing the 6-dose regimen with the 4-dose regimen. Baseline parasite density ranged from 500/μL - 200,000/μL (0.01% to 4% parasitemia) in the majority of patients. Studies were conducted in otherwise healthy, partially immune or non-immune adults and children (≥5kg body weight) with uncomplicated malaria in Thailand, sub-Saharan Africa, Europe, and South America.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

Efficacy endpoints consisted of:

- 28-day cure rate, proportion of patients with clearance of asexual parasites within 7 days without recrudescence by day 28
- parasite clearance time (PCT), defined as time from first dose until first total and continued disappearance of asexual parasite which continues for a further 48 hours
- fever clearance time (FCT), defined as time from first dose until the first time body temperature fell below 37.5°C and remained below 37.5°C for at least a further 48 hours (only for patients with temperature >37.5°C at baseline)

The modified intent to treat (mITT) population includes all patients with malaria diagnosis confirmation who received at least one dose of study drug. Evaluable patients generally are all patients who had a day 7 and a day 28 parasitological assessment or experienced treatment failure by day 28. The results are presented in the table below:

**Table 2 Clinical efficacy results**

<b>Study No.</b>	<b>Age</b>	<b>Polymerase chain reaction (PCR)-corrected 28-day cure rate<sup>1</sup> n/N (%) in evaluable patients</b>	<b>Median FCT<sup>2</sup> [25<sup>th</sup>, 75<sup>th</sup>percentile]</b>	<b>Median PCT<sup>2</sup> [25<sup>th</sup>, 75<sup>th</sup>percentile]</b>	<b>Year/ Study location</b>
A025 <sup>4</sup>	3-62 years	93/96 (96.9)	n <sup>3</sup> =59 35 hours [20, 46]	n=118 44 hours [22, 47]	1996-97 Thailand
A026	2-63 years	130/133 (97.7)	n <sup>3</sup> =87 22 hours [19, 44]	NA	1997-98 Thailand
A028	12-71 years	148/154 (96.1)	n <sup>3</sup> =76 29 hours [8, 51]	n=164 29 hours [18, 40]	1998-99 Thailand
A2401	16-66 years	119/124 (96.0)	n <sup>3</sup> =100 37 hours [18, 44]	n=162 42 hours [34, 63]	2001-05 Europe, Columbia
A2403	2 months-9 years	289/299 (96.7)	n <sup>3</sup> =309 8 hours [8, 24]	n=310 24 hours [24, 36]	2002-03 3 countries in Africa
B2303 <sup>CT</sup>	3 months-12 years	403/419 (96.2)	n <sup>3</sup> =323 8 hours [8, 23]	n=452 35 hours [24, 36]	2006-07 5 countries in Africa

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

B2303 <sup>DT</sup>	3 months-12 years	394/416 (94.7)	n <sup>3</sup> =311 8 hours [8, 24]	n=446 34 hours [24, 36]	2006-07 5 countries in Africa
---------------------	-------------------	----------------	--	----------------------------	----------------------------------

<sup>1</sup> Efficacy cure rate based on blood smear microscopy

<sup>2</sup> mITT population

<sup>3</sup> For patients who had a body temperature >37.5°C at baseline only

<sup>4</sup> Only the 6-dose regimen over 60 hours group data is presented

<sup>CT</sup> –Artelumex tablets administered as crushed tablets

<sup>DT</sup> –Artelumex Dispersible tablets

Artelumex is not indicated for, and has not been evaluated in, the treatment of malaria due to *P. vivax*, *P. malariae* or *P. ovale*, although some patients in clinical studies had co-infection with *P. falciparum* and *P. vivax* at baseline. Artelumex is active against blood stages of *Plasmodium vivax*, but is not active against hypnozoites.

Paediatric population

Two studies have been conducted

Study A2403 was conducted in Africa in 310 infants and children aged 2 months to 9 years, weighing 5 kg to 25 kg, with an axillary temperature  $\geq 37.5^\circ\text{C}$ . Results of 28-day cure rate (PCR-corrected), median parasite clearance time (PCT), and fever clearance time (FCT) are reported in table 3 below.

Study B2303 was conducted in Africa in 452 infants and children, aged 3 months to 12 years, weighing 5 kg to <35 kg, with fever ( $\geq 37.5^\circ\text{C}$  axillary or  $\geq 38^\circ\text{C}$  rectally) or history of fever in the preceding 24 hours. This study compared crushed tablets and dispersible tablets. Results of 28-day cure rate (PCR-corrected), median parasite clearance time (PCT), and fever clearance time (FCT) for crushed tablets are reported in table 3 below.

**Table 3 Clinical efficacy by weight for pediatric studies**

Study No. Weight category	Median PCT <sup>1</sup> [25 <sup>th</sup> , 75 <sup>th</sup> percentile]	PCR-corrected 28-day cure rate <sup>2</sup> n/N (%) in evaluable patients
Study A2403		
5 - <10 kg	24 hours [24, 36]	145/149 (97.3)
10 - <15 kg	35 hours [24, 36]	103/107 (96.3)
15 -25 kg	24 hours [24, 36]	41/43 (95.3)
Study B2303 <sup>CT</sup>		
5 - <10 kg	36 hours [24, 36]	65/69 (94.2)
10 - <15 kg	35 hours [24, 36]	174/179 (97.2)
15 -<25 kg	35 hours [24, 36]	134/140 (95.7)
25-35 kg	26 hours [24, 36]	30/31 (96.8)

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

<sup>1</sup> mITT population

<sup>2</sup> Efficacy cure rate based on blood smear microscopy

<sup>CT</sup> Artelumex tablets administered as crushed tablets

**QT/QTc Prolongation:**

Adults and children with malaria

For information on the risk of QT/QTc prolongation in patients.

**Healthy adults**

In a healthy adult volunteer parallel group study including a placebo and moxifloxacin control group (n=42 per group), the administration of the six dose regimen of Artelumex was associated with prolongation of QTcF. The mean changes from baseline at 68, 72, 96, and 108 hours post first dose were 7.45, 7.29, 6.12 and 6.84 msec, respectively. At 156 and 168 hours after first dose, the changes from baseline for QTcF had no difference from zero. No subject had a >30 msec increase from baseline nor an absolute increase to >500 msec. Moxifloxacin control was associated with a QTcF increase as compared to placebo for 12 hours after the single dose with a maximal change at 1 hour after dose of 14.1 msec.

In the adult/adolescent population included in clinical trials, 8 patients (0.8%) receiving Artelumex experienced a QTcB >500 msec and 3 patients (0.4%) a QTcF >500 msec. Prolongation of QTcF interval >30 msec was observed in 36% of patients.

In clinical trials conducted in children with the 6-dose regimen, no patient had post-baseline QTcF >500 msec whereas 29.4% had QTcF increase from baseline >30 msec and 5.1% >60 msec. In clinical trials conducted in adults and adolescents with the 6-dose regimen, post-baseline QTcF prolongation of >500 msec was reported in 0.2% of patients, whereas QTcF increase from baseline >30 msec was reported in 33.9% and >60 msec in 6.2% of patients.

In the infant/children population included in clinical trials, 3 patients (0.2%) experienced a QTcB >500 msec. No patient had QTcF >500 msec. Prolongation of QTcF intervals >30 msec was observed in 34% of children weighing 5-10 kg, 31% of children weighing 10-15 kg and 24% of children weighing 15-25 kg, and 32% of children weighing 25-35 kg.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

**5.2 Pharmacokinetic Properties**

Pharmacokinetic characterisation of Artelumex is limited by the lack of an intravenous formulation, and the very high inter-and intra-subject variability of artemether and lumefantrine plasma concentrations and derived pharmacokinetic parameters (AUC, Cmax).

**Absorption**

Artemether is absorbed fairly rapidly and dihydroartemisinin, the active metabolite of artemether, appears rapidly in the systemic circulation with peak plasma concentrations of both compounds reached about 2 hours after dosing. Mean Cmax and AUC values of artemether ranged between 60.0-104 ng/mL and 146-338 ng·h/mL, respectively, in fed healthy adults after a single dose of Artelumex, 80 mg artemether/480 mg lumefantrine. Mean Cmax and AUC values of dihydroartemisinin ranged between 49.7-104 ng/mL and 169-308 ng·h/mL, respectively. Absorption of lumefantrine, a highly lipophilic compound, starts after a lag-time of up to 2 hours, with peak plasma concentration (mean between 5.10-9.80 µg/mL) about 6-8 hours after dosing. Mean AUC values of lumefantrine ranged between 108 and 243 µg·h/mL. Food enhances the absorption of both artemether and lumefantrine: in healthy volunteers the relative bioavailability of artemether was increased more than two-fold, and that of lumefantrine sixteen-fold compared with fasted conditions when Artelumex was taken after a high-fat meal.

Food has also been shown to increase the absorption of lumefantrine in patients with malaria, although to a lesser extent (approximately two-fold), most probably due to the lower fat content of the food ingested by acutely ill patients. The food interaction data indicate that absorption of lumefantrine under fasted conditions is very poor (assuming 100% absorption after a high-fat meal, the amount absorbed under fasted conditions would be <10% of the dose). Patients should therefore be encouraged to take the medication with a normal diet as soon as food can be tolerated.

**Distribution**

Artemether and lumefantrine are both highly bound to human serum proteins *in vitro* (95.4% and 99.7%, respectively). Dihydroartemisinin is also bound to human serum proteins (47-76%).

**Metabolism**

Artemether is rapidly and extensively metabolised (substantial first-pass metabolism) both *in vitro* and in humans. Human liver microsomes metabolise artemether to the biologically active main metabolite dihydroartemisinin (demethylation), predominantly through the isoenzyme CYP3A4/5. This metabolite has also been detected in humans *in vivo*.

Dihydroartemisinin is further converted to inactive metabolites.

The pharmacokinetics of artemether in adults is time-dependent. During repeated administration of Artelumex, plasma artemether levels decreased significantly, while levels of the active metabolite (dihydroartemisinin) increased, although not to a statistically significant degree. The

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

ratio of day 3/day 1 AUC for artemether was between 0.19 and 0.44, and was between 1.06 and 2.50 for dihydroartemisinin. This suggests that there was induction of the enzyme responsible for the metabolism of artemether. Artemether and dihydroartemisinin were reported to have a mild inducing effect on CYP3A4 activity.

Lumefantrine is N-debutylated, mainly by CYP3A4, in human liver microsomes. *In vivo* in animals (dogs and rats), glucuronidation of lumefantrine takes place directly and after oxidative biotransformation. In humans, the exposure to lumefantrine increases with repeated administration of Artelumex over the 3-day treatment period, consistent with the slow elimination of the compound (see section 5.2 Elimination). Systemic exposure to the metabolite desbutyl-lumefantrine, for which the *in vitro* antiparasitic effect is 5 to 8 fold higher than that for lumefantrine, was less than 1% of the exposure to the parent drug. Desbutyl-lumefantrine data is not available specifically for an African population. *In vitro*, lumefantrine significantly inhibits the activity of CYP2D6 at therapeutic plasma concentrations.

Elimination

Artemether and dihydroartemisinin are rapidly cleared from plasma with a terminal half-life of about 2 hours. Lumefantrine is eliminated very slowly with an elimination half-life of 2 to 6 days. Demographic characteristics such as sex and weight appear to have no clinically relevant effects on the pharmacokinetics of Artelumex.

Limited urinary excretion data are available for humans. In 16 healthy volunteers, neither lumefantrine nor artemether was found in urine after administration of Artelumex, and only traces of dihydroartemisinin were detected (urinary excretion of dihydroartemisinin amounted to less than 0.01% of the artemether dose).

In animals (rats and dogs), no unchanged artemether was detected in faeces and urine due to its rapid and extensive first-pass metabolism, but numerous metabolites (partly identified) have been detected in faeces, bile and urine. Lumefantrine was excreted unchanged in faeces and with traces only in urine. Metabolites of lumefantrine were eliminated in bile/faeces.

Dose proportionality

No specific dose proportionality studies were performed. Limited data suggest a dose-proportional increase of systemic exposure to lumefantrine when doubling the Artelumex dose. No conclusive data is available for artemether.

Bioavailability/bioequivalence studies

Systemic exposure to lumefantrine, artemether and dihydroartemisinin was similar following administration of Artelumex as dispersible tablets and crushed tablets in healthy adults.

Systemic exposure to lumefantrine was similar following administration of Artelumex dispersible tablets and intact tablets in healthy adults. However, exposure to artemether and

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

dihydroartemisinin was significantly lower (by 20-35%) for the dispersible than for the intact tablet. These findings are not considered to be clinically relevant for the use of the dispersible tablets in the paediatric population since adequate efficacy of Artelumex dispersible tablets was demonstrated in this population. The dispersible tablet is not recommended for use in adults.

Special populations

No specific pharmacokinetic studies have been performed in elderly patients. However, there is no information suggesting that the dosage in patients over 65 years of age should be different than in younger adults.

In paediatric malaria patients, mean C<sub>max</sub> (CV%) of artemether (observed after first dose of Artelumex) were 223 (139%), 198 (90%) and 174 ng/mL (83%) for body weight groups 5-<15, 15-<25 and 25-<35 kg, respectively, compared to 186 ng/mL (67%) in adult malaria patients. The associated mean C<sub>max</sub> of DHA were 54.7 (108%), 79.8 (101%) and 65.3 ng/mL (36%), respectively compared to 101 ng/mL (57%) in adult malaria patients. AUC of lumefantrine (population mean, covering the six doses of Artelumex) were 577, 699 and 1150 µg•h/mL for paediatric malaria patients in body weight groups 5-<15, 15-<25 and 25-<35 kg, respectively, compared to a mean AUC of 758 µg•h/mL (87%) in adult malaria patients. The elimination half-lives of artemether and lumefantrine in children are unknown.

No specific pharmacokinetic studies have been performed either in patients with hepatic or renal insufficiency or elderly patients. The primary clearance mechanism of both artemether and lumefantrine may be affected in patients with hepatic impairment. In patients with severe hepatic impairment, a clinically significant increase of exposure to artemether and lumefantrine and/or their metabolites cannot be ruled out. Therefore caution should be exercised in dosing patients with severe hepatic impairment. Based on the pharmacokinetic data in 16 healthy subjects showing no or insignificant renal excretion of lumefantrine, artemether and dihydroartemisinin, no dose adjustment for the use of Artelumex in patients with renal impairment is advised.

**5.3 Preclinical safety data**

General toxicity

The main changes observed in repeat-dose toxicity studies were associated with the expected pharmacological action on erythrocytes, accompanied by responsive secondary haematopoiesis.

Neurotoxicity

Studies in dogs and rats have shown that intramuscular injections of artemether resulted in brain lesions. Changes observed mainly in brainstem nuclei included chromatolysis, eosinophilic cytoplasmic granulation, spheroids, apoptosis and dark neurons. Lesions were observed in rats dosed with artemether at 25 mg/kg for 7 or 14 days and dogs dosed at 20 mg/kg for 8 days or longer, but lesions were not observed after shorter courses of drug or after oral dosing. The



**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

estimated artemether 24 h AUC after 7 days of dosing at the no observed effect level (10 mg/kg/day given intramuscularly) is approximately 7-fold greater than the estimated artemether 24 h AUC in humans on day 1 of the standard 3-day oral treatment regimen; oral exposure in humans decreases on subsequent days, thus the exposure margin increases. Dogs dosed orally with 143 mg/kg artemether showed a statistically measureable effect on the hearing threshold at 20 dB. This dose is equivalent to about 29 times the highest artemether clinical dose (160 mg/day) based on body surface area comparisons. Most nervous system disorder adverse events in the studies of the 6-dose regimen were mild in intensity and resolved by the end of the study.

Mutagenicity

No evidence of mutagenicity was detected in *in vitro* or *in vivo* tests with an artemether:lumefantrine combination (consisting of 1 part artemether:6 parts lumefantrine). In the micronucleus test myelotoxicity was seen at all dose levels (500, 1,000 and 2,000 mg/kg), but recovery was almost complete 48 hours after dosing.

Carcinogenicity

Carcinogenicity studies with the artemether:lumefantrine combination were not conducted.

Reproductive toxicity studies

Reproductive toxicity studies performed with the artemether:lumefantrine combination caused maternal toxicity and increased post-implantation loss in rats and rabbits at doses  $\geq 50$  mg/kg/day (corresponding to approximately 7 mg/kg/day artemether) and 175 mg/kg/day (corresponding to 25 mg/kg/day artemether) respectively. These effects were not observed at lower doses.

Lumefantrine alone caused no sign of reproductive or development toxicity at doses up to 1,000 mg/kg/day in rats and rabbits.

Embryotoxicity has been observed in rat and rabbit reproductive toxicity studies conducted with artemether, a derivative of artemisinin. Artemisinins (e.g. artesunate) are known to be embryotoxic.

Artemether caused increases in post-implantation loss and teratogenicity (characterised as a low incidence of cardiovascular and skeletal malformations) in rats at 19.4 mg/kg, and in rabbits at 30 mg/kg. Maternal toxicity was also observed in rabbits at 30 mg/kg/day. No other adverse effects were observed at lower doses in rabbits. The no observed effect dose was 3 mg/kg/day in rats and 25 mg/kg/day in rabbits.

The embryotoxic artemether dose, 20 mg/kg/day in the rat, yields artemether and dihydroartemisinin exposures similar to those achieved in humans.

Artesunate, a structurally related compound, also caused increases in post-implantation loss and teratogenicity (low incidence of cardiovascular and skeletal malformations) in rats at 6 mg/kg and in the lowest dose tested in the rabbits, 5 mg/kg/day.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

Fertility

After artemether-lumefantrine administration for 10 weeks in males and 2 weeks in females, reduced fertility occurred at 1000 mg/kg/day where altered sperm motility, abnormal sperm, reduced epididymal sperm count, increased testes weight, and embryotoxicity and other reproductive effects (decreased implants and viable embryos, increased preimplantation loss) were also observed. General toxicity was observed in males and females at doses  $\geq$  300 mg/kg/day. The no adverse effect level for fertility was 300 mg/kg/day. The relevance to this finding in humans is unknown.

Juvenile toxicity studies

A specific study to investigate the neurotoxicity of artemether in juvenile rats involved oral administration of artemether during four different dosing intervals, at doses of 30 or 80 mg/kg/day on post-partum days 7 to 13, and at doses of 30 or 120 mg/kg/day on post-partum days 14 to 21, 22 to 28, or 29 to 36. Mortality, clinical signs and reductions in body weight parameters occurred most notably during the first two dosing intervals. Despite the systemic toxicity noted, there were no effects of artemether on any of the functional tests performed and there was no evidence of a direct neurotoxic effect of orally administered artemether on the brain of juvenile rats.

Juvenile studies in the rat indicate that very young animals (aged 7-21 days) are more sensitive to artemether than adult animals. There is no difference in sensitivity in slightly older (3-5 weeks of age) animals following 13 weeks of artemether/lumefantrine administration. Consistent with the later data, clinical studies have established the safety of artemether and lumefantrine administration in patients weighing 5 kg and above.

Cardiovascular Safety Pharmacology

In toxicity studies in dogs at doses  $\geq$ 600 mg/kg/day only, there was some evidence of prolongation of the QTc interval (safety margin of 1.3-fold to 2.2-fold for artemether using calculated free C<sub>max</sub>), at higher doses than intended for use in man. In an *in vitro* assay of HERG channels stably expressed in HEK293 cells, lumefantrine and the main metabolite desbutyl-lumefantrine showed some inhibitory potential in one of the currents responsible for cardiac repolarization. The potency was lower than the other antimalarial drugs tested. From the estimated IC<sub>50</sub> values, the order of potency of HERG current block was halofantrine (IC<sub>50</sub> = 0.04  $\mu$ M) >chloroquine (2.5  $\mu$ M) >mefloquine 2.6  $\mu$ M) >desbutyl-lumefantrine (5.5  $\mu$ M) >lumefantrine (8.1  $\mu$ M).

Additional studies were performed to evaluate the *in vitro* effects of artemether and its active metabolite, dihydroartemisinin, on the HERG current. At concentrations that produced significant inhibition, the safety margins for artemether and dihydroartemisinin are greater than 100 if they are estimated using the total therapeutic concentration at C<sub>max</sub> or greater than 1000 if

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

they are estimated using the calculated free C<sub>max</sub>. Based on the available non-clinical data, a potential for QTc prolongation in the human cannot be discounted.

**6. Pharmaceutical Particulars**

**6.1 List of Excipients**

1. Croscarmellose Sodium BP
2. Hydroxypropylmethylcellulose USP (Methocel E5 Premium)
3. Polysorbate 80 BP (Silvosorb 8020)
4. Isopropyl Alcohol BP
5. Purified Water BP
6. Microcrystalline Cellulose BP (PH 112)
7. Colloidal Anhydrous Silica BP
8. Purified Talc BP
9. Magnesium Stearate BP
10. IC-MS-218 IH
11. Methylene Chloride
12. Wincoat Ready mix, WT-1875 Yellow

**BP: British Pharmacopoeia**

**USP-NF: United State Pharmacopoeia-National Formulary**

**MS: Meyer Specifications**

**6.2 Incompatibilities:**

Not applicable.

**6.3 Shelf life:**

36 months from the date of manufacture.

**6.4 Special precautions for storage:**

Store below 30°C, at a dry place.

Protect from light.

Keep the medicine out of reach of children.

**6.5 Nature and contents of container:**

Each Carton contains **1 blister strip**, each containing **24 no. of ARTELUMEX TABLETS**.

**6.6 Special precautions for disposal and other handling:**

No data available.

**MODULE -1**  
**ADMINISTRATIVE AND PRODUCT INFORMATION**  
**ARTELUMEX TABLET**

**7. MANUFACTURER**

**Company Name:** Medibios Laboratories Pvt. Ltd.

**Address:** Plot No. J-76, MIDC, Tarapur, Taluka – Palghar, Dist. Thane 401 506, Maharashtra,  
INDIA

**Country:** INDIA.

**EXPORTED BY**

**Company Name:** Meyer Organics Pvt. Ltd.

**Address:** A-177, Wagle Estate, Thane-400 604 (Mumbai) INDIA.

**Country:** India

**Telephone:** (91) (22) 25817000

**Telefax:** (91) (22) 25817002

**E-Mail:** [info@meyer.co.in](mailto:info@meyer.co.in)