

Name of the medicinal product

Ibupkris® Oral Suspension

2. Qualitative and quantitative composition

Ibuprofen USP

The suspension contains 100 mg/5 mL ibuprofenUSP.

Excipients with known effect:

Sucrose	660 mg/mL
Sorbitol (70% non-crystallizing)	100 mg/mL
Sunset yellow (E110)	0.1 mg/mL
Methyl para-hydroxybenzoate (E218)	1 mg/mL
Propyl para-hydroxybenzoate (E216)	0.5 mg/mL

For the full list of excipients, see section 6.1.

3. Pharmaceutical form

An orange-coloured, orange-flavoured, syrupy suspension.

4. Clinical particulars

4.1 Therapeutic indications

Ibupkris® Oral Suspension is indicated for its analgesic and anti-inflammatory effects in the treatment of rheumatoid arthritis (including juvenile rheumatoid arthritis or Still's disease), ankylosing spondylitis, osteoarthritis and other non-rheumatoid (seronegative) arthropathies.

In the treatment of non-articular rheumatic conditions, Ibupkris® Oral Suspension is indicated in peri-articular conditions such as frozen shoulder (capsulitis), bursitis, tendinitis, tenosynovitis and low back pain; Ibupkris® Oral Suspension can also be used in soft-tissue injuries such as sprains and strains.

Ibupkris® Oral Suspension is also indicated for its analgesic effect in the relief of mild to moderate pain such as dysmenorrhoea, dental and post-operative pain and for symptomatic relief of headache including migraine headache.

Ibupkris® Oral Suspension is indicated in short-term use for the treatment of pyrexia in children over one year of age.

4.2 Posology and method of administration

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms (see section 4.4).

Adults and children over 12 years of age: The recommended dosage of Ibupkris® Oral Suspension

is 1200-1800 mg daily in divided doses. Some patients can be maintained on 600-1200 mg daily. Total daily dose should not exceed 2400 mg.

Children: The daily dosage of Ibupkris® Oral Suspension is 100 mg/kg of bodyweight in divided doses. This can be achieved as follows:

1-2 years: One 2.5 ml spoonful (50 mg) three to four times a day.

3-7 years: One 5 ml spoonful (100 mg) three to four times a day.

8-12 years: Two 5 ml spoonful (200 mg) three to four times a day.

Not recommended for children weighing less than 7 kg.

In juvenile rheumatoid arthritis, up to 40 mg/kg of bodyweight daily in divided doses may be taken.

The lowest effective dose should be used for the shortest duration necessary to relieve symptoms (see section 4.4).

Elderly: The elderly is at increased risk of serious consequences of adverse reactions. If an NSAID is considered necessary, the lowest effective dose should be used and for the shortest possible duration. The

patient should be monitored regularly for GI bleeding during NSAID therapy. If renal or hepatic function is impaired, dosage should be assessed individually.

Renal impairment:

Patients with mild to moderate renal impairment, (see section 4.4 - Special warnings and precautions for use) and patients with severe renal insufficiency (see section 4.3 – Contraindications).

Hepatic impairment:

For patients with mild to moderate hepatic impairment (see section 4.4 Special warnings and precautions for use) and patients with severe hepatic dysfunction (see section 4.3-Contraindications).

For oral administration. It is recommended that patients with sensitive stomachs take Ibupkris® Oral Suspension with food. If taken shortly after eating, the onset of action of Ibupkris® Oral Suspension may be delayed. To be taken preferably with or after food.

Ensure the bottle is thoroughly shaken before use. A transient sensation of burning in the mouth or throat may occur with Ibupkris® Oral Suspension.

4.3 Contraindications

Ibupkris® Oral Suspension is contraindicated in patients with hypersensitivity to the active substance or to any of the excipients.

Ibupkris® Oral Suspension should not be used in patients who have previously shown hypersensitivity reactions (e.g., asthma, urticaria, angioedema or rhinitis) after taking ibuprofen, aspirin or other NSAIDs.

Ibupkris® Oral Suspension is also contraindicated in patients with a history of gastrointestinal bleeding or perforation, related to previous NSAID therapy. Ibupkris® Oral Suspension should not be used in patients with active, or history of, recurrent peptic ulcer or gastrointestinal haemorrhage (two or more distinct episodes of proven ulceration or bleeding).

Ibupkris® Oral Suspension should not be given to patients with conditions involving an increased tendency to bleeding.

Ibupkris® Oral Suspension is contraindicated in patients with severe heart failure (NYHA Class IV), hepatic failure and renal failure (see section 4.4).

Ibupkris® Oral Suspension is contraindicated during the last trimester of pregnancy (see section 4.6).

4.4 Special warnings and precautions for use

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms (see section 4.2, and GI and cardiovascular risks below).

Excipients:

- Sucrose - patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltose insufficiency should not take this medicine. Contains 3g sucrose per 5 mL dose, this should be considered in patients with diabetes mellitus and may be harmful to the teeth.
- Sorbitol - this medicine contains 500 mg sorbitol in each 5 mL oral solution. Patients with hereditary fructose intolerance (HFI) should not take this medicinal product. Sorbitol may cause gastrointestinal discomfort and mild laxative effect.
- Methyl para-hydroxybenzoate (E218) and Propyl para-hydroxybenzoate (E216) – may cause allergic reactions (possibly delayed).
- Sunset yellow (E110) - may cause allergic reactions.
- This medicine contains less than 1 mmol sodium (23 mg) per 5ml dose, that is to say essentially 'sodium-free'
- This medicine contains 0.5 mg Sodium Benzoate (E221) in each 5ml dose

As with other NSAIDs, ibuprofen may mask the signs of infection.

The use of Ibupkris® Oral Suspension with concomitant NSAIDs, including cyclooxygenase-2 selective inhibitors, should be avoided due to the increased risk of ulceration or bleeding (see section 4.5).

The diagnosis of medication overuse headache (MOH) should be suspected in patients who have frequent or daily headaches despite (or because of) the regular use of analgesic medication. Patients with medication

overuse headache should not be treated by increasing the dose of the analgesic. In such cases the use of analgesics should be discontinued.

The concomitant consumption of excessive alcohol with NSAIDs, including ibuprofen, may increase the risk of adverse effects on the gastrointestinal tract, such as GI haemorrhage or the central nervous system possibly due to an additive effect.

Elderly

The elderly have an increased frequency of adverse reactions to NSAIDs, especially gastrointestinal bleeding and perforation, which may be fatal (see section 4.2).

Paediatric population

There is a risk of renal impairment in dehydrated children and adolescents.

Gastrointestinal bleeding, ulceration and perforation

GI bleeding, ulceration or perforation, which can be fatal, has been reported with all NSAIDs at anytime during treatment, with or without warning symptoms or a previous history of serious GI events.

The risk of GI bleeding, ulceration or perforation is higher with increasing NSAID doses, in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation (see section 4.3), and in the elderly. These patients should commence treatment on the lowest dose available. Combination therapy with protective agents (e.g. misoprostol or proton pump inhibitors) should be considered for these patients, and also for patients requiring concomitant low dose aspirin, or other drugs likely to increase gastrointestinal risk (see below and section 4.5).

Patients with a history of gastrointestinal disease, particularly when elderly, should report any unusual abdominal symptoms (especially gastrointestinal bleeding) particularly in the initial stages of treatment.

Caution should be advised in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as oral corticosteroids, anticoagulants such as warfarin, selective serotonin-reuptake inhibitors or anti-platelet agents such as aspirin (see section 4.5).

When GI bleeding or ulceration occurs in patients receiving Ibupkris® Oral Suspension, the treatment should be withdrawn. NSAIDs should be given with care to patients with a history of ulcerative colitis or Crohn's disease as these conditions may be exacerbated (see section 4.8).

Respiratory disorders and hypersensitivity reactions

Caution is required if Ibupkris® Oral Suspension is administered to patients suffering from, or with a previous history of, bronchial asthma, chronic rhinitis or allergic diseases since NSAIDs have been reported to precipitate bronchospasm, urticaria or angioedema in such patients.

Cardiac, renal and hepatic impairment

The administration of an NSAID may cause a dose dependent reduction in prostaglandin formation and precipitate renal failure. The habitual concomitant intake of various similar painkillers further increases this risk. Patients at greatest risk of this reaction are those with impaired renal function, cardiac impairment, liver dysfunction, those taking diuretics and the elderly. For these patients, use the lowest effective dose, for the shortest possible duration and monitor renal function especially in long-term treated patients (see also section 4.3).

Brufen should be given with care to patients with a history of heart failure or hypertension since oedema has been reported in association with ibuprofen administration.

Cardiovascular and cerebrovascular effects

Appropriate monitoring and advice are required for patients with a history of hypertension and/or mild to moderate congestive heart failure as fluid retention and oedema have been reported in association with NSAID therapy.

Clinical studies suggest that use of ibuprofen, particularly at a high dose (2400 mg/day) may be associated with a small increased risk of arterial thrombotic events such as myocardial infarction or stroke. Overall, epidemiological studies do not suggest that low dose ibuprofen (e.g. \leq 1200mg/day) is associated with an increased risk of arterial thrombotic events.

Patients with uncontrolled hypertension, congestive heart failure (NYHA II-III), established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease should only be treated with ibuprofen

after careful consideration and high doses (2400mg/day) should be avoided. Careful consideration should also be exercised before initiating long-term treatment of patients with risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking), particularly if high doses of ibuprofen (2400mg/day) are required.

Renal effects

Caution should be used when initiating treatment with ibuprofen in patients with considerable dehydration. There is a risk of renal impairment especially in dehydrated children, adolescents and the elderly.

As with other NSAIDs, long-term administration of ibuprofen has resulted in renal papillary necrosis and other renal pathologic changes. Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of an NSAID may cause a dose-dependant reduction in prostaglandin formation and, secondarily, in renal blood flow, which may cause renal failure. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pre-treatment state.

SLE and mixed connective tissue disease

In patients with systemic lupus erythematosus (SLE) and mixed connective tissue disorders there may be an increased risk of aseptic meningitis (see below and section 4.8).

Severe skin reactions

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs (see section 4.8). Patients appear to be at highest risk of these reactions early in the course of therapy, the onset of the reaction occurring within the first month of treatment in the majority of cases. Acute generalised exanthematous pustulosis (AGEP) has been reported in relation to ibuprofen-containing products. Ibupkris® Oral Suspension should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

In exceptional cases, varicella can be at the origin of serious cutaneous and soft tissues infectious complications. To date, the contributing role of NSAIDs in the worsening of these infections cannot be ruled out. Thus, it is advisable to avoid use of Ibuprofen in case of varicella.

Masking of symptoms of underlying infections

Ibupkris® Oral Suspension can mask symptoms of infection, which may lead to delayed initiation of appropriate treatment and thereby worsening the outcome of the infection. This has been observed in bacterial community acquired pneumonia and bacterial complications to varicella. When Ibupkris® Oral Suspension is administered for fever or pain relief in relation to infection, monitoring of infection is advised. In nonhospital settings, the patient should consult a doctor if symptoms persist or worsen.

Haematological effects

Ibuprofen, like other NSAIDs, can interfere with platelet aggregation and prolong bleeding time in normal subjects.

Aseptic meningitis

Aseptic meningitis has been observed on rare occasions in patients on ibuprofen therapy. Although it is probably more likely to occur in patients with systemic lupus erythematosus and related connective tissue diseases, it has been reported in patients who do not have an underlying chronic disease.

Impaired female fertility

The use of Ibupkris® Oral Suspension may impair female fertility and is not recommended in women attempting to conceive. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of Ibupkris® Oral Suspension should be considered.

Ibupkris® Oral Suspension contains sucrose and sorbitol. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine. Contains 3 g sucrose and 0.5 g sorbitol per 5 mL dose. This should be taken into account in patients with diabetes mellitus. May be harmful to the teeth.

This medicine contains 500 mg sorbitol in each 5 ml oral solution. Patients with hereditary fructose intolerance (HFI) should not take/be given this medicinal product. Sorbitol may cause gastrointestinal discomfort and mild laxative effect.

Ibukris® Oral Suspension contains methyl parahydroxybenzoate and propyl parahydroxybenzoate. May cause allergic reactions (possibly delayed).

Ibukris® Oral Suspension contains sunset yellow (E110). May cause allergic reactions.

This medicine contains less than 1 mmol sodium (23 mg) per dosage unit, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

Care should be taken in patients treated with any of the following drugs as interactions have been reported in some patients. Antihypertensives, beta-blockers and diuretics: NSAIDs may reduce the effect of antihypertensives, such as ACE inhibitors, angiotensin-II receptor antagonists, beta-blockers and diuretics.

Diuretics can also increase the risk of nephrotoxicity of NSAIDs.

Cardiac glycosides: NSAIDs may exacerbate cardiac failure, reduce GFR and increase plasma cardiac glycoside levels.

Cholestyramine; The concomitant administration of ibuprofen and cholestyramine may reduce the absorption of ibuprofen in the gastrointestinal tract. However, the clinical significance is unknown.

Lithium: Decreased elimination of lithium.

Methotrexate: NSAIDs may inhibit the tubular secretion of methotrexate and reduce clearance of methotrexate.

Ciclosporin: Increased risk of nephrotoxicity.

Mifepristone: A decrease in the efficacy of the medicinal product can theoretically occur due to the antiprostaglandin properties of NSAIDs. Limited evidence suggests that coadministration of NSAIDs on the day of prostaglandin administration does not adversely influence the effects of mifepristone or the prostaglandin on cervical ripening or uterine contractility and does not reduce the clinical efficacy of medicinal termination of pregnancy.

Other analgesics and cyclooxygenase-2 selective inhibitors: Avoid concomitant use of two or more NSAIDs, including Cox-2 inhibitors, as this may increase the risk of adverse effects (see section 4.4).

Aspirin (Acetylsalicylic acid): As with other products containing NSAIDs, concomitant administration of ibuprofen and aspirin is not generally recommended because of the potential of increased adverse effects.

Experimental data suggest that ibuprofen may competitively inhibit the effect of low dose aspirin on platelet aggregation when they are dosed concomitantly. Although there are uncertainties regarding extrapolation of these data to the clinical situation, the possibility that regular, long-term use of ibuprofen may reduce the cardioprotective effect of low-dose acetylsalicylic acid cannot be excluded. No clinically relevant effect is considered to be likely for occasional use (see section 5.1).

Corticosteroids: Increased risk of gastrointestinal ulceration or bleeding with NSAIDs (see section 4.4).

Anticoagulants: NSAIDs may enhance the effects of anticoagulants, such as warfarin (see section 4.4).

Quinolone antibiotics: Animal data indicate that NSAIDs can increase the risk of convulsions associated with quinolone antibiotics. Patients taking NSAIDs and quinolones may have an increased risk of developing convulsions.

Sulfonylureas: NSAIDs may potentiate the effects of sulfonylurea medications. There have been rare reports of hypoglycaemia in patients on sulfonylurea medications receiving ibuprofen.

Anti-platelet agents and selective serotonin reuptake inhibitors (SSRIs): Increased risk of gastrointestinal bleeding with NSAIDs (see section 4.4).

Tacrolimus: Possible increased risk of nephrotoxicity when NSAIDs are given with tacrolimus.

Zidovudine: Increased risk of haematological toxicity when NSAIDs are given with zidovudine. There is evidence of an increased risk of haemarthroses and haematoma in HIV(+) haemophiliacs receiving concurrent treatment with zidovudine and ibuprofen.

Aminoglycosides: NSAIDs may decrease the excretion of aminoglycosides.

Herbal extracts: Ginkgo biloba may potentiate the risk of bleeding with NSAIDs.

CYP2C9 Inhibitors: Concomitant administration of ibuprofen with CYP2C9 inhibitors may increase the exposure to ibuprofen (CYP2C9 substrate). In a study with voriconazole and fluconazole (CYP2C9 inhibitors), an increased S(+)-ibuprofen exposure by approximately 80 to 100% has been shown. Reduction of the ibuprofen dose should be considered when potent CYP2C9 inhibitors are administered concomitantly, particularly when high-dose ibuprofen is administered with either voriconazole or fluconazole.

4.6 Pregnancy and lactation

Pregnancy

Inhibition of prostaglandin synthesis may adversely affect the pregnancy and/or embryo/foetal development. Data from epidemiological studies suggest an increased risk of miscarriage and of cardiac malformation and gastroschisis after the use of a prostaglandin synthesis inhibitor in early pregnancy. The risk is believed to increase with dose and duration of therapy. In animals, the administration of a prostaglandin synthesis inhibitor has been shown to result in increased pre- and post-implantation losses and embryo/foetal lethality. In addition, increased incidences of various malformations, including cardiovascular, have been reported in animals given a prostaglandin synthesis inhibitor during the organogenetic period.

During the first and second trimester of pregnancy, Ibupkris® Oral Suspension should not be given unless clearly necessary. If Ibupkris® Oral Suspension is used by a woman attempting to conceive, or during the first or second trimester of pregnancy, the dose should be kept as low and duration of treatment as short as possible.

During the third trimester of pregnancy, all prostaglandin synthesis inhibitors may expose the foetus to the following:

- Cardiopulmonary toxicity (with premature closure of the ductus arteriosus and pulmonary hypertension)
- Renal dysfunction, which may progress to renal failure with oligohydramnios.

At the end of pregnancy, prostaglandin synthesis inhibitors may expose the mother and the neonate to the following:

- Possible prolongation of bleeding time
- Inhibition of uterine contractions, which may result in delayed or prolonged labour.

Consequently, Ibupkris® Oral Suspension is contraindicated during the third trimester of pregnancy.

Lactation

In the limited studies so far available, NSAIDs can appear in the breast milk in very low concentrations. NSAIDs should, if possible, be avoided when breastfeeding.

See section 4.4 Special warnings and precautions for use, regarding female fertility.

4.7 Effects on ability to drive and use machines

Undesirable effects such as dizziness, drowsiness, fatigue and visual disturbances are possible after taking NSAIDs. If affected, patients should not drive or operate machinery.

4.8 Undesirable effects

Gastrointestinal disorders: The most observed adverse events are gastrointestinal in nature. Peptic ulcers, perforation, or GI bleeding, sometimes fatal, particularly in the elderly, may occur (see section 4.4). Nausea, vomiting, diarrhoea, flatulence, constipation, dyspepsia, abdominal pain, melaena, haematemesis, ulcerative stomatitis, gastrointestinal haemorrhage and exacerbation of colitis and Crohn's disease (see section 4.4) have been reported following ibuprofen administration. Less frequently, gastritis, duodenal ulcer, gastric ulcer and gastrointestinal perforation have been observed

A transient sensation of burning in the mouth or throat may occur with Brufen Syrup.

Immune system disorders: Hypersensitivity reactions have been reported following treatment with NSAIDs. These may consist of (a) non-specific allergic reaction and anaphylaxis, (b) respiratory tract reactivity comprising asthma, aggravated asthma, bronchospasm or dyspnoea, or (c) assorted skin disorders, including rashes of various types, pruritus, urticaria, purpura, angioedema and, very rarely, erythema multiforme, bullous dermatoses (including Stevens- Johnson syndrome and toxic epidermal necrolysis).

Cardiac disorders and vascular disorders: Oedema, hypertension and cardiac failure have been reported in association with NSAID treatment. Clinical studies suggest that use of ibuprofen, particularly at high dose (2400 mg/day) may be associated with a small increased risk of arterial thrombotic events such as myocardial infarction or stroke (see section 4.4).

Infections and infestations: Rhinitis and aseptic meningitis (especially in patients with existing autoimmune disorders, such as systemic lupus erythematosus and mixed connective tissue disease) with symptoms of stiff neck, headache, nausea, vomiting, fever or disorientation (see section 4.4).

Exacerbation of infection-related inflammations coinciding with the use of NSAIDs has been described. If signs of an infection occur or get worse during use of Ibuprofen the patient is therefore recommended to go to a doctor without delay.

Skin and subcutaneous tissue disorders: In exceptional cases, severe skin infections and soft-tissue complications may occur during a varicella infection (see also "Infections and infestations")

The following adverse reactions possibly related to ibuprofen and displayed by MedDRA frequency convention and system organ classification. Frequency groupings are classified according to the subsequent conventions: very common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1,000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1,000$), Very rare ($< 1/10,000$) and Not known (cannot be estimated from the available data).

System organ class	Frequency	Adverse reaction
Infections and infestations	Uncommon	Rhinitis
	Rare	Meningitis aseptic (see section 4.4)
Blood and lymphatic system disorders	Rare	Leukopenia, thrombocytopenia, neutropenia, agranulocytosis, aplastic anaemia, haemolytic anaemia
Immune system disorders	Uncommon	Hypersensitivity
	Rare	Anaphylactic reaction
Psychiatric disorders	Uncommon	Insomnia, anxiety
	Rare	Depression, confusional state
Nervous system disorders	Common	Headache, dizziness
	Uncommon	Paraesthesia, somnolence
	Rare	Optic neuritis
Eye disorders	Uncommon	Visual impairment
	Rare	Toxic optic neuropathy
Ear and labyrinth disorders	Uncommon	Hearing impaired, tinnitus, vertigo
Respiratory, thoracic and mediastinal disorders	Uncommon	Asthma, bronchospasm, dyspnoea
Gastrointestinal disorders	Common	Dyspepsia, diarrhoea, nausea, vomiting, abdominal pain, flatulence, constipation, melaena, haematemesis, gastrointestinal haemorrhage
	Uncommon	Gastritis, duodenal ulcer, gastric ulcer, mouth ulceration, gastrointestinal perforation
	Very rare	Pancreatitis
	Not known	Exacerbation of Colitis and Crohn´s disease
Hepatobiliary disorders	Uncommon	Hepatitis, jaundice, hepatic function abnormal
	Very Rare	Hepatic failure
Skin and subcutaneous tissue disorders	Common	Rash
	Uncommon	Urticaria, pruritus, purpura, angioedema, photosensitivity reaction
	Very rare	Severe forms of skin reactions (e.g. Erythema multiforme, bullous reactions, including Stevens-Johnson syndrome, and toxic epidermal necrolysis)
	Not known	Drug reaction with eosinophilia and systemic symptoms (DRESS syndrome) Acute generalised exanthematous pustulosis (AGEP)

Renal and urinary disorders	Uncommon	Nephrotoxicity in various forms e.g. Tubulointerstitial nephritis, nephrotic syndrome and renal failure
General disorders and administration site conditions	Common	Fatigue
	Rare	Oedema
Cardiac disorders	Very rare	Cardiac failure, myocardial infarction (also see section 4.4)
Vascular disorders	Very rare	Hypertension

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions.

4.9 Overdose

Toxicity

Signs and symptoms of toxicity have generally not been observed at doses below 100 mg/kg in children or adults. However, supportive care may be needed in some cases. Children have been observed to manifest signs and symptoms of toxicity after ingestion of 400 mg/kg or greater.

Symptoms

Most patients who have ingested significant amounts of ibuprofen will manifest symptoms within 4 to 6 hours.

The most frequently reported symptoms of overdose include nausea, vomiting, abdominal pain, lethargy and drowsiness. Central nervous system (CNS) effects include headache, tinnitus, dizziness, convulsion, and loss of consciousness. Nystagmus, metabolic acidosis, hypothermia, renal effects, gastrointestinal bleeding, coma, apnoea, diarrhoea and depression of the CNS and respiratory system have also been rarely reported. In serious poisoning metabolic acidosis may occur. Disorientation, excitation, fainting and cardiovascular toxicity, including hypotension, bradycardia and tachycardia have been reported. In cases of significant overdose, renal failure and liver damage are possible. Large overdoses are generally well tolerated when no other drugs are being taken.

Therapeutic measures

Patients should be treated symptomatically as required. Within one hour of ingestion of a potentially toxic amount, activated charcoal should be considered. Alternatively, in adults, gastric lavage should be considered within one hour of ingestion of a potentially life-threatening overdose.

Good urine output should be ensured.

Renal and liver function should be closely monitored.

Patients should be observed for at least four hours after ingestion of potentially toxic amounts.

Frequent or prolonged convulsions should be treated with intravenous diazepam. Other measures may be indicated by the patient's clinical condition.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic classification: Anti-inflammatory and antirheumatic products, nonsteroidal; propionic acid derivatives.

ATC code: M01AE01

Ibuprofen is a propionic acid derivative with analgesic, anti-inflammatory and anti-pyretic activity. The drug's therapeutic effects as an NSAID is thought to result from its inhibitory effect on the enzyme cyclo-oxygenase, which results in a marked reduction in prostaglandin synthesis.

Experimental data suggest that ibuprofen may competitively inhibit the effect of low dose aspirin on platelet aggregation when they are dosed concomitantly. Some pharmacodynamic studies show that when single doses of ibuprofen 400mg were taken within 8 hours before or within 30 minutes after immediate release aspirin dosing (81mg), a decreased effect of aspirin on the formation of thromboxane or platelet aggregation occurred. Although there are uncertainties regarding extrapolation of these data to the clinical situation, the possibility that regular, long-term use of ibuprofen may reduce the cardioprotective effect of low-dose

acetylsalicylic acid cannot be excluded. No clinically relevant effect is likely for occasional ibuprofen use. (see section 4.5)

5.2 Pharmacokinetic properties

Ibuprofen is rapidly absorbed from the gastrointestinal tract, peak serum concentrations occurring 1-2 hours after administration. The elimination half-life is approximately 2 hours.

Ibuprofen is metabolised in the liver to two inactive metabolites and these, together with unchanged ibuprofen, are excreted by the kidney either as such or as conjugates. Excretion by the kidney is both rapid and complete.

Ibuprofen is extensively bound to plasma proteins.

5.3 Preclinical safety data

Not applicable.

6. Pharmaceutical particulars

6.1 List of excipients

Methyl hydroxybenzoate (E218)

Propyl hydroxybenzoate (E216)

Refined sugar

Citric acid monohydrate granular

Sodium benzoate (E221)

Agar powder

Glycerine

Sorbitol solution 70% (non-crystallising)

Polysorbate 80

Sunset yellow (E110)

Orange flavour D717

Purified water

6.2 Incompatibilities

None known.

6.3 Shelf life

36 months.

Once the bottle is opened, the Ibupkris® Oral Suspension should be used within 12 months.

6.4 Special precautions for storage

Store below 30°C and protect from light and moisture.

6.5 Nature and contents of container

An amber-coloured polyethylene terephthalate (PET) bottle with an aluminium cap fitted with a low-density polyethylene liner.

Pack sizes of 100ml.

6.6 Special precautions for disposal and other handling

None stated.

Administrative Data

7. Manufactured by

Krishat Pharma Industries Limited

KM 15, Lagos-Ibadan Expressway, Ibadan, Oyo State,

NIGERIA.

Email: info@krishatpharma.com

Company contacts details

operations@krishatpharma.com