

## **1. NAME OF THE MEDICINAL PRODUCT**

**TEMPOVATE GEL**

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Clobetasol propionate 0.05% w/w

Excipients with known effect:

Propylene glycol

For the full list of excipients, see section 6.1

## **3. PHARMACEUTICAL FORM**

Gel

Light Pink Gel

## **4. Clinical particulars**

### **4.1 Therapeutic indications**

Short – courses treatment of more resistant dermatoses e.g psoriasis, recalcitrant eczemas, lichen planus, discoid lupus erythematosus, and other conditions which do not respond satisfactorily to less active steroids.

### **4.2 Posology and method of administration**

#### Posology

Adult and children

Apply sparingly to the affected area once or twice daily until improvement occurs. As with other highly-active topical steroid preparations, therapy should be discontinued when control is achieved. In the more responsive conditions this may be within a few days. If a longer course is necessary, it is recommended that the drug should not be continued for more than four weeks without the patient's conditions being reviewed.

Repeated short courses of Tempovate Gel steroid therapy may be used to control exacerbations. If continuous steroid treatment is necessary, a less potent corticosteroid preparations should be used.

In very resistant lesions, especially where there is hyperkeratosis, the anti-inflammatory effect of Tempovate Gel can be enhanced, if necessary, by occluding the treatment area with polythene film. Overnight occlusion alone is usually adequate to bring about a satisfactory response. Thereafter, improvement can be further maintained by application without occlusion.

Rarely, occlusion is necessary. In cases where an occlusive dressing is applied, caution is

needed in order to avoid the risk of local and systemic adverse events.

Tempovate Gel should only be used for no more than 5 days on the face and eyelids.

#### Method of administration

For topical administration.

Application of gel in adults:

- Two fingertips of gel will cover both hands or one foot
- Three fingertips of gel will cover one arm
- Six fingertips of gel will cover one leg
- Fourteen fingertips of gel will cover the front and back of the body.
- If no improvement is seen within two to four weeks, reassessment of the diagnosis, or referral, may be necessary.
- Application of gel in children 1 year and older:
- The smaller the child the less you will need to use.
- A child of 4 years needs about a third of the adult amount.
- A course of treatment for a child should not normally last more than 5 days - unless your doctor has told you to use it for longer. The doctor may want to see the child every week, whilst using the gel.

Children below 1 year

Tempovate Gel is contraindicated in children below 1 year.

Children are more susceptible to local and systemic adverse events caused by topical corticosteroids, and should generally be treated for a shorter duration and with less potent substances than adults. Clobetasol should be used with caution in children, to ensure that the smallest possible dose is applied within the therapeutic range.

### **4.3 Contraindications**

- Rosacea
- Acne vulgaris
- Perioral dermatitis
- Perianal and genital pruritus
- Primary cutaneous viral infections (e.g. herpes simplex, chickenpox)
- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- The use of Tempovate Gel not indicated in the treatment of primary infected skin lesions caused by infection with fungi (e.g. candidiasis, tinea) or bacteria (e.g. impetigo)
- Dermatoses in children under one year of age, including dermatitis and napkin eruptions

#### **4.4 Special warnings and precautions for use**

Long term continuous therapy with Tempovate Gel should be avoided, particularly in infants and children, in whom adrenal suppression occurs readily. If Tempovate Gel is required for use in children, it is recommended that the treatment should be reviewed weekly. It should be noted that the infant's napkin may act as an occlusive dressing.

If used in children or on the face, courses should be limited if possible to five days and occlusion should not be used.

The face, more than other areas of the body, may exhibit atrophic changes after prolonged treatment with potent topical corticosteroids. This must be borne in mind when treating such conditions as psoriasis, discoid lupus erythematosus and severe eczema.

If applied to the eyelids, care is needed to ensure that the preparation does not enter the eye, as glaucoma might result. If Tempovate Gel does enter the eye, the affected eye should be bathed in copious amounts of water.

##### Visual disturbance

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids

Topical steroids may be hazardous in psoriasis for a number of reasons including rebound relapses, development of tolerance, risk of generalised pustular psoriasis and development of local or systemic toxicity due to impaired barrier function of the skin. If used in psoriasis careful patient supervision is important.

Appropriate antimicrobial therapy should be used whenever the treated inflammatory lesions which have become infected. Any spread of infection requires withdrawal of topical corticosteroid therapy and systemic administration of antimicrobial agents. Bacterial infection is encouraged by the warm, moist conditions induced by occlusive dressings, and so the skin should be cleansed before a fresh dressing is applied.

During application of corticosteroids on large areas, especially under (plastic) occlusion or in skin folds, increased absorption may occur, which could lead to adrenal function inhibition.

There have been a few reports in the literature of the development of cataracts in patients who have been using corticosteroids for prolonged periods of time. Although it is not possible to rule out systemic corticosteroids as a known factor, prescribers should be aware of the possible role of corticosteroids in cataract development.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

No interaction studies have been performed.

#### **4.6 Pregnancy and Lactation**

##### Pregnancy

There is inadequate evidence of safety in human pregnancy. Topical administration of corticosteroids to pregnant animals can cause abnormalities of fetal development including cleft palate and intrauterine growth retardation. The relevance of this finding to humans has not been established, therefore, topical steroids should not be used extensively in pregnancy, i.e. in large

amounts or for prolonged periods.

### Breast-feeding

The safe use of topical corticosteroids during lactation has not been established.

It is not known whether the topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable amounts in breast milk.

Administration of clobetasol during lactation should only be considered if the expected benefit to the mother outweighs the risk to the infant.

If used during lactation clobetasol should not be applied to the breasts to avoid accidental ingestion by the infant.

### Fertility

There are insufficient fertility data available to indicate whether clobetasol propionate has any effect on fertility.

## **4.7 Effects on ability to drive and use machines**

Tempovate Gel is not expected to have any effects.

## **4.8 Undesirable effects**

The following adverse reactions have been identified during post-approval use of clobetasol propionate. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. The frequency of these adverse events has therefore been classified as “unknown”.

### **Immune system disorders**

#### Hypersensitivity

- Local hypersensitivity reactions such as erythema, rash, pruritus, urticaria and allergic contact dermatitis may occur at the site of application and may resemble symptoms of the condition under treatment.
- If signs of hypersensitivity appear, application should be stopped immediately.

### **Endocrine disorders**

#### Features of Cushing's syndrome

- As with other topical corticosteroids, prolonged use of large amounts, or treatment of extensive areas can result in sufficient systemic absorption to produce the features of Cushing's syndrome. This effect is more likely to occur in infants and children, and if occlusive dressings are used. In infants, the nappy may act as an occlusive dressing.
- Provided the weekly dosage is less than 50 g in adults, any suppression of the HPA axis is likely to be transient with a rapid return to normal values once the short course of steroid therapy has ceased. The same applies to children given proportionate dosage.

### **Eye disorders**

- Vision, blurred (see also section 4.4).

### **Vascular disorders**

Dilatation of the superficial blood vessels

- Prolonged and intensive treatment with highly-active corticosteroid preparations may cause dilatation of the superficial blood vessels, particularly when occlusive dressings are used, or when skin folds are involved.

### **Skin and subcutaneous tissue disorders**

Local skin burning, local atrophy, striae, thinning, pigmentation changes, hypertrichosis, exacerbation of underlying symptoms, pustular psoriasis.

- Prolonged and intensive treatment with highly-active corticosteroid preparations may cause local atrophic changes, such as thinning and striae.
- Treatment of psoriasis with corticosteroids (or its withdrawal) is thought to have provoked the pustular form of the disease.
- Clobetasol may induce steroid-rosacea and steroid-acne.

## **4.9 Overdose**

Acute overdosage is very unlikely to occur, however, in the case of chronic overdosage or misuse, the features of hypercortisolism may appear and in this situation topical steroids should be reduced or discontinued gradually, under medical supervision.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamics properties**

Pharmacotherapeutic group: Corticosteroids, very potent, dermatological preparations (group IV)

ATC code: D07AD01

Clobetasol propionate is a highly active corticosteroid with topical anti-inflammatory activity. The major effect of clobetasol propionate on skin is a non-specific anti-inflammatory response, partially due to vasoconstriction and decrease in collagen synthesis.

### **5.2 Pharmacokinetic properties**

Percutaneous penetration of clobetasol propionate varies among individuals and can be increased by the use of occlusive dressings, or when the skin is inflamed or diseased.

Mean peak plasma clobetasol propionate concentrations of 0.63 ng/ml occurred in one study eight hours after the second application (13 hours after an initial application) of 30 g clobetasol propionate 500 micrograms/g ointment to normal individuals with healthy skin. Following the application of a second dose of 30 g clobetasol propionate gel 500 micrograms/g mean peak

plasma concentrations were slightly higher than the ointment and occurred 10 hours after application.

In a separate study, mean peak plasma concentrations of approximately 2.3 ng/ml and 4.6 ng/ml occurred respectively in patients with psoriasis and eczema three hours after a single application of 25 g clobetasol propionate 500 micrograms/g ointment.

Following percutaneous absorption of clobetasol propionate, the drug probably follows the metabolic pathway of systemically administered corticosteroids, i.e. metabolised primarily by the liver and then excreted by the kidneys. However, systemic metabolism of clobetasol has never been fully characterized or quantified.

### **5.3 Preclinical safety data**

Parenteral administration of corticosteroids, including clobetasol propionate, to pregnant animals can cause abnormalities of fetal development including cleft palate and intrauterine growth retardation. Animal studies have indicated that intrauterine exposure to corticosteroids may contribute to the development of cardiovascular and metabolic diseases in adult life, but there is a lack of evidence for the occurrence of such effects in humans.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Carbopol 990  
DPD-3111, Floral Aldehyde BQT  
Methyl Hydroxybenzoate  
Oxyde De Titane Organophile W877  
Propylene Glycol  
Propyl Hydroxybenzoate  
Sicofarm Erythrosine 85 E127, CI 45430  
Triethanolamine  
Purified water

### **6.2 Incompatibilities**

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

Store below 30°C. Keep All medicines out of the reach of children. Use only as directed by the physician for external use only.

### **6.5 Nature and contents of container**

Tube with inner lacquer and rubber end seal.  
Pack size: 30 g

## **6.6 Special precautions for disposal and other handling**

Patients should be advised to wash their hands after applying Tempovate Gel unless it is the hands that are being treated.

## **7. APPLICANT/MANUFACTURER**

### **Applicant**

#### **Tempo Scan Africa Ltd.**

Sky Water Tower, No 6 Otunba Yomi Oshikoya Drive, Off Mobolaji Bank Anthony Way Ikeja, Lagos State, Nigeria.

### **Manufacturer**

PT. Supra Ferbindo Farma

EJIP Industrial Park Plot 8J, Cikarang, Bekasi, 17550 - INDONESIA

Released by

PT. Tempo Scan Pacific

EJIP Industrial Park, Plot 1 G-H, Cikarang, Bekasi 17550 - INDONESIA