

1. Name of the medicinal product

Emzor Griseofulvin 500mg Tablets

2. Qualitative and quantitative composition

Each film-coated tablet contains 500mg Emzor Griseofulvin

For a full list of excipients, see section 6.1.

3. Pharmaceutical form

Film-coated tablet.

White to off white, round, biconvex film-coated tablet.

4. Clinical particulars

4.1 Therapeutic indications

The treatment of fungal infections of the skin, scalp, hair, or nails (Tinea barbae, Tinea capitis, Tinea corporis, Tinea cruris, Tinea pedis, Tinea unguium) where topical therapy is considered inappropriate, or the infection has proven refractory to topical therapy.

Oral administration of Emzor Griseofulvin for systemic therapy of fungal infections enables newly formed keratin of the skin, hair, and nails to resist fungal attack. As the new keratin extends, the old infected keratin is shed.

Prior to therapy, the type of fungi responsible should be identified. The use of Emzor Griseofulvin is not justified in the treatment of minor or trivial infections that will respond to topical therapy.

Before prescribing Emzor Griseofulvin Tablets, consideration should be given to national and/or local guidance on the appropriate use of antifungals.

4.2 Posology and method of administration

General:

For oral administration.

Tablets should be swallowed whole with a glass of water. Emzor Griseofulvin is recommended to be taken after a high fat meal, for increased absorption and minimising GI distress, see section 5.2.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Concomitant use of appropriate topical agents is usually required, particularly in treatment of tinea pedis. In some forms of tinea pedis, yeasts and bacteria may be involved as well as fungi. Emzor Griseofulvin will not eradicate the bacterial or candidal infections.

Adults

The usual adult dose is 500 mg to 1000 mg daily. The dose should not be less than 10 mg / Kg bodyweight / day. The dose may be administered as a single daily dose, or it may be administered twice daily. The twice daily dosing regimen may be more effective in those patients who respond poorly.

Hepatic impairment

Emzor Griseofulvin is contraindicated in patients with severe hepatic impairment, see section 4.3.

For patients with moderate to mild hepatic impairment, no dosage adjustment is required. However Emzor Griseofulvin may lead to further impairment of hepatic function, therefore regular monitoring of liver function is mandated, see section 4.4.

Renal impairment

No dosage adjustment is required in renally impaired patients; renal insufficiency does not lead to accumulation.

Elderly

No dosage adjustment is required in the elderly. Consideration should be given that such patients may also have a degree of hepatic impairment, see section 4.4.

Children

The dosage form, film-coated tablet, is only suitable for children of an age to swallow the tablet.

The usual dose is 10 mg / Kg bodyweight / day, in divided doses.

Duration of therapy

The duration of therapy depends upon the thickness of keratin at the site of infection, and the clinical response. The following duration of therapy are indicative:

Tinea corporis: 2-4 weeks

Tinea capitis: 4-8 weeks, in refractory cases, 8-12 weeks

Tinea pedis: 4-8 weeks

Tinea unguium: 6-12 months

Therapy should be continued for at least two weeks after all signs of infection have disappeared.

4.3 Contraindications

Emzor Griseofulvin is contraindicated in patients who have:

- Hypersensitivity to Emzor Griseofulvin or to any of the excipients, see section 6.1
- Porphyria
- Severe hepatic impairment
- Systemic Lupus Erythematosus (SLE)
- Pregnancy, see section 4.6
- Breastfeeding, see section 4.6

4.4 Special warnings and precautions for use

Emzor Griseofulvin is recommended after a high fat meal for increased absorption and minimising GI distress.

Emzor Griseofulvin is contraindicated in patients with severe hepatic impairment, see section 4.3. In patients with minor to moderate hepatic impairment, Emzor Griseofulvin may cause further deterioration of hepatic function. Therefore care should be exercised with such patients, and it is recommended to perform regular periodic liver function tests, see section 4.8.

Emzor Griseofulvin is contraindicated in patients with Systemic Lupus Erythematosus (SLE), see section 4.3; Emzor Griseofulvin has been reported to exacerbate the conditions, and care should be taken to exclude patients with pre-existing SLE from therapy.

Animal data, see section 5.3, indicates long term administration of high dose Emzor Griseofulvin induces tumours in some species, but not others. The clinical relevance of this to man is unknown, but Emzor Griseofulvin should not be used prophylactically.

Emzor Griseofulvin is a liver microsomal enzyme inducer and thus may impair the effectiveness of oral contraceptives. Therefore in women of child bearing age using oral contraception, additional barrier methods of contraception must be used during therapy and for 4 weeks following therapy cessation, see sections 4.5 and 4.6.

Emzor Griseofulvin causes chromosomal abnormalities in animals, see section 5.3. Therefore sexually active males should be cautioned to use an effective barrier method of contraception throughout therapy and for 6 months after therapy termination, see section 4.6.

A theoretical possibility of cross sensitivity in patients known to be allergic to penicillins exists, therefore caution should be exercised in administration of Emzor Griseofulvin to such patients. It should be noted that such patients have been satisfactorily treated with Emzor Griseofulvin without sequelae.

Patients should be cautioned to avoid excessive and unnecessary exposure to sunlight or U.V sources, including sunbeds, during Emzor Griseofulvin therapy as photosensitivity reactions can occur, see section 4.8.

Consumption of alcohol in association with Emzor Griseofulvin can result in an "Antabuse" type reaction, see section 4.5. Patients should be cautioned to avoid consumption of alcoholic beverages, and medicines containing alcohol, while undergoing Emzor Griseofulvin therapy.

In patients undergoing long term Emzor Griseofulvin therapy, i.e for tinea unguium, consideration should be given to periodic monitoring of blood chemistry, particularly for patients with pre-existing blood disorders, since Emzor Griseofulvin may cause blood disorders, see section 4.8.

In common with any antibiotic, therapy with Emzor Griseofulvin may result in the overgrowth of non-susceptible organisms, i.e bacteria or yeasts, or non-dermatophyte fungi, that are often cofactors in tinea infections, especially tinea pedis. Additional therapy is required to control or eradicate such organisms, as Emzor Griseofulvin is ineffective.

Emzor Griseofulvin is not effective in infections due to *Candida albicans*, *Aspergillus sp.*, *Malassezia furfur* (*Pityriasis versicolor*) and *Nocardia sp.* It has no antibacterial effects.

4.5 Interaction with other medicinal products and other forms of interaction

Medicinal Products:

Emzor Griseofulvin may depress plasma levels, and therefore the efficacy, of concomitantly administered medicinal products that are metabolised by cytochrome P450 3A4.

Interactions of Emzor Griseofulvin with other drugs:

Ciclosporin: concomitant administration may result in a reduction of ciclosporin plasma levels, necessitating a dosage adjustment. Plasma levels of ciclosporin should be monitored during griseofulvin therapy, and necessary dosage adjustments made.

Coumarin anticoagulants: the efficacy may be reduced, necessitating dosage adjustment. It is recommended that both prothrombin and INR are regularly monitored, for the duration of Emzor Griseofulvin therapy, and for 8 days post therapy cessation.

Methadone: depression of methadone plasma levels may occur during Emzor Griseofulvin therapy. Patients should be closely monitored for any loss of efficacy, or plasma levels of methadone be monitored, and corresponding dosage adjustments made.

Oral contraceptives: efficacy of oral contraception is reduced during Emzor Griseofulvin therapy and for four weeks post therapy cessation. In view of the contraindication in pregnancy, see section 4.3, and of the possible sequelae of male patients fathering a child during therapy, all sexually active patients should use additional barrier contraception, such as condoms, throughout Emzor Griseofulvin therapy, and for four weeks (female) and 6 months (male) post therapy cessation. See also sections 4.3, 4.4, 4.6, and 5.3 for additional information.

Interactions of other drugs with Emzor Griseofulvin:

Concurrent administration of other medicinal products that induce metabolising enzymes may result in a reduction of Emzor Griseofulvin blood plasma levels and thus efficacy. The following drugs are known to have this effect:

Barbiturates, such as phenobarbitone

Doxercalciferol

Phenylbutazone

Primidone

Other sedative and hypnotic drugs that induce metabolising enzymes.

Food: administration of Emzor Griseofulvin after food, results in increased absorption, and thus higher plasma levels. This effect is enhanced if the meal contains high fat content. Administration after food is recommended, see section 4.2.

Alcohol: there are reports that Emzor Griseofulvin enhances the central nervous system effects of alcohol. There are also reports that Emzor Griseofulvin and alcohol use result in an "Antabuse" type reaction. Patients should be cautioned to avoid alcohol and all alcohol containing products while undergoing Emzor Griseofulvin therapy, See also section 4.8.

4.6 Pregnancy and lactation

Pregnancy:

There are case reports of human foetal abnormalities associated with Emzor Griseofulvin.

There are no adequate and well controlled studies in man, and inadequate epidemiological data. Emzor Griseofulvin has been shown to be teratogenic and embryotoxic in mice and rats. (**see section 5.3**).

Emzor Griseofulvin is suspected to cause serious birth defects when administered during pregnancy.

Emzor Griseofulvin is contraindicated (see section 4.3) in pregnancy.

Women of childbearing potential have to use effective contraception during (and up to 4 weeks after) treatment (see section 4.5) in respect of effect on oral contraceptives, and contraceptive precautions.

Male-mediated effects on pregnancy

Emzor Griseofulvin has been shown to induce chromosomal aberrations in animal spermatocytes (see section 5.3). Therefore men should take effective contraceptive precautions, i.e barrier contraception, to avoid fathering children for the duration of Emzor Griseofulvin therapy, and for 6 months post therapy cessation.

Lactation:

It is unknown if Emzor Griseofulvin is excreted in breast milk, but the possibility does exist. There is inadequate data on the safety of Emzor Griseofulvin in breast feeding, and the potential risk to the infant cannot be assessed, therefore Emzor Griseofulvin is contraindicated in breast feeding (see section 4.3).

4.7 Effects on ability to drive and use machines

Emzor Griseofulvin has no or negligible influence on the ability to drive and use machines. However, it may cause drowsiness, confusion dizziness, and impaired co-ordination, see section 4.8. Patients should therefore be cautioned not to drive or operate machines until they are sure they are not affected.

4.8 Undesirable effects

The following frequencies are used for the description of the occurrence of undesirable effects:

Very common: $\geq 1 / 10$

Common: $\geq 1 / 100, < 1 / 10$

Uncommon: $\geq 1 / 1,000, < 1 / 100$

Rare: $\geq 1 / 10,000, < 1 / 1,000$

Very rare: $< 1 / 10,000$

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Headache and gastric discomfort are the most common effects on starting treatment, but usually disappear as treatment is continued.

Blood and lymphatic system disorder:

Rare: leucopenia, neutropenia, anaemia-these usually resolve on therapy cessation

Nervous system disorders:

Common: headache

Uncommon: impaired co-ordination, peripheral neuropathy, confusion, dizziness, drowsiness, insomnia, irritability.

Gastrointestinal disorders:

Common: diarrhoea, vomiting, nausea, gastric discomfort

Uncommon: anorexia, taste sensation changes

Skin and subcutaneous tissue disorders:

Uncommon: toxic epidermal necrolysis, erythema multiforme, photosensitivity on exposure to intense natural or artificial sunlight.

Rare: precipitation of Systemic Lupus Erythematosus, bullous reactions including Lyell's syndrome, urticarial reactions, skin rashes.

Hepatobiliary disorders:

Very rare: alteration in liver function tests, with elevation to more than three times upper normal limit, intrahepatic cholestasis, hepatitis.

4.9 Overdose

No case of overdose has been reported.

Symptoms:

The likely symptoms of any overdose would be nausea, vomiting, headache, numbness and tingling, confusion, and vertigo. Urticaria or porphyria could occur.

Treatment:

There is no specific antidote to Emzor Griseofulvin. Gastric lavage, or the induction of emesis may be of help, if ingestion is recent. Administration of activated charcoal may also be of use. Treatment should be symptomatic and supportive. Laboratory monitoring of haemopoietic, hepatic and nephritic parameters and electrolytes is recommended.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antifungals for systemic use

ATC code: D01BA01

Emzor Griseofulvin is an antifungal antibiotic that is active *in vivo* against common dermatophytes. The antifungal effect is manifested by binding to tubulin, at distinct binding sites, thus interfering with the microtubule function and causing inhibition of mitosis, and arresting cell division.

The inhibition of fungal mitosis leads to the production of multinucleate cells of characteristic morphology.

On entering the systemic circulation, Emzor Griseofulvin binds to keratin in keratin precursor cells, thereby making them resistant to fungal infections. The drug only reaches the site of action when hair or skin is replaced by the keratin-Emzor Griseofulvin complex.

Emzor Griseofulvin then enters the dermatophyte through energy dependent transport processes and binds to the fungal microtubules, interfering with, and inhibiting mitosis, and the deposition of fungal cell walls.

Mycology:

Emzor Griseofulvin has antifungal activity against the following dermatophytes, although there is species and strain variability in susceptibility.

Trichophyton rubrum, *T. tonsurans*, *T. mentagrophytes*, *T. interdigitalis*, *T. verrucosum*, *T. megnini*, *T. gallinae*, *T. Crateriform*, *T. sulphureum* and *T. schoenleinii*.

Microsporum audouinii, *M. Canis*, *M. gypseum*.

Epidermophyton floccosum.

Emzor Griseofulvin has no activity against dermatophyte fungi of other genera, non-dermatophyte fungi, yeasts, gram positive bacteria, or gram negative bacteria. If any of these are cofactors in the pathology of infection, suitable additional therapy will be required for their eradication.

5.2 Pharmacokinetic properties

Absorption:

The absorption of Emzor Griseofulvin from the gastrointestinal tract is variable and incomplete. On average, less than 50% of the oral dose is absorbed, but administration after a fatty meal, and a reduction in particle size will increase the rate and extent of the absorption.

Following oral administration there is a phase of rapid absorption, and thereafter a phase of slower prolonged absorption.

Peak plasma levels, 0.5 µg / ml-1.5 µg / ml after a 500 mg dose, and 1.5 µg / ml-3.0 µg / ml after a 1000 mg dose, are reached in 2-4 hours, and are maintained for some 10-20 hours.

Emzor Griseofulvin exhibits linear pharmacokinetics.

Distribution:

The volume of distribution is about 0.7 L / Kg, and Emzor Griseofulvin is ca 80 % bound to plasma proteins, predominantly serum albumin.

Emzor Griseofulvin crosses the placenta, and may be excreted in breast milk. There is selective deposition of Emzor Griseofulvin in newly formed keratin of hair, skin, and nails, which gradually moves to the surface of these appendages.

Metabolism:

Emzor Griseofulvin undergoes metabolism to inactive metabolites, principally 6- desmethylEmzor Griseofulvin, or its glucuronide conjugate.

Excretion:

The terminal plasma half life ranges from 9.5-21 hours, with considerable intersubject variability. The majority of the dose, as 6-desmethylEmzor Griseofulvin or the glucuronide conjugate, and other metabolites is excreted in the urine, with less than 1% administered dose being excreted as unchanged Emzor Griseofulvin. The remainder of the dose, principally as metabolites, is excreted in bile and faeces.

Renal insufficiency does not lead to accumulation.

5.3 Preclinical safety data

Emzor Griseofulvin can induce aneuploidy and meiotic delay in mouse oocytes following oral administration of high doses, i.e. 250mg/kg or greater. In addition, Emzor Griseofulvin caused increases in numerical and structural chromosome aberrations in mouse spermatocytes at doses of 500mg/kg and above. Aneuploidy was observed at doses of 1500mg/kg.

Emzor Griseofulvin administered to rats and mice during pregnancy has been associated with foetotoxicity and foetal malformations. Long-term administration of high doses of Emzor Griseofulvin with food has been reported to induce hepatomas in mice and thyroid tumours in rats but not hamsters (see contraindications). The effects in mice may be due to a species specific effect on porphyrin metabolism.

6. Pharmaceutical particulars

6.1 List of excipients

Tablet core:	Maize starch
	Microcrystalline cellulose E 460
	Sodium laurilsulphate
	Povidone E 1201
	Magnesium stearate E 470b
Film coat:	Hypromellose E 464
	Ethylcellulose E 462

Polysorbate 80 E 433
Propylene glycol E 1520

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

Unopened container: 36 Months

Opened container: 6 Months

6.4 Special precautions for storage

Do not store above 25°C.

6.5 Nature and contents of container

Polypropylene (PP) tablet container with linear low density polyethylene (LLDPE) screw closure.

Tablet containers with 90 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Any unused product or waste material should be disposed of in accordance with local requirements.

7. Marketing authorisation holder

Emzor Pharmaceutical Industries Limited

Km 1 Flowergate Mixed Development Scheme, Sagamu/Benin Expressway, Makun, Sagamu, Ogun-State

8. Marketing authorisation number(s)

NA

9. Date of first authorisation/renewal of the authorisation

NA

10. Date of revision of the text

NA