



MECURE INDUSTRIES PLC

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

SUMMARY OF PRODUCT CHARACTERISTICS

1. Name of the medicinal product

Tramez 100 mg Capsules (Tramadol HCL 100mg)

2. Qualitative and quantitative composition

Each capsule contains 100 mg of Tramadol hydrochloride BP

For the full list of excipients, see section 6.1.

3. Pharmaceutical form

Capsule

Green colour cap and yellow colour body gelatin capsules with white powder, printed “tramez” on cap and body

4. Clinical particulars

4.1 Therapeutic indications

Treatment of moderate to severe pain

4.2 Posology and method of administration

Posology:

The dose should be adjusted to the intensity of the pain and the sensitivity of the individual patient. The lowest effective dose for analgesia should generally be selected. The total daily dose of 400 mg active substance should not be exceeded, except in special circumstances.

Unless otherwise prescribed, Tramadol should be administered as follows:

Adults and adolescents above the age of 12 years:

The usual initial dose is 50-100 mg tramadol hydrochloride twice daily, morning and evening. If pain relief is insufficient, the dose may be titrated upwards to 150 mg or 200 mg tramadol hydrochloride twice daily (see section 5.1).

Children

Tramez is not suitable for children below the age of 12 years.

Geriatric patients

A dose adjustment is not usually necessary in patients up to 75 years without clinically manifest hepatic or renal insufficiency. In elderly patients over 75 years elimination may be prolonged. Therefore, if necessary the dosage interval is to be extended according to the patient's requirements.

Renal insufficiency/dialysis and hepatic insufficiency

In patients with renal and/or hepatic insufficiency the elimination of tramadol is delayed. In these patients prolongation of the dosage interval should be carefully considered according to the patients requirements.

Method of administration

The tablets are to be taken whole, not divided or chewed, with sufficient liquid, independent of meals.

Duration of administration

Tramadol should under no circumstances be administered for longer than absolutely necessary. If long-term pain treatment with tramadol is necessary in view of the nature and severity of the illness, then careful and regular monitoring should be carried out (if necessary with breaks in treatment) to establish whether and to what extent further treatment is necessary.

4.3 Contraindications

Tramadol is contraindicated

- in hypersensitivity to the active substance or any of the excipients listed in section 6.1,
- in acute intoxication with alcohol, hypnotics, analgesics, opioids, or psychotropic medicinal products,
- in patients who are receiving MAO inhibitors or who have taken them within the last 14 days
- in patients with epilepsy not adequately controlled by treatment,
- for use in narcotic withdrawal treatment.

4.4 Special warnings and precautions for use

Tramadol may only be used with particular caution in opioid-dependent patients, patients with head injury, shock, a reduced level of consciousness of uncertain origin, disorders of the respiratory centre or function, increased intracranial pressure.

In patients sensitive to opiates tramadol should only be used with caution.

Concomitant use of tramadol and sedating medicinal products such as benzodiazepines or related substances, may result in sedation, respiratory depression, coma and death. Because of these risks, concomitant prescribing with these sedating medicinal products should be reserved for patients for whom alternative treatment options are not possible. If a decision is made to prescribe tramadol concomitantly with sedating medicinal products, the lowest effective dose of tramadol should be used, and the duration of the concomitant treatment should be as short as possible.

The patients should be followed closely for signs and symptoms of respiratory depression and sedation. In this respect, it is strongly recommended to inform patients and their caregivers to be aware of these symptoms (see section 4.5).

Care should be taken when treating patients with respiratory depression, or if concomitant CNS depressant drugs are being administered (see section 4.5), or if the recommended dosage is significantly exceeded (see section 4.9) as the possibility of respiratory depression cannot be excluded in these situations.

Sleep-related breathing disorders

Opioids can cause sleep-related breathing disorders including central sleep apnea (CSA) and sleep-related hypoxemia. Opioid use increases the risk of CSA in a dose-dependent fashion. In patients who present with CSA, consider decreasing the total opioid dosage.

Convulsions have been reported in patients receiving tramadol at the recommended dose levels. The risk may be increased when doses of tramadol exceed the recommended upper daily dose limit (400 mg). In addition, tramadol may increase the seizure risk in patients taking other medicinal products that lowers the seizure threshold (see section 4.5). Patients with epilepsy or those susceptible to seizures should be only treated with tramadol if there are compelling circumstances.

Serotonin syndrome

Serotonin syndrome, a potentially life-threatening condition, has been reported in patients receiving tramadol in combination with other serotonergic agents or tramadol alone (see sections 4.5, 4.8 and 4.9).

If concomitant treatment with other serotonergic agents is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose escalations.

Symptoms of serotonin syndrome may include mental status changes, autonomic instability, neuromuscular abnormalities and/or gastrointestinal symptoms.

Serotonin syndrome is likely when one of the following is observed:

If serotonin syndrome is suspected, a dose reduction or discontinuation of therapy should be considered depending on the severity of the symptoms. Withdrawal of the serotonergic drugs usually brings about a rapid improvement.

Drug dependence, tolerance and potential for abuse

For all patients, prolonged use of this product may lead to drug dependence (addiction), even at therapeutic doses. The risks are increased in individuals with current or past history of substance misuse disorder (including alcohol misuse) or mental health disorder (e.g., major depression). Additional support and monitoring may be necessary when prescribing for patients at risk of opioid misuse.

Drug withdrawal syndrome

Prior to starting treatment with any opioids, a discussion should be held with patients to put in place a withdrawal strategy for ending treatment with tramadol.

Drug withdrawal syndrome may occur upon abrupt cessation of therapy or dose reduction. When a patient no longer requires therapy, it is advisable to taper the dose gradually to minimise symptoms of withdrawal. Tapering from a high dose may take weeks to months.

The opioid drug withdrawal syndrome is characterised by some or all of the following: restlessness, lacrimation, rhinorrhoea, yawning, perspiration, chills, myalgia, mydriasis and palpitations. Other symptoms may also develop including irritability, agitation, anxiety, hyperkinesia, tremor, weakness, insomnia, anorexia, abdominal cramps, nausea, vomiting, diarrhoea, increased blood pressure, increased respiratory rate or heart rate.

If women take this drug during pregnancy, there is a risk that their newborn infants will experience neonatal withdrawal syndrome.

Tramadol is not suitable as a substitute in opioid-dependent patients. Although it is an opioid agonist, tramadol cannot suppress morphine withdrawal symptoms

Children with compromised respiratory function

Tramadol is not recommended for use in children in whom respiratory function might be compromised including neuromuscular disorders, severe cardiac or respiratory conditions, upper respiratory or lung infections, multiple trauma or extensive surgical procedures. These factors may worsen symptoms of opioid toxicity.

This medicinal product contains lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Tramadol should not be combined with MAO inhibitors (see section 4.3).

In patients treated with MAO inhibitors in the 14 days prior to the use of the opioid pethidine, life-threatening interactions on the central nervous system, respiratory and cardiovascular function have been observed. The same interactions with MAO inhibitors cannot be ruled out during treatment with tramadol.

Concomitant administration of tramadol with other centrally depressant medicinal products including alcohol may potentiate the CNS effects (see section 4.8).

The concomitant use of opioids with sedating medicinal products such as benzodiazepines or related substances increases the risk of sedation, respiratory depression, coma and death because of additive CNS depressant effect.

The dose of tramadol and the duration of the concomitant use should be limited (see section 4.4).

The results of pharmacokinetic studies have so far shown that on the concomitant or previous administration of cimetidine (enzyme inhibitor) clinically relevant interactions are unlikely to occur. Simultaneous or previous administration of carbamazepine (enzyme inducer) may reduce the analgesic effect and shorten the duration of action.

Tramadol can induce convulsions and increase the potential for selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants, antipsychotics and other seizure threshold-lowering medicinal product (such as bupropion, mirtazapine, tetrahydrocannabinol) to cause convulsions.

Concomitant therapeutic use of tramadol and serotonergic drugs, such as selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors (SNRIs), MAO inhibitors (see section 4.3), tricyclic antidepressants and mirtazapine may cause serotonin syndrome, a potentially life-threatening condition (see sections 4.4 and 4.8).

Caution should be exercised during concomitant treatment with tramadol and coumarin derivatives (e.g. warfarin) due to reports of increased INR with major bleeding and ecchymoses in some patients.

Other active substances known to inhibit CYP3A4, such as ketoconazole and erythromycin, might inhibit the metabolism of tramadol (N-demethylation) probably also the metabolism of the active O-demethylated metabolite. The clinical importance of such an interaction has not been studied (see section 4.8).

In a limited number of studies the pre- or postoperative application of the antiemetic 5-HT₃ antagonist ondansetron increased the requirement of tramadol in patients with postoperative pain.

4.6 Pregnancy and lactation

Pregnancy:

Animal studies with tramadol revealed at very high doses effects on organ development, ossification and neonatal mortality. Tramadol crosses the placenta. There is inadequate evidence available on the safety of tramadol in human pregnancy. Therefore tramadol should not be used in pregnant women.

Regular use during pregnancy may cause drug dependence in the foetus, leading to withdrawal symptoms in the neonate.

If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available.

Tramadol - administered before or during birth - does not affect uterine contractility.

Administration during labour may depress respiration in the neonate and an antidote for the child should be readily available.

Breast-feeding

Administration to nursing women is not recommended as tramadol may be secreted in breast milk and may cause respiratory depression in the infant.

Fertility

Post marketing surveillance does not suggest an effect of tramadol on fertility. Animal studies did not show an effect of tramadol on fertility.

4.7 Effects on ability to drive and use machines

Even when taken according to instructions, tramadol may cause effects such as somnolence and dizziness and therefore may impair the reactions of drivers and machine operators. This applies particularly in conjunction with other psychotropic substances, particularly alcohol.

When prescribing this medicine, patients should be told:

- The medicine is likely to affect your ability to drive
- Do not drive until you know how the medicine affects you
- It is an offence to drive while under the influence of this medicine
- However, you would not be committing an offence (called 'statutory defence') if:

4.8 Undesirable effects

The most commonly reported adverse reactions are nausea and dizziness, both occurring in more than 10 % of patients.

The frequencies are defined as follows:

Very common: $\geq 1/10$

Common: $\geq 1/100, < 1/10$

Uncommon: $\geq 1/1000, < 1/100$

Rare: $\geq 1/10\ 000, < 1/1000$

Very rare: $< 1/10\ 000$

Not known: cannot be estimated from the available data

Cardiac disorders:

Uncommon: cardiovascular regulation (palpitation, tachycardia). These adverse reactions may occur especially on intravenous administration and in patients who are physically stressed.

Rare: bradycardia

Investigations:

Rare: increase in blood pressure

Vascular disorders:

Uncommon: cardiovascular regulation (postural hypotension or cardiovascular collapse). These adverse reactions may occur especially on intravenous administration and in patients who are physically stressed.

Metabolism and nutrition disorders:

Rare: changes in appetite

Respiratory, thoracic and mediastinal disorders:

Rare: respiratory depression, dyspnoea

If the recommended doses are considerably exceeded and other centrally depressant substances are administered concomitantly (see section 4.5), respiratory depression may occur.

Worsening of asthma has been reported, though a causal relationship has not been established.

Nervous system disorders:

Very common: dizziness

Common: headache, somnolence

Rare: speech disorders, paraesthesia, tremor, epileptiform convulsions, involuntary muscle contractions, abnormal coordination, syncope.

Not known: Serotonin syndrome

Convulsions occurred mainly after administration of high doses of tramadol or after concomitant treatment with medicinal products which can lower the seizure threshold (see sections 4.4 and 4.5).

Psychiatric disorders:

Rare: hallucinations, confusion, sleep disturbance, delirium, anxiety and nightmares. Psychic adverse reactions may occur following administration of tramadol which vary individually in intensity and nature (depending on personality and duration of treatment). These include changes in mood (usually elation, occasionally dysphoria), changes in activity (usually suppression, occasionally increase) and changes in cognitive and sensorial capacity (e.g. decision behaviour, perception disorders).

Frequency unknown: drug dependence (see section 4.4)

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after the authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions to the appropriate authority.

4.9 Overdose

Patients should be informed of the signs and symptoms of overdose and to ensure that family and friends are also aware of these signs and to seek immediate medical help if they occur.

Symptoms

In principle, on intoxication with tramadol symptoms similar to those of other centrally acting analgesics (opioids) are to be expected. These include in particular miosis, vomiting, cardiovascular collapse, consciousness disorders up to coma, convulsions and respiratory depression up to respiratory arrest.

Serotonin syndrome has also been reported.

Treatment

The general emergency measures apply. Keep open the respiratory tract (aspiration!), maintain respiration and circulation depending on the symptoms. The antidote for respiratory depression is naloxone. In animal experiments naloxone had no effect on convulsions. In such cases diazepam should be given intravenously.

In case of intoxication orally, gastrointestinal decontamination with activated charcoal or by gastric lavage is only recommended within 2 hours after tramadol intake. Gastrointestinal decontamination at a later time point may be useful in case of intoxication with exceptionally large quantities or prolonged-release formulations.

Tramadol is minimally eliminated from the serum by haemodialysis or haemo-filtration. Therefore treatment of acute intoxication with Tramadol with haemodialysis or haemofiltration alone is not suitable for detoxification.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: other opioids; ATC code: N02 AX02

Tramadol is a centrally acting opioid analgesic. It is a non-selective pure agonist at μ , δ and κ opioid receptors with a higher affinity for the μ receptor. Other mechanisms which contribute to its analgesic effect are inhibition of neuronal reuptake of noradrenaline and enhancement of serotonin release.

Tramadol has an antitussive effect. In contrast to morphine, analgesic doses of tramadol over a wide range have no respiratory depressant effect. Also gastrointestinal motility is less affected. Effects on the cardiovascular system tend to be slight. The potency of tramadol is reported to be 1/10 (one tenth) to 1/6 (one sixth) that of morphine.

Paediatric population

Effects of enteral and parenteral administration of tramadol have been investigated in clinical trials involving more than 2000 paediatric patients ranging in age from neonate to 17 years of age. The indications for pain treatment studied in those trials included pain after surgery (mainly abdominal), after surgical tooth extractions, due to fractures, burns and traumas as well as other painful conditions likely to require analgesic treatment for at least 7 days.

At single doses of up to 2 mg/kg or multiple doses of up to 8 mg/kg per day (to a maximum of 400 mg per day) efficacy of tramadol was found to be superior to placebo, and superior or equal to paracetamol, nalbuphine, pethidine or low dose morphine. The conducted trials confirmed the efficacy of tramadol. The safety profile of tramadol was similar in adult and paediatric patients older than 1 year (see section 4.2).

5.2 Pharmacokinetic properties

More than 90% of Tramadol HCL is absorbed after oral administration. The mean absolute bioavailability is approximately 70 %, irrespective of the concomitant intake of food. The difference between absorbed and non-metabolised available tramadol is probably due to the low first-pass effect. The first-pass effect after oral administration is a maximum of 30 %.

Tramadol has a high tissue affinity ($V_{d,B} = 203 \pm 40$ l). It has a plasma protein binding of about 20 %.

After administration of Tramadol 100 mg the peak plasma concentration $C_{max} = 141 \pm 40$ ng/ml is reached after 4.9 h. Tramadol passes the blood-brain and placental barriers. Very small amounts of the substance and its O-desmethyl derivative are found in the breast-milk (0.1 % and 0.02 % respectively of the applied dose).

Elimination half-life $t_{1/2,B}$ is approximately 6 h, irrespective of the mode of administration. In patients above 75 years of age it may be prolonged by a factor of approximately 1.4.

In humans tramadol is mainly metabolised by means of N- and O-demethylation and conjugation of the O-demethylation products with glucuronic acid. Only O-desmethyltramadol is pharmacologically active. There are considerable interindividual quantitative differences between the other metabolites. So far, eleven metabolites have been found in the urine. Animal experiments have shown that O-desmethyltramadol is more potent than the parent substance by the factor 2 - 4. Its half-life $t_{1/2,B}$ (6 healthy volunteers) is 7.9 h (range 5.4 - 9.6 h) and is approximately that of tramadol.

The inhibition of one or both types of the isoenzymes CYP3A4 and CYP2D6 involved in the biotransformation of tramadol may affect the plasma concentration of tramadol or its active metabolite.

Tramadol and its metabolites are almost completely excreted via the kidneys. Cumulative urinary excretion is 90 % of the total radioactivity of the administered dose. In cases of impaired hepatic and renal function the half-life may be slightly prolonged. In patients with cirrhosis of the liver, elimination half-lives of 13.3 ± 4.9 h (tramadol) and 18.5 ± 9.4 h (O-desmethyltramadol), in an extreme case 22.3 h and 36 h respectively, have been determined. In patients with renal insufficiency (creatinine clearance < 5 ml/min) the values were 11 ± 3.2 h and 16.9 ± 3 h, in an extreme case 19.5 h and 43.2 h respectively.

Tramadol has a linear pharmacokinetic profile within the therapeutic dosage range.

The relationship between serum concentrations and the analgesic effect is dose-dependent, but varies considerably in isolated cases. A serum concentration of 100 - 300 ng/ml is usually effective.

Paediatric population

The pharmacokinetics of tramadol and O-desmethyltramadol after single-dose and multiple-dose oral administration to subjects aged 1 year to 16 years were found to be generally similar to those

in adults when adjusting for dose by body weight, but with a higher between-subject variability in children aged 8 years and below.

In children below 1 year of age, the pharmacokinetics of tramadol and O-desmethyltramadol have been investigated, but have not been fully characterized. Information from studies including this age group indicates that the formation rate of O-desmethyltramadol via CYP2D6 increases continuously in neonates, and adult levels of CYP2D6 activity are assumed to be reached at about 1 year of age. In addition, immature glucuronidation systems and immature renal function may result in slow elimination and accumulation of O-desmethyltramadol in children under 1 year of age.

5.3 Preclinical safety data

Not applicable

6. Pharmaceutical particulars

6.1 List of excipients

Colloidal Silicon Dioxide, Talcum, Micro Crystalline Cellulose, Sodium starch glycolate and Magnesium Stearate, Empty hard gelatin capsule

6.2 Incompatibilities

None known

6.3 Shelf life

36 months

6.4 Special precautions for storage

Store in a cool dry place at temperature below 30 °C. Store in the original packaging.

6.5 Nature and contents of container

PVC pack in a printed carton with a printed insert.

Pack size: Blister pack of 100 x 10's.

6.6 Special precautions for disposal and other handling

Not applicable.

7. Marketing authorization holder

Me Cure Industries Limited

Plot 6 Block H, Debo Industries Compound,

Oshodi Industrial Scheme, Oshodi,

Lagos, Nigeria.

8.0 NAFDAC Registration Number: A4-6295