



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## **MODULE-1 ADMINISTRATIVE INFORMATION**

### **1.3.1 Summary Of Product Characteristics (Smpe)**

#### **1. Name of drug product**

CEFTRIAZONE FOR INJECTION USP 1 GM

#### **1.1 (Trade) name of product**

NORACEF

#### **1.2 Strength**

Each vial Contains:

Ceftriazone sodium (sterile) USP

Eq. to Anhydrous Ceftriazone..... 1000 mg

#### **1.3 Pharmaceutical Dosage Form**

Powder for injection



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## **MODULE-1 ADMINISTRATIVE INFORMATION**

### **2. Qualitative & Quantitative composition**

#### **2.1 Qualitative Declaration**

Each vial Contains:

Ceftriaxone sodium (sterile) USP

Eq. to Anhydrous Ceftriaxone..... 1000 mg

#### **2.2 Quantitative Declaration**

##### **Batch Formula:**

**Batch size: 10,000 Vial**

<b>Sr. No.</b>	<b>Ingredients</b>	<b>Specification</b>	<b>Rational</b>	<b>Label claim</b>	<b>Overages</b>	<b>Quantity/ Unit (mg)</b>	<b>Quantity/ Batch (Kg)</b>
<b>1.</b>	Ceftriaxone sodium (sterile) Eq. to Anhydrous Ceftriaxone	USP	Active	1000 mg	---	1200.00	12.00



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## **MODULE-1 ADMINISTRATIVE INFORMATION**

### **3. Pharmaceutical dosage form**

IV

### **4. Clinical particulars**

#### **4.1 Therapeutic indications**

Ceftriaxone is indicated in the treatment of the following infections in adults and children including term neonates (from birth):

Bacterial Meningitis

Community acquired pneumonia

Hospital acquired pneumonia

Acute otitis media

Intra-abdominal infections

Complicated urinary tract infections (including pyelonephritis)

Infections of bones and joints

Complicated skin and soft tissue infections

Gonorrhoea

Syphilis

Bacterial endocarditis

Ceftriaxone may be used:

For treatment of acute exacerbations of chronic obstructive pulmonary disease in adults

For treatment of disseminated Lyme borreliosis (early (stage II) and late (stage III)) in adults and children including neonates from 15 days of age.

For Pre-operative prophylaxis of surgical site infections

In the management of neutropenic patients with fever that is suspected to be due to a bacterial infection

In the treatment of patients with bacteraemia that occurs in association with, or is suspected to be associated with, any of the infections listed above

Ceftriaxone should be co-administered with other antibacterial agents whenever the possible range of causative bacteria would not fall within its spectrum (see section 4.4).

Consideration should be given to official guidance on the appropriate use of antibacterial agents.



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## MODULE-1 ADMINISTRATIVE INFORMATION

### 4.2 Posology and method of administration

#### Posology

The dose depends on the severity, susceptibility, site and type of infection and on the age and hepato-renal function of the patient.

The doses recommended in the tables below are the generally recommended doses in these indications. In particularly severe cases, doses at the higher end of the recommended range should be considered.

Adults and children over 12 years of age ( $\geq 50$  kg)

<b>Ceftriaxone Dosage*</b>	<b>Treatment frequency**</b>	<b>Indications</b>
1-2 g	Once daily	Community acquired pneumonia
		Acute exacerbations of chronic obstructive pulmonary disease
		Intra-abdominal infections
		Complicated urinary tract infections (including pyelonephritis)
2 g	Once daily	Hospital acquired pneumonia
		Complicated skin and soft tissue infections
		Infections of bones and joints
2-4 g	Once daily	Management of neutropenic patients with fever that is suspected to be due to a bacterial infection
		Bacterial endocarditis
		Bacterial meningitis

\* In documented bacteraemia, the higher end of the recommended dose range should be considered.

\*\* Twice daily (12 hourly) administration may be considered where doses greater than 2 g daily are administered.

Indications for adults and children over 12 years of age ( $\geq 50$  kg) that require specific dosage schedules:

#### Syphilis

The generally recommended doses are 500 mg-1 g once daily increased to 2 g once daily for neurosyphilis for 10-14 days. The dose recommendations in syphilis, including neurosyphilis, are based on limited data. National or local guidance should be taken into consideration.

Disseminated Lyme borreliosis (early [Stage II] and late [Stage III])



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAXONE FOR INJECTION USP 1 GM</b>

### MODULE-1 ADMINISTRATIVE INFORMATION

2 g once daily for 14-21 days. The recommended treatment durations vary and national or local guidelines should be taken into consideration.

Paediatric population

Neonates, infants and children 15 days to 12 years of age (< 50 kg)

For children with bodyweight of 50 kg or more, the usual adult dosage should be given.

Ceftriaxone dosage*	Treatment frequency**	Indications
50-80 mg/kg	Once daily	Intra-abdominal infections
		Complicated urinary tract infections (including pyelonephritis)
		Community acquired pneumonia
		Hospital acquired pneumonia
50-100 mg/kg (Max 4 g)	Once daily	Complicated skin and soft tissue infections
		Infections of bones and joints
		Management of neutropenic patients with fever that is suspected to be due to a bacterial infection
80-100 mg/kg (max 4 g)	Once daily	Bacterial meningitis
100 mg/kg (max 4 g)	Once daily	Bacterial endocarditis

\* In documented bacteraemia, the higher end of the recommended dose range should be considered.

\*\* Twice daily (12 hourly) administration may be considered where doses greater than 2 g daily are administered.

Indications for neonates, infants and children 15 days to 12 years (< 50 kg) that require specific dosage schedules:

Syphilis

The generally recommended doses are 75-100 mg/kg (max 4 g) once daily for 10-14 days. The dose recommendations in syphilis, including neurosyphilis, are based on very limited data. National or local guidance should be taken into consideration.

Disseminated Lyme borreliosis (early [Stage II] and late [Stage III])

50–80 mg/kg once daily for 14-21 days. The recommended treatment durations vary and national or local guidelines should be taken into consideration.

Neonates 0-14 days

Ceftriaxone is contraindicated in premature neonates up to a postmenstrual age of 41 weeks (gestational age + chronological age).



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAXONE FOR INJECTION USP 1 GM</b>

## MODULE-1 ADMINISTRATIVE INFORMATION

Ceftriaxone dosage*	Treatment frequency	Indications
20-50 mg/kg	Once daily	Intra-abdominal infections
		Complicated skin and soft tissue infections
		Complicated urinary tract infections (including pyelonephritis)
		Community acquired pneumonia
		Hospital acquired pneumonia
		Infections of bones and joints
		Management of neutropenic patients with fever that is suspected to be due to a bacterial infection
50 mg/kg	Once daily	Bacterial meningitis
		Bacterial endocarditis

\* In documented bacteraemia, the higher end of the recommended dose range should be considered.

A maximum daily dose of 50 mg/kg should not be exceeded.

Indications for neonates 0-14 days that require specific dosage schedules:

Acute otitis media

Pre-operative prophylaxis of surgical site infections

20-50 mg/kg as a single pre-operative dose.

Syphilis

The generally recommended dose is 50 mg/kg once daily for 10-14 days. The dose recommendations in syphilis, including neurosyphilis, are based on very limited data. National or local guidance should be taken into consideration.

Duration of therapy

The duration of therapy varies according to the course of the disease. As with antibiotic therapy in general, administration of ceftriaxone should be continued for 48 - 72 hours after the patient has become afebrile or evidence of bacterial eradication has been achieved.

Older people

The dosages recommended for adults require no modification in older people provided that renal and hepatic function is satisfactory.

Patients with hepatic impairment

Available data do not indicate the need for dose adjustment in mild or moderate liver function impairment provided renal function is not impaired.



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

### **MODULE-1 ADMINISTRATIVE INFORMATION**

There are no study data in patients with severe hepatic impairment (see section 5.2).

Patients with renal impairment:

In patients with impaired renal function, there is no need to reduce the dosage of ceftriaxone provided hepatic function is not impaired. Only in cases of preterminal renal failure (creatinine clearance < 10 ml/min) should the ceftriaxone dosage not exceed 2 g daily.

In patients undergoing dialysis no additional supplementary dosing is required following the dialysis. Ceftriaxone is not removed by peritoneal- or haemodialysis. Close clinical monitoring for safety and efficacy is advised.

Patients with severe hepatic and renal impairment

In patients with both severe renal and hepatic dysfunction, close clinical monitoring for safety and efficacy is advised.

#### **Method of administration**

Intravenous administration

For IV injection 1 g ceftriaxone is dissolved in 10 ml of water for injections PhEur. The injection should be administered over 5 minutes, directly into the vein or via the tubing of an intravenous infusion.

### **4.3 Contraindications**

Hypersensitivity to the active substance, to any other cephalosporin or to any of the excipients listed in section 6.1

History of severe hypersensitivity (e.g. anaphylactic reaction) to any other type of beta-lactam antibacterial agent (penicillins, monobactams and carbapenems).

Ceftriaxone is contraindicated in:

- Premature neonates up to a postmenstrual age of 41 weeks (gestational age + chronological age)\*
- Full-term neonates (up to 28 days of age):
  - with hyperbilirubinaemia, jaundice, or who are hypoalbuminaemic or acidotic because these are conditions in which bilirubin binding is likely to be impaired\*
  - if they require (or are expected to require) intravenous calcium treatment, or calcium-containing infusions due to the risk of precipitation of a ceftriaxone-calcium salt (see sections 4.4, 4.8 and 6.2).



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## **MODULE-1 ADMINISTRATIVE INFORMATION**

\* In vitro studies have shown that ceftriaxone can displace bilirubin from its serum albumin binding sites leading to a possible risk of bilirubin encephalopathy in these patients.

### **4.4 Special warnings and precautions for use**

#### **Hypersensitivity reactions**

As with all beta-lactam antibacterial agents, serious and occasionally fatal hypersensitivity reactions have been reported (see section 4.8). In case of severe hypersensitivity reactions, treatment with ceftriaxone must be discontinued immediately and adequate emergency measures must be initiated. Before beginning treatment, it should be established whether the patient has a history of severe hypersensitivity reactions to ceftriaxone, to other cephalosporins or to any other type of beta-lactam agent. Caution should be used if ceftriaxone is given to patients with a history of non-severe hypersensitivity to other beta-lactam agents.

Severe cutaneous adverse reactions (Stevens Johnson syndrome or Lyell's syndrome/toxic epidermal necrolysis) and drug reaction with eosinophilia and systemic symptoms (DRESS)) which can be life-threatening or fatal, have been reported in association with ceftriaxone treatment; however, the frequency of these events is not known (see section 4.8).

#### **Jarisch-Herxheimer reaction (JHR)**

Some patients with spirochete infections may experience a Jarisch-Herxheimer reaction (JHR) shortly after ceftriaxone treatment is started. JHR is usually a self-limiting condition or can be managed by symptomatic treatment. The antibiotic treatment should not be discontinued if such reaction occurs.'

#### **Interaction with calcium containing products**

Cases of fatal reactions with calcium-ceftriaxone precipitates in lungs and kidneys in premature and full-term neonates aged less than 1 month have been described. At least one of them had received ceftriaxone and calcium at different times and through different intravenous lines. In the available scientific data, there are no reports of confirmed intravascular precipitations in patients, other than neonates, treated with ceftriaxone and calcium-containing solutions or any other calcium-containing products. In vitro studies demonstrated that neonates have an increased risk of precipitation of ceftriaxone-calcium compared to other age groups.

In patients of any age ceftriaxone must not be mixed or administered simultaneously with any calcium-containing intravenous solutions, even via different infusion lines or at different infusion





<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

### **MODULE-1 ADMINISTRATIVE INFORMATION**

sites. However, in patients older than 28 days of age ceftriaxone and calcium-containing solutions may be administered sequentially one after another if infusion lines at different sites are used or if the infusion lines are replaced or thoroughly flushed between infusions with physiological salt-solution to avoid precipitation. In patients requiring continuous infusion with calcium-containing total parenteral nutrition (TPN) solutions, healthcare professionals may wish to consider the use of alternative antibacterial treatments which do not carry a similar risk of precipitation. If the use of ceftriaxone is considered necessary in patients requiring continuous nutrition, TPN solutions and ceftriaxone can be administered simultaneously, albeit via different infusion lines at different sites. Alternatively, infusion of TPN solution could be stopped for the period of ceftriaxone infusion and the infusion lines flushed between solutions (see sections 4.3, 4.8, 5.2 and 6.2).

#### **Paediatric population**

Safety and effectiveness of Ceftriaxone in neonates, infants and children have been established for the dosages described under Posology and Method of Administration (see section 4.2). Studies have shown that ceftriaxone, like some other cephalosporins, can displace bilirubin from serum albumin.

Ceftriaxone is contraindicated in premature and full-term neonates at risk of developing bilirubin encephalopathy (see section 4.3).

#### **Immune mediated haemolytic anaemia**

An immune mediated haemolytic anaemia has been observed in patients receiving cephalosporin class antibacterials including Ceftriaxone (see section 4.8). Severe cases of haemolytic anaemia, including fatalities, have been reported during Ceftriaxone treatment in both adults and children. If a patient develops anaemia while on ceftriaxone, the diagnosis of a cephalosporin-associated anaemia should be considered and ceftriaxone discontinued until the aetiology is determined.

#### **Long term treatment**

During prolonged treatment complete blood count should be performed at regular intervals.

#### **Colitis/Overgrowth of non-susceptible microorganisms**

Antibacterial agent-associated colitis and pseudo-membranous colitis have been reported with nearly all antibacterial agents, including ceftriaxone, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhoea during or subsequent to the administration of ceftriaxone (see section 4.8). Discontinuation of therapy with ceftriaxone and the administration of specific treatment for



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

### **MODULE-1 ADMINISTRATIVE INFORMATION**

Clostridium difficile should be considered. Medicinal products that inhibit peristalsis should not be given.

Superinfections with non-susceptible micro-organisms may occur as with other antibacterial agents.

Severe renal and hepatic insufficiency

In severe renal and hepatic insufficiency, close clinical monitoring for safety and efficacy is advised (see section 4.2).

Interference with serological testing

Interference with Coombs tests may occur, as Ceftriaxone may lead to false-positive test results. Ceftriaxone can also lead to false-positive test results for galactosaemia (see section 4.8).

Non-enzymatic methods for the glucose determination in urine may give false-positive results. Urine glucose determination during therapy with Ceftriaxone should be done enzymatically. The presence of ceftriaxone may falsely lower estimated blood glucose values obtained with some blood glucose monitoring systems. Please refer to instructions for use for each system. Alternative testing methods should be used if necessary.

Sodium

This medicinal product contains 82mg sodium per 1g vial, equivalent to 4.1% of the WHO recommended maximum daily intake of 2 g sodium for an adult.

Antibacterial spectrum

Ceftriaxone has a limited spectrum of antibacterial activity and may not be suitable for use as a single agent for the treatment of some types of infections unless the pathogen has already been confirmed (see section 4.2). In polymicrobial infections, where suspected pathogens include organisms resistant to ceftriaxone, administration of an additional antibiotic should be considered.

Use of lidocaine

Biliary lithiasis

When shadows are observed on sonograms, consideration should be given to the possibility of precipitates of calcium ceftriaxone. Shadows, which have been mistaken for gallstones, have been detected on sonograms of the gallbladder and have been observed more frequently at ceftriaxone doses of 1 g per day and above. Caution should be particularly considered in the paediatric population. Such precipitates disappear after discontinuation of ceftriaxone therapy. Rarely precipitates of calcium ceftriaxone have been associated with symptoms. In symptomatic cases,



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

### **MODULE-1 ADMINISTRATIVE INFORMATION**

conservative nonsurgical management is recommended and discontinuation of ceftriaxone treatment should be considered by the physician based on specific benefit risk assessment (see section 4.8).

#### **Biliary stasis**

Cases of pancreatitis, possibly of biliary obstruction aetiology, have been reported in patients treated with Ceftriaxone (see section 4.8). Most patients presented with risk factors for biliary stasis and biliary sludge e.g. preceding major therapy, severe illness and total parenteral nutrition. A trigger or cofactor of Ceftriaxone-related biliary precipitation cannot be ruled out.

#### **Renal lithiasis**

Cases of renal lithiasis have been reported, which is reversible upon discontinuation of ceftriaxone (see section 4.8). In symptomatic cases, sonography should be performed. Use in patients with history of renal lithiasis or with hypercalciuria should be considered by the physician based on specific benefit risk assessment.

#### **Encephalopathy**

Encephalopathy has been reported with the use of ceftriaxone (see section 4.8), particularly in elderly patients with severe renal impairment (see section 4.2) or central nervous system disorders. If ceftriaxone-associated encephalopathy is suspected (e.g. decreased level of consciousness, altered mental state, myoclonus, convulsions), discontinuation of ceftriaxone should be considered.

### **4.5 Interaction with other medicinal products and other forms of interaction**

Calcium-containing diluents, such as Ringer's solution or Hartmann's solution, should not be used to reconstitute Ceftriaxone vials or to further dilute a reconstituted vial for intravenous administration because a precipitate can form.

Precipitation of ceftriaxone-calcium can also occur when ceftriaxone is mixed with calcium-containing solutions in the same intravenous administration line.

Ceftriaxone must not be administered simultaneously with calcium-containing intravenous solutions, including continuous calcium-containing infusions such as parenteral nutrition via a Y-site. However, in patients other than neonates, ceftriaxone and calcium-containing solutions may be administered sequentially of one another if the infusion lines are thoroughly flushed between infusions with a compatible fluid.



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAXONE FOR INJECTION USP 1 GM</b>

## **MODULE-1 ADMINISTRATIVE INFORMATION**

In vitro studies using adult and neonatal plasma from umbilical cord blood demonstrated that neonates have an increased risk of precipitation of ceftriaxone-calcium (see sections 4.2, 4.3, 4.4, 4.8 and 6.2).

Concomitant use with oral anticoagulants may increase the anti-vitamin K effect and the risk of bleeding. It is recommended that the International Normalised Ratio (INR) is monitored frequently and the posology of the anti-vitamin K drug adjusted accordingly, both during and after treatment with ceftriaxone (see section 4.8).

There is conflicting evidence regarding a potential increase in renal toxicity of aminoglycosides when used with cephalosporins. The recommended monitoring of aminoglycoside levels (and renal function) in clinical practice should be closely adhered to in such cases.

In an in-vitro study antagonistic effects have been observed with the combination of chloramphenicol and ceftriaxone. The clinical relevance of this finding is unknown.

In patients treated with ceftriaxone, the Coombs' test may lead to false-positive test results.

Ceftriaxone, like other antibiotics, may result in false-positive tests for galactosaemia.

Likewise, non-enzymatic methods for glucose determination in urine may yield false-positive results. For this reason, glucose level determination in urine during therapy with ceftriaxone should be carried out enzymatically.

No impairment of renal function has been observed after concurrent administration of large doses of ceftriaxone and potent diuretics (e.g. furosemide).

Simultaneous administration of probenecid does not reduce the elimination of ceftriaxone.

## **4.6 Fertility, pregnancy and lactation**

### **Pregnancy**

Ceftriaxone crosses the placental barrier. There are limited amounts of data from the use of ceftriaxone in pregnant women. Animal studies do not indicate direct or indirect harmful effects with respect to embryonal/foetal, perinatal and postnatal development (see section 5.3). Ceftriaxone should only be administered during pregnancy and in particular in the first trimester of pregnancy if the benefit outweighs the risk.



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## MODULE-1 ADMINISTRATIVE INFORMATION

### Breastfeeding

Ceftriaxone is excreted into human milk in low concentrations but at therapeutic doses of ceftriaxone no effects on the breastfed infants are anticipated. However, a risk of diarrhoea and fungal infection of the mucous membranes cannot be excluded. The possibility of sensitisation should be taken into account. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from ceftriaxone therapy, taking into account the benefit of breast feeding for the child and the benefit of therapy for the woman.

### Fertility

Reproductive studies have shown no evidence of adverse effects on male or female fertility

### 4.7 Effects on ability to drive and use machines

During treatment with ceftriaxone, undesirable effects may occur (e.g. dizziness), which may influence the ability to drive and use machines (see section 4.8). Patients should be cautious when driving or operating machinery.

### 4.8 Undesirable effects

The most frequently reported adverse reactions for ceftriaxone are eosinophilia, leucopenia, thrombocytopenia, diarrhoea, rash, and hepatic enzymes increased.

Data to determine the frequency of ceftriaxone ADRs was derived from clinical trials.

The following convention has been used for the classification of frequency:

Very common ( $\geq 1/10$ )

Common ( $\geq 1/100$  -  $< 1/10$ )

Uncommon ( $\geq 1/1000$  -  $< 1/100$ )

Rare ( $\geq 1/10000$  -  $< 1/1000$ )

Not known (cannot be estimated from the available data)

System Organ Class	Common	Uncommon	Rare	Not Known <sup>a</sup>
Infections and infestations		Genital fungal infection	Pseudomembranous colitis <sup>b</sup>	Superinfection <sup>b</sup>
Blood and lymphatic system disorders	Eosinophilia Leucopenia Thrombocytopenia	Granulocytopenia Anaemia Coagulopathy		Haemolytic anaemia <sup>b</sup> Agranulocytosis



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

### MODULE-1 ADMINISTRATIVE INFORMATION

Immune system disorders				Anaphylactic shock Anaphylactic reaction Anaphylactoid reaction Hypersensitivity <sup>b</sup> Jarisch-Herxheimer reaction <sup>b</sup>
Nervous system disorders		Headache Dizziness	Encephalopathy	Convulsion
Ear and labyrinth disorders				Vertigo
Respiratory, thoracic and mediastinal disorders			Bronchospasm	
Gastrointestinal disorders	Diarrhoea <sup>b</sup> Loose stools	Nausea Vomiting		Pancreatitis <sup>b</sup> Stomatitis Glossitis
Hepatobiliary disorders	Hepatic enzyme increased			Gall bladder precipitation <sup>b</sup> Kernicterus Hepatitis <sup>c</sup> Hepatitis cholestatic <sup>b,c</sup>
Skin and subcutaneous tissue disorders	Rash	Pruritus	Urticaria	Stevens Johnson Syndrome <sup>b</sup> Toxic epidermal necrolysis <sup>b</sup> Erythema multiforme Acute generalised exanthematous pustulosis drug reaction with eosinophilia and systemic symptoms (DRESS) <sup>b</sup>
Renal and urinary disorders			Haematuria Glycosuria	Oliguria Renal precipitation (reversible)
General disorders and administration site conditions		Phlebitis Injection site pain Pyrexia	Oedema Chills	



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## MODULE-1 ADMINISTRATIVE INFORMATION

Investigations		Blood creatinine increased		Coombs test false positive <sup>b</sup> Galactosaemia test false positive <sup>b</sup> Non enzymatic methods for glucose determination false positive <sup>b</sup>
----------------	--	----------------------------	--	--

<sup>a</sup>Based on post-marketing reports. Since these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency which is therefore categorised as not known.

<sup>b</sup> See section 4.4

<sup>c</sup> Usually reversible upon discontinuation of ceftriaxone

Description of selected adverse reactions

### Infections and infestations

Reports of diarrhoea following the use of ceftriaxone may be associated with *Clostridium difficile*.

Appropriate fluid and electrolyte management should be instituted (see section 4.4).

### Ceftriaxone-calcium salt precipitation

Rarely, severe, and in some cases, fatal, adverse reactions have been reported in pre-term and full-term neonates (aged < 28 days) who had been treated with intravenous ceftriaxone and calcium. Precipitations of ceftriaxone-calcium salt have been observed in lung and kidneys post-mortem. The high risk of precipitation in neonates is a result of their low blood volume and the longer half-life of ceftriaxone compared with adults (see sections 4.3, 4.4, and 5.2).

Cases of ceftriaxone precipitation in the urinary tract have been reported, mostly in children treated with high doses (e.g.  $\geq 80$  mg/kg/day or total doses exceeding 10 grams) and who have other risk factors (e.g. dehydration, confinement to bed). This event may be asymptomatic or symptomatic, and may lead to ureteric obstruction and postrenal acute renal failure, but is usually reversible upon discontinuation of ceftriaxone (see section 4.4).

Precipitation of ceftriaxone calcium salt in the gallbladder has been observed, primarily in patients treated with doses higher than the recommended standard dose. In children, prospective studies have shown a variable incidence of precipitation with intravenous application - above 30 % in some studies. The incidence appears to be lower with slow infusion (20 - 30 minutes). This effect is usually asymptomatic, but the precipitations have been accompanied by clinical symptoms such as pain, nausea and vomiting in rare cases. Symptomatic treatment is recommended in these cases. Precipitation is usually reversible upon discontinuation of ceftriaxone (see section 4.4).





<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## MODULE-1 ADMINISTRATIVE INFORMATION

### 4.9 Overdose

In overdose, the symptoms of nausea, vomiting and diarrhoea can occur. Ceftriaxone concentrations cannot be reduced by haemodialysis or peritoneal dialysis. There is no specific antidote. Treatment is symptomatic.

## 5. Pharmacological properties

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antibacterials for systemic use, Third-generation cephalosporins

ATC code: J01DD04

#### Mechanism of action

Ceftriaxone inhibits bacterial cell wall synthesis following attachment to penicillin binding proteins (PBPs). This results in the interruption of cell wall (peptidoglycan) biosynthesis, which leads to bacterial cell lysis and death.

#### Resistance

Bacterial resistance to ceftriaxone may be due to one or more of the following mechanisms:

- hydrolysis by beta-lactamases, including extended-spectrum beta-lactamases (ESBLs), carbapenemases and Amp C enzymes that may be induced or stably derepressed in certain aerobic Gram-negative bacterial species.
- reduced affinity of penicillin-binding proteins for ceftriaxone.
- outer membrane impermeability in Gram-negative organisms.
- bacterial efflux pumps.

#### Susceptibility testing Breakpoints

Minimum inhibitory concentration (MIC) breakpoints established by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) are as follows:

Pathogen	Dilution Test (MIC, mg/L)	
	Susceptible	Resistant
Enterobacteriaceae	≤ 1	> 2
Staphylococcus spp	a.	a.
Streptococcus spp. (Groups A, B, C and G)	b.	b.
Streptococcus pneumoniae	≤ 0.5 <sup>c</sup>	> 2





<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAXONE FOR INJECTION USP 1 GM</b>

## MODULE-1 ADMINISTRATIVE INFORMATION

<i>Viridans group Streptococci</i>	≤0.5	>0.5
<i>Haemophilus influenzae</i>	≤ 0.12 <sup>c</sup>	> 0.12
<i>Moraxella catarrhalis</i>	≤ 1	> 2
<i>Neisseria gonorrhoeae</i>	≤ 0.12	> 0.12
<i>Neisseria meningitidis</i>	≤ 0.12 <sup>c</sup>	> 0.12
<i>Non-species related</i>	≤ 1 <sup>d</sup>	> 2

a. Susceptibility inferred from cefoxitin susceptibility.

b. Susceptibility inferred from penicillin susceptibility.

c. Isolates with a ceftriaxone MIC above the susceptible breakpoint are rare and, if found, should be re-tested and, if confirmed, should be sent to a reference laboratory.

d. Breakpoints apply to a daily intravenous dose of 1 g x 1 and a high dose of at least 2 g x 1.

Clinical efficacy against specific pathogens

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of ceftriaxone in at least some types of infections is questionable.

### Commonly susceptible species

Gram-positive aerobes  
*Staphylococcus aureus* (methicillin-susceptible)<sup>‡</sup>  
*Staphylococci coagulase-negative* (methicillin-susceptible)<sup>‡</sup>  
*Streptococcus pyogenes* (Group A)  
*Streptococcus agalactiae* (Group B)  
*Streptococcus pneumoniae*  
*Viridans Group Streptococci*  
Gram-negative aerobes  
*Borrelia burgdorferi*  
*Haemophilus influenzae*  
*Haemophilus parainfluenzae*  
*Moraxella catarrhalis*  
*Neisseria gonorrhoea*  
*Neisseria meningitidis*  
*Proteus mirabilis*  
*Providencia spp*  
*Treponema pallidum*

### Species for which acquired resistance may be a problem

Gram-positive aerobes  
*Staphylococcus epidermidis*<sup>+</sup>  
*Staphylococcus haemolyticus*<sup>+</sup>



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## MODULE-1 ADMINISTRATIVE INFORMATION

*Staphylococcus hominis*<sup>+</sup>  
*Gram-negative aerobes*  
*Citrobacter freundii*  
*Enterobacter aerogenes*  
*Enterobacter cloacae*  
*Escherichia coli*<sup>%</sup>  
*Klebsiella pneumoniae*<sup>%</sup>  
*Klebsiella oxytoca*<sup>%</sup>  
*Morganella morganii*  
*Proteus vulgaris*  
*Serratia*  
*marcescens*  
*Anaerobes*  
*Bacteroides spp.*  
*Fusobacterium spp.*  
*Peptostreptococcus*  
*spp.*  
*Clostridium perfringens*

### Inherently resistant organisms

*Gram-positive aerobes*  
*Enterococcus spp.*  
*Listeria monocytogenes*  
*Gram-negative aerobes*  
*Acinetobacter baumannii*  
*Pseudomonas*  
*aeruginosa*  
*Stenotrophomonas maltophilia*  
*Anaerobes*  
*Clostridium difficile*  
*Others:*  
*Chlamydia spp.*  
*Chlamydophila spp.*  
*Mycoplasma spp.*  
*Legionella spp.*  
*Ureaplasma*  
*urealyticum*

£ All methicillin-resistant staphylococci are resistant to ceftriaxone.

<sup>+</sup> Resistance rates >50% in at least one region

<sup>%</sup> ESBL producing strains are always resistant

## 5.2 Pharmacokinetic properties

### Absorption



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

#### **MODULE-1 ADMINISTRATIVE INFORMATION**

Intravenous administration

After intravenous bolus administration of ceftriazone 500 mg and 1 g, mean peak plasma ceftriazone levels are approximately 120 and 200 mg/l respectively. After intravenous infusion of



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

### **MODULE-1 ADMINISTRATIVE INFORMATION**

ceftriaxone 500 mg, 1 g and 2 g, the plasma ceftriaxone levels are approximately 80, 150 and 250 mg/l respectively.

#### **Distribution**

The volume of distribution of ceftriaxone is 7 – 12 l. Concentrations well above the minimal inhibitory concentrations of most relevant pathogens are detectable in tissue including lung, heart, biliary tract/liver, tonsil, middle ear and nasal mucosa, bone, and in cerebrospinal, pleural, prostatic and synovial fluids. An 8 - 15 % increase in mean peak plasma concentration ( $C_{max}$ ) is seen on repeated administration; steady state is reached in most cases within 48 - 72 hours depending on the route of administration.

#### **Penetration into particular tissues**

Ceftriaxone penetrates the meninges. Penetration is greatest when the meninges are inflamed. Mean peak ceftriaxone concentrations in CSF in patients with bacterial meningitis are reported to be up to 25 % of plasma levels compared to 2 % of plasma levels in patients with uninflamed meninges. Peak ceftriaxone concentrations in CSF are reached approximately 4-6 hours after intravenous injection. Ceftriaxone crosses the placental barrier and is excreted in the breast milk at low concentrations (see section 4.6).

#### **Protein binding**

Ceftriaxone is reversibly bound to albumin. Plasma protein binding is about 95 % at plasma concentrations below 100 mg/l. Binding is saturable and the bound portion decreases with rising concentration (up to 85 % at a plasma concentration of 300 mg/l).

#### **Biotransformation**

Ceftriaxone is not metabolised systemically; but is converted to inactive metabolites by the gut flora.

#### **Elimination**

Plasma clearance of total ceftriaxone (bound and unbound) is 10 - 22 ml/min. Renal clearance is 5 - 12 ml/min. 50 - 60 % of ceftriaxone is excreted unchanged in the urine, primarily by glomerular filtration, while 40 - 50 % is excreted unchanged in the bile. The elimination half-life of total ceftriaxone in adults is about 8 hours.

Patients with renal or hepatic impairment



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

### **MODULE-1 ADMINISTRATIVE INFORMATION**

In patients with renal or hepatic dysfunction, the pharmacokinetics of ceftriaxone are only minimally altered with the half-life slightly increased (less than two fold), even in patients with severely impaired renal function.

The relatively modest increase in half-life in renal impairment is explained by a compensatory increase in non-renal clearance, resulting from a decrease in protein binding and corresponding increase in non-renal clearance of total ceftriaxone.

In patients with hepatic impairment, the elimination half-life of ceftriaxone is not increased, due to a compensatory increase in renal clearance. This is also due to an increase in plasma free fraction of ceftriaxone contributing to the observed paradoxical increase in total drug clearance, with an increase in volume of distribution paralleling that of total clearance.

#### **Older people**

In older people aged over 75 years the average elimination half-life is usually two to three times that of young adults.

#### **Paediatric population**

The half-life of ceftriaxone is prolonged in neonates. From birth to 14 days of age, the levels of free ceftriaxone may be further increased by factors such as reduced glomerular filtration and altered protein binding. During childhood, the half-life is lower than in neonates or adults.

The plasma clearance and volume of distribution of total ceftriaxone are greater in neonates, infants and children than in adults.

#### **Linearity/non-linearity**

The pharmacokinetics of ceftriaxone are non-linear and all basic pharmacokinetic parameters, except the elimination half-life, are dose dependent if based on total drug concentrations, increasing less than proportionally with dose. Non-linearity is due to saturation of plasma protein binding and is therefore observed for total plasma ceftriaxone but not for free (unbound) ceftriaxone.

#### **Pharmacokinetic/pharmacodynamic relationship**

As with other beta-lactams, the pharmacokinetic-pharmacodynamic index demonstrating the best correlation with in vivo efficacy is the percentage of the dosing interval that the unbound concentration remains above the minimum inhibitory concentration (MIC) of ceftriaxone for individual target species (i.e. %T > MIC).



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## **MODULE-1 ADMINISTRATIVE INFORMATION**

### **5.3 Preclinical safety data**

There is evidence from animal studies that high doses of ceftriaxone calcium salt led to formation of concrements and precipitates in the gallbladder of dogs and monkeys, which proved to be reversible. Animal studies produced no evidence of toxicity to reproduction and genotoxicity. Carcinogenicity studies on ceftriaxone were not conducted.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Not applicable

### **6.2 Incompatibilities**

Based on literature reports, ceftriaxone is not compatible with amsacrine, vancomycin, fluconazole and aminoglycosides and labetalol.

Solutions containing ceftriaxone should not be mixed with or added to other agents except those mentioned in section 6.6

In particular, diluents containing calcium, (e.g. Ringer's solution, Hartmann's solution) should not be used to reconstitute ceftriaxone vials or to further dilute a reconstituted vial for IV administration because a precipitate can form. Ceftriaxone must not be mixed or administered simultaneously with calcium containing solutions including total parenteral nutrition (see section 4.2, 4.3, 4.4 and 4.8).

If treatment with a combination of another antibiotic with Ceftriaxone is intended, administration should not occur in the same syringe or in the same infusion solution.

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

### **6.3 Shelf life**

36 months

### **6.4 Special precautions for storage**

Do not store above 30°C. Protect from light. Keep out of reach of children.

### **6.5 Nature and contents of container**

1200 mg powder filled in glass vial with chlorobutyl rubber stopper sealed with Aluminium flip



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

#### **MODULE-1 ADMINISTRATIVE INFORMATION**

off seal with one ampoule of 10 ml sterile water for injection packed in monocarton along with package leaflet.

#### **6.6 Special precautions for disposal and other handling**

Concentrations for the intravenous injection: 100 mg/ml

Concentrations for the intravenous infusion: 50 mg/ml

(Please refer to section 4.2 for further information).

#### **Reconstitution table Water for Injection BP (Intravenous Injection):**

<b>Vial size</b>	<b>Volume of Diluent to be added</b>
1g	10mL

#### **Reconstitution table 1% Lidocaine Injection BP (Intramuscular Injection):**

<b>Vial size</b>	<b>Volume of Diluent to be added</b>
1g	3.5mL

The use of freshly prepared solutions is recommended. For storage conditions of the reconstituted medicinal product, see section 6.3.

Ceftriazone should not be mixed in the same syringe with any drug other than 1% Lidocaine Injection BP (for intramuscular injection only).

The reconstituted solution should be clear. Do not use if particles are present.

Ceftriazone sodium when dissolved in Water for Injections Ph Eur forms a pale yellow to amber solution. Variations in the intensity of colour of the freshly prepared solutions do not indicate a change in potency or safety.

For single use only. Discard any unused contents.



<b>BRAND NAME:</b>	<b>NORACEF</b>
<b>GENERIC NAME:</b>	<b>CEFTRIAZONE FOR INJECTION USP 1 GM</b>

## **MODULE-1 ADMINISTRATIVE INFORMATION**

### **7. Marketing authorisation holder**

**MEDNORAL PHARMACEUTICAL LTD.**

ALLIANCE BIOTECH, 440/3, VILL. KATHA, BADDI, DISTT. SOLAN (H.P)-173205, INDIA

3, OLAIDE BENSON STREET,  
MARYLAND LAGOS. NIGERIA.

### **8. Marketing authorisation number(s)**

### **9. Date of first authorisation/renewal of the authorisation**

### **10. Date of revision of the text**