

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA****1.3.1 Summary of Product Characteristics (SmPC)****1. Name of Medicinal Product****BALMAL 20/120 TABLETS****ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****2. Qualitative and Quantitative Composition****2.1. Qualitative declaration:****Composition of the Drug product:**

Each Uncoated Tablet Contains

Artemether..... 20 mg

Lumefantrine USP..... 120 mg

Excipients..... Q.S

Qualitative & Quantitative Composition Formula:**Batch Size: 10,00,000 Tablets**

Sr. no.	Ingredients	Spec.	Quantity/ tab (mg)	Overages (%)	Quantity /tab with Overages (mg)	Quantity per Batch (kg)	Reasonfor inclusion
DRY MIXING							
1	Artemether	IHS	20.00	---	20.00	20.00	Active
2	Lumefantrine	USP	120.00	---	120.00	120.00	Active
3	Micro Crystalline Cellulose	BP	25.00	--	25.00	25.00	Diluent
4	Maize Starch	BP	55.00	---	55.00	55.00	Diluent
5	Dicalcium Phosphate	BP	30.00	---	30.00	30.00	Diluent
PASTE PREPARATION							
6	Maize Starch	BP	14.00	---	14.00	14.00	Binder
7	Methyl Paraben	BP	0.23	--	0.23	0.23	Preservative
8	Propyl Paraben	BP	0.04	---	0.04	0.04	Preservative
9	P.V.P.K -30	BP	2.00	---	2.00	2.00	Binder
10	Purified Water	BP	--	---	--	--	Solvent
LUBRICATION							

AR LIFESCIENCES**819/B, RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST.GANDHINAGAR, GUJARAT, INDIA.**

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

11	Purified Talcum	BP	14.00	---	14.00	14.00	Glidant
12	Magnesium stearate	BP	8.75	---	8.75	8.75	Lubricant
13	Colloidal Silicon Dioxide	BP	3.00	---	3.00	3.00	Adsorbant
14	Sodium Starch Glycolate	BP	8.00	---	8.00	8.00	Super Disintegrant
Total weight of tablets					300.00 mg	300.0 kg	

3. Pharmaceutical form

Uncoated tablets

Yellow colour, round shaped, uncoated, biconvex tablets Plain on both sides.

4. Clinical particulars**4.1 Therapeutic indications**

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is indicated for the treatment of uncomplicated cases of malaria due to Plasmodium falciparum in adults and children of 35 kg and above.

The most recent official guidelines on the appropriate use of antimalarial agents and local information on the prevalence of resistance to antimalarial drugs must be taken into consideration for deciding on the appropriateness of therapy with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG

4.2 Posology and method of administration

Oral use

Treatment should be administered at the time of initial diagnosis or at the onset of symptoms. It is preferable that the patient has a positive diagnostic test before administration.

One tablet should be taken twice a day for three days (total six doses).The first dose should be followed by a second dose after 8 hours. The following two days the doses of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should be given twice daily, morning and evening (i.e. 12 hours apart).

AR LIFESCIENCES**819/B,RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST.GANDHINAGAR, GUJARAT, INDIA.**

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

To increase absorption, ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should be taken with food or a milky drink (see section 5.2). If a patient is unable to tolerate food, ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should still be administered, but the systemic exposure may be reduced.

Patients who vomit within 1 hour of taking the medication should repeat the dose.

If a dose is missed, it should be taken as soon as realized and then the recommended regimen continued until the full course of treatment has been completed.

Renal or hepatic impairment

No dose adjustments are necessary in patients with renal or hepatic impairment. However, caution is advised when administering ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG to patients with severe renal or hepatic problems (see section 4.4).

Paediatric patients weighing less than 35 kg:

Appropriate dose adjustments cannot be achieved with this product. Other formulations containing lower amounts of artemether/lumefantrine are available for these patients.

Elderly

No special precautions or dosage adjustments are necessary in such patients.

4.3 Contraindications

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is contraindicated in:

- patients with known hypersensitivity to the active substances or to any of the excipients listed in section 6.1.
 - patients with severe malaria according to WHO definition*.
 - Patients who are taking any drug which is metabolised by the cytochrome enzyme CYP2D6 (e.g. metoprolol, imipramine, amitriptyline, clomipramine).
 - patients with a family history of sudden death or of congenital prolongation of the QTc interval on electrocardiograms, or with any other clinical condition known to prolong the QTc interval.
 - patients taking drugs that are known to prolong the QTc interval (proarrhythmic). These drugs include:
 - antiarrhythmics of classes IA and III,
 - neuroleptics, antidepressive agents,
 - certain antibiotics including some agents of the following classes: macrolides, fluoroquinolones, imidazole and triazole antifungal agents,
 - certain non-sedating antihistamines (terfenadine, astemizole),
 - cisapride.
- flecainide
-

BALMAL 20/120 TABLETS

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG

INDIA

- patients with a history of symptomatic cardiac arrhythmias or with clinically relevant bradycardia or with congestive cardiac failure accompanied by reduced left ventricle ejection fraction.
- Patients with disturbances of electrolyte balance e.g. hypokalemia or hypomagnesemia.
- patients taking drugs that are strong inducers of CYP3A4 such as rifampin, carbamazepine, phenytoin, St. John's wort (*Hypericum perforatum*).

(*Presence of one or more of the following clinical or laboratory features:

Clinical manifestation: Prostration; impaired consciousness or unarousable coma; failure to feed; deep breathing, respiratory distress (acidotic breathing); multiple convulsions; circulatory collapse or shock; pulmonary edema (radiological); abnormal bleeding; clinical jaundice; hemoglobinuria
Laboratory test: Severe normocytic anemia; hemoglobuniuria; hypoglycemia; metabolic acidosis; renal impairment; hyperlactatemia; hyperparasitemia)

4.4 Special warnings and precautions for use

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is not recommended during the first trimester of pregnancy in situations where other suitable and effective antimalarials are available (see section 4.6).

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG has not been evaluated for the treatment of severe malaria, including cases of cerebral malaria or other severe manifestations such as pulmonary oedema or renal failure.

Due to limited data on safety and efficacy, ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should not be given concurrently with any other antimalarial agent (see section 4.5) unless there is no other treatment option.

If a patient deteriorates whilst taking ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG, alternative treatment for malaria should be started without delay. In such cases, monitoring of the ECG is recommended and steps should be taken to correct any electrolyte disturbances.

The long elimination half-life of lumefantrine must be taken into account when administering quinine in patients previously treated with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG.

If quinine is given after ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG, close monitoring of the ECG is advised (see section 4.5).

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

If ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is given after mefloquine, close monitoring of food intake is advised (see section 4.5).

In patients previously treated with halofantrine, ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should not be administered earlier than one month after the last halofantrine dose.

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is not indicated and has not been evaluated for prophylaxis of malaria.

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should be used cautiously in patients on anti-retroviral drugs (ARTs) since decreased artemether, DHA, and/or lumefantrine concentrations may result in a decrease of antimalarial efficacy of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG, (see section 4.5).

Like other antimalarials (e.g. halofantrine, quinine and quinidine) ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG has the potential to cause QT prolongation (see section 5.1).

Caution is recommended when combining ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG with drugs exhibiting variable patterns of inhibition, moderate induction or competition for CYP3A4 as the therapeutic effects of some drugs could be altered. Drugs that have a mixed inhibitory/induction effect on CYP3A4, especially anti-retroviral drugs such as HIV protease inhibitors and non-nucleoside reverse transcriptase inhibitors should be used with caution in patients taking ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG (see sections 4.5 and 5.2).

Caution is recommended when combining ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG with hormonal contraceptives. ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG may reduce the effectiveness of hormonal contraceptives. Therefore, patients using oral, transdermal patch, or other systemic hormonal contraceptives should be advised to use an additional non-hormonal method of birth control for about one month (see sections 4.5).

Patients who remain averse to food during treatment should be closely monitored as the risk of recrudescence may be greater.

Renal impairment

No specific studies have been carried out in this group of patients. There is no significant renal excretion of lumefantrine, artemether and dihydroartemisinin in studies conducted in healthy

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

volunteers and clinical experience is limited. No dose adjustment for the use of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG in patients with renal impairment is recommended. Caution is advised when administering ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG to patients with severe renal impairment. In these patients, ECG and blood potassium monitoring is advised.

Hepatic impairment

No specific studies have been carried out in this group of patients. In patients with severe hepatic impairment, a clinically relevant increase of exposure to artemether and lumefantrine and/or their metabolites cannot be ruled out. Therefore caution should be exercised in dosing patients with severe hepatic impairment (see section 5.2). In these patients, ECG and blood potassium monitoring is advised. No dose adjustment is recommended for patients with mild to moderate hepatic impairment.

New infections

Data for a limited number of patients in a malaria endemic area show that new infections can be treated with a second course of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG. In the absence of carcinogenicity study data, and due to lack of clinical experience, more than two courses of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG cannot be recommended.

Excipient with known effect:

This medicine contains less than 1 mmol sodium (23 mg) per tablet, i.e. is essentially “sodium-free.”

4.5 Interaction with other medicinal products and other forms of interactionInteraction with drugs that are known to prolong the QTc interval

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is contraindicated with concomitant use of drugs (they may cause prolonged QTc interval and Torsade de Pointes) such as: antiarrhythmics of classes IA and III, neuroleptics and antidepressant agents, certain antibiotics including some agents of the following classes: macrolides, fluoroquinolones, imidazole, and triazole antifungal agents, certain non-sedating antihistamines (terfenadine, astemizole), cisapride, flecainide (see section 4.3)

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**Interaction with drugs metabolized by CYP2D6

Lumefantrine was found to inhibit CYP2D6 in vitro. This may be of particular clinical relevance for compounds with a low therapeutic index. Co-administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG with drugs that are metabolised by this iso-enzyme is contraindicated (e.g. neuroleptics, metoprolol, and tricyclic antidepressants such as imipramine, amitriptyline, clomipramine) is contraindicated (see sections 4.3 and 5.2).

Interaction with strong inducers of CYP3A4 such as rifampin

Oral administration of rifampin (600 mg daily), a strong CYP3A4 inducer, with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG Tablets (6-dose regimen over 3 days) in six HIV-1 and tuberculosis coinfecting adults without malaria resulted in significant decreases in exposure to artemether (89%), DHA (85%) and lumefantrine (68%) when compared to exposure values after ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG alone. Concomitant use of strong inducers of CYP3A4 such as rifampin, carbamazepine, phenytoin, St. John's Wort is contraindicated with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG (see section 4.3).

Inducers should not be administered at least one month after ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG administration, unless critical to use as judged by the prescriber.

Concomitant use not recommendedInteraction with other antimalarial drugs (see section 4.4)

Data on safety and efficacy are limited, and ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should therefore not be given concurrently with other antimalarials unless there is no other treatment option (see section 4.4).

If ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is given following administration of mefloquine or quinine, close monitoring of food intake (for mefloquine) or of the ECG (for quinine) is advised. The long elimination half-life of lumefantrine must be taken into account when administering quinine in patients previously treated with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG. In patients previously treated with halofantrine, ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should not be administered earlier than one month after the last halofantrine dose (see section 4.4).

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**Mefloquine

A drug interaction study with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG in man involved administration of a 6-dose regimen over 60 hours in healthy volunteers which was commenced at 12 hours after completion of a 3-dose regimen of mefloquine or placebo. Plasma mefloquine concentrations from the time of addition of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG were not affected compared with a group which received mefloquine followed by placebo.

Pre-treatment with mefloquine had no effect on plasma concentrations of artemether or the artemether/dihydroartemisinin ratio but there was a significant reduction in plasma levels of lumefantrine, possibly due to lower absorption secondary to a mefloquine-induced decrease in bile production. Patients should be encouraged to eat at dosing times to compensate for the decrease in bioavailability.

Quinine

A drug interaction study in healthy male volunteers showed that the plasma concentrations of lumefantrine and quinine were not affected when i.v. quinine (10 mg/kg BW over 2 hours) was given sequentially 2 hours after the last (sixth) dose of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG (so as to produce concurrent plasma peak levels of lumefantrine and quinine). Plasma concentrations of artemether and dihydroartemisinin (DHA) appeared to be lower. In this study, administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG to 14 subjects had no effect on QTc interval. Infusion of quinine alone in 14 other subjects caused a transient prolongation of QTc interval, which was consistent with the known cardiotoxicity of quinine. This effect was slightly, but significantly, greater when quinine was infused after ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG in 14 additional subjects. It would thus appear that the inherent risk of QTc prolongation associated with i.v. quinine was enhanced by prior administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG.

Concomitant use requiring caution**Interactions affecting the use of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG**Interaction with CYP3A4 inhibitors

AR LIFESCIENCES**819/B, RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST. GANDHINAGAR, GUJARAT, INDIA.**

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

Both artemether and lumefantrine are metabolised predominantly by the cytochrome enzyme CYP3A4, but do not inhibit this enzyme at therapeutic concentrations.

Ketoconazole

The concurrent oral administration of ketoconazole with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG led to a modest increase (≤ 2 -fold) in artemether, DHA, and lumefantrine exposure in healthy adult subjects. This increase in exposure to the antimalarial combination was not associated with increased side effects or changes in electrocardiographic parameters. Based on this study, dose adjustment of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is considered unnecessary in falciparum malaria patients when administered in association with ketoconazole or other potent CYP3A4 inhibitors.

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should be used cautiously with drugs that inhibit CYP3A4 and are contraindicated with drugs which additionally are known to prolong QTc (see Section 4.3 Contraindications), due to potential for increased concentrations of lumefantrine which could lead to QT prolongation.

Interaction with weak to moderate inducers of CYP3A4

When ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is co-administered with moderate inducers of CYP3A4, it may result in decreased concentrations of artemether and/or lumefantrine and loss of antimalarial efficacy (see section 4.4).

Interaction with anti-retroviral drugs such as protease inhibitors and non-nucleoside reverse transcriptase inhibitors

Both artemether and lumefantrine are metabolized by CYP3A4. ARTs, such as protease inhibitors and non- nucleoside reverse transcriptase inhibitors, are known to have variable patterns of inhibition, induction or competition for CYP3A4. ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should be used cautiously in patients on ARTs since decreased artemether, DHA, and/or lumefantrine concentrations may result in a decrease of antimalarial efficacy of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG, and increased lumefantrine concentrations may cause QT prolongation (see Section 4.4).

Lopinavir/ ritonavir

In a clinical study in healthy volunteers, lopinavir/ritonavir decreased the systemic exposures to artemether and DHA by approximately 40% but increased the exposure to lumefantrine by

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

approximately 2.3- fold. Exposures to lopinavir/ritonavir were not significantly affected by concomitant use of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG.

Nevirapine

In a clinical study in HIV-infected adults, nevirapine significantly reduced the median C_{max} and AUC of artemether by approximately 61% and 72%, respectively and reduced the median C_{max} and AUC of dihydroartemisinin by approximately 45% and 37%, respectively. Lumefantrine C_{max} and AUC were non-significantly reduced by nevirapine. Artemether/lumefantrine reduced the median C_{max} and AUC of nevirapine by approximately 43% and 46% respectively.

Efavirenz

Efavirenz decreased the exposures to artemether, DHA, and lumefantrine by approximately 50%, 45%, and 20%, respectively. Exposures to efavirenz were not significantly affected by concomitant use of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG.

Interactions resulting in effects of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG on other drugsInteraction with drugs metabolized by CYP450 enzymes

When ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is co-administered with substrates of CYP3A4 it may result in decreased concentrations of the substrate and potential loss of substrate efficacy. Studies in humans have demonstrated that artemisinins have some capacity to induce CYP3A4 and CYP2C19 and inhibit CYP2D6 and CYP1A2. Although the magnitude of the changes was generally low it is possible that these effects could alter the therapeutic response of drugs that are predominantly metabolised by these enzymes (see sections 4.4 and 5.2).

Interaction with hormonal contraceptives

In vitro, the metabolism of ethinyl estradiol and levonorgestrel was not induced by artemether, DHA, or lumefantrine. However, artemether has been reported to weakly induce, in humans, the activity of CYP2C19, CYP2B6, and CYP3A. Therefore, ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG may potentially reduce the effectiveness of hormonal contraceptives. Patients using oral, transdermal patch, or other systemic hormonal contraceptives should be advised to use an additional nonhormonal method of birth control for about one month (see sections 4.4 and 4.6).

Drug-food/drink interactions

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should be taken with food or drinks rich in fat such as milk as the absorption of both artemether and lumefantrine is increased (see Section 4.2).

Grapefruit juice should be used cautiously during ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG treatment. Administration of artemether with grapefruit juice in healthy adult subjects resulted in an approximately two fold increase in systemic exposure to the parent drug.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential

Women using oral, transdermal patch, or other systemic hormonal contraceptives should be advised to use an additional non-hormonal method of birth control for about one month (see section 4.4).

Pregnancy

A meta-analysis of observational studies including over 500 artemether- lumefantrine exposed women in their first trimester of pregnancy assessed adverse pregnancy outcomes. The data showed that compared to quinine, artemisinin treatment, including artemether- lumefantrine, was not associated with an increased risk of miscarriage, stillbirth or congenital anomalies. However, due to the limitations of these studies, the risk of adverse pregnancy outcomes for artemether- lumefantrine exposed women in early pregnancy cannot be excluded.

Safety data from pregnancy studies including over 1200 pregnant women who were exposed to artemether- lumefantrine during the second or third trimester did not show an increase in adverse pregnancy outcomes or teratogenic effects over background rates.

Studies in animals have shown reproductive toxicity (see section 5.3).

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG treatment is not recommended during the first trimester of pregnancy in situations where other suitable and effective antimalarials are available (see section 4.4). However, it should not be withheld in life-threatening situations, where no other effective antimalarials are available. During the second and third trimester, ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG treatment should be considered if the expected benefit to the mother outweighs the risk to the foetus.

Breast-feeding

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

Animal data suggest excretion into breast milk but no data are available in humans. Women taking ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should not breast- feed during their treatment. Due to the long elimination half- life of lumefantrine (2 to 6 days), it is recommended that breast- feeding should not resume until at least one week after the last dose of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG unless potential benefits to the mother and child outweigh the risks of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG treatment.

Fertility

There is no information on the effects of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG on human fertility (see section 5.3).

4.7 Effects on ability to drive and use machines

Patients receiving ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG should be warned that dizziness or fatigue/asthenia may occur in which case they should not drive or use machines.

4.8 Undesirable effects

The safety of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG has been evaluated in 20 clinical trials with more than 3500 patients. A total of 1810 adults and adolescents above 12 years of age as well as 1788 infants and children of 12 years of age and below have received ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG in clinical trials.

Adverse reactions reported from clinical studies and post-marketing experience are listed below according to system organ class.

Adverse reactions are ranked under headings of frequency using the MedDRA frequency convention:

Very common ($\geq 1/10$)

Common ($\geq 1/100$ to $< 1/10$)

Uncommon ($\geq 1/1,000$ to $< 1/100$)

AR LIFESCIENCES**819/B, RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST. GANDHINAGAR, GUJARAT, INDIA.**

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**Rare ($\geq 1/10,000$ to $< 1/1,000$)Very rare ($< 1/10,000$)

Not known (cannot be estimated from available data).

Table 1 Frequency of Undesirable effects

	Adults and adolescents above 12 years of age	Infants and children of 12 years of age and below (incidence estimates)
Blood and lymphatic system disorders		
Delayed haemolytic anaemia [#]	Not Known	Not Known
Immune system disorders		
Hypersensitivity	Not known	Rare
Metabolism and nutrition disorders		
Decreased appetite	Very common	Very common (16.8 %)
Psychiatric disorders		
Sleep disorders	Very common	Common (6.4 %)
Insomnia	Common	Uncommon
Nervous system disorders		
Headache	Very common	Very common (17.1 %)
Dizziness	Very common	Common (5.5 %)
Paraesthesia	Common	--
Ataxia, hypoaesthesia	Uncommon	--
Somnolence	Uncommon	Uncommon
Clonus	Common	Uncommon
Cardiac disorders		
Palpitations	Very common	Common (1.8 %)
Electrocardiogram QT prolonged	Common	Common (5.3 %)

AR LIFESCIENCES**819/B, RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST. GANDHINAGAR, GUJARAT, INDIA.**

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA****Respiratory, thoracic and mediastinal disorders**

Cough	Common	Very common (22.7 %)
-------	--------	----------------------

Gastrointestinal disorders

Vomiting	Very common	Very common (20.2 %)
----------	-------------	----------------------

Abdominal pain	Very common	Very common (12.1 %)
----------------	-------------	----------------------

Nausea	Very common	Common (6.5 %)
--------	-------------	----------------

Diarrhoea	Common	Common (8.4 %)
-----------	--------	----------------

Hepatobiliary disorders

Liver function tests increased	Uncommon	Common (4.1 %)
--------------------------------	----------	----------------

Skin and subcutaneous tissue disorders

Rash	Common	Common (2.7 %)
------	--------	----------------

Pruritus	Common	Uncommon
----------	--------	----------

Urticaria	Uncommon	Uncommon
-----------	----------	----------

Angioedema*	Not known	Not known
-------------	-----------	-----------

Musculoskeletal and connective tissue disorders

Arthralgia	Very common	Common (2.1 %)
------------	-------------	----------------

Myalgia	Very common	Common (2.2 %)
---------	-------------	----------------

General disorders and administration site conditions

Asthenia	Very common	Common (5.2 %)
----------	-------------	----------------

Fatigue	Very common	Common (9.2 %)
---------	-------------	----------------

Gait disturbance	Common	--
------------------	--------	----

*: These adverse reactions were reported during post- marketing experience. Because these spontaneously reported events are from a population of uncertain size, it is difficult to estimate their frequency.

#: Has been reported up to a few weeks after treatment has been stopped.

AR LIFESCIENCES**819/B, RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST.GANDHINAGAR, GUJARAT, INDIA.**

4.9 Overdose

In cases of suspected overdosage symptomatic and supportive therapy should be given as appropriate, which should include ECG and blood potassium monitoring.

5. Pharmacological properties**5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: antimalarials, blood schizontocide, ATC code: P01 BF01.

Pharmacodynamic effects

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG comprises a fixed ratio of 1:6 parts of artemether and lumefantrine, respectively. The site of antiparasitic action of both components is the food vacuole of the malarial parasite, where they are thought to interfere with the conversion of haem, a toxic intermediate produced during haemoglobin breakdown, to the nontoxic haemozoin, malaria pigment. Lumefantrine is thought to interfere with the polymerisation process, while artemether generates reactive metabolites as a result of the interaction between its peroxide bridge and haem iron. Both artemether and lumefantrine have a secondary action involving inhibition of nucleic acid- and protein synthesis within the malarial parasite. ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG has been reported to have potent activity in terms of clearing gametocytes.

By 2015, resistance to artemisinins emerged in Southeast Asia. Studies with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG in this region showed delayed parasite clearance (manifested as a higher proportion of patients with parasitemia on Day 3 after initiation of treatment), although overall efficacy as measured by cure rates after 28 days, remained high (WHO 2014). In Africa, only isolated reports on delayed parasite clearance are available and a clear trend towards resistance development was not observed.

Treatment of Acute Uncomplicated P. falciparum Malaria

The efficacy of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG Tablets was evaluated for the treatment of acute, uncomplicated malaria (defined as symptomatic P. falciparum malaria without signs and symptoms of severe malaria or evidence of vital organ dysfunction) in five 6-dose regimen studies and one study comparing the 6-dose regimen with the 4-dose regimen. Baseline parasite density ranged from 500/µl - 200,000/µl (0.01% to 4% parasitemia) in the majority of patients. Studies were conducted in otherwise healthy, partially immune or non-

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

immune adults and children (≥ 5 kg body weight) with uncomplicated malaria in Thailand, sub-Saharan Africa, Europe, and South America.

Efficacy endpoints consisted of:

- 28-day cure rate, proportion of patients with clearance of asexual parasites within 7 days without recrudescence by day 28
- parasite clearance time (PCT), defined as time from first dose until first total and continued disappearance of asexual parasite which continues for a further 48 hours
- fever clearance time (FCT), defined as time from first dose until the first time body temperature fell below 37.5°C and remained below 37.5°C for at least a further 48 hours (only for patients with temperature $>37.5^{\circ}\text{C}$ at baseline)

The modified intent to treat (mITT) population includes all patients with malaria diagnosis confirmation who received at least one dose of study drug. Evaluable patients generally are all patients who had a day 7 and a day 28 parasitological assessment or experienced treatment failure by day 28. The results are presented in the table below:

Table 2 Clinical efficacy results

Study No.	Age	Polymerase chain reaction (PCR)-corrected 28-day cure rate ¹ n/N (%) in evaluable patients	Median FCT ² [25 th , 75 th percentile]	Median PCT ² [25 th , 75 th percentile]	Year/ Study location
A025 ⁴	3-62 years	93/96 (96.9)	n ³ =59 35 hours [20, 46]	n=118 44 hours [22, 47]	1996-97 Thailand
A026	2-63 years	130/133 (97.7)	n ³ =87 22 hours [19, 44]	NA	1997-98 Thailand
A028	12-71 years	148/154 (96.1)	n ³ =76 29 hours [8, 51]	n=164 29 hours [18, 40]	1998-99 Thailand
A2401	16-66 years	119/124 (96.0)	n ³ =100 37 hours [18, 44]	n=162 42 hours [34, 63]	2001-05 Europe, Columbia

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

A2403	2 months-9 years	289/299 (96.7)	n ³ =309 8 hours [8, 24]	n=310 24 hours [24, 36]	2002-03 3 countries in Africa
B2303 ^{CT}	3 months-12 years	403/419 (96.2)	n ³ =323 8 hours [8, 23]	n=452 35 hours [24, 36]	2006-07 5 countries in Africa
B2303 ^{DT}	3 months-12 years	394/416 (94.7)	n ³ =311 8 hours [8, 24]	n=446 34 hours [24, 36]	2006-07 5 countries in Africa

¹ Efficacy cure rate based on blood smear microscopy

² mITT population

³ For patients who had a body temperature >37.5°C at baseline only

⁴ Only the 6-dose regimen over 60 hours group data is presented

^{CT} –ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG tablets administered as crushed tablets

^{DT} –ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG Dispersible tablets

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is not indicated for, and has not been evaluated in, the treatment of malaria due to *P. vivax*, *P. malariae* or *P. ovale*, although some patients in clinical studies had co-infection with *P. falciparum* and *P. vivax* at baseline. In 319 adult patients in whom gametocytes were present, the median time to gametocyte clearance with ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG was 96 hours. ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is active against blood stages of *Plasmodium vivax*, but is not active against hypnozoites.

Paediatric population

Three studies have been conducted

Study A2403 was conducted in Africa in 310 infants and children aged 2 months to 9 years, weighing 5 kg to 25 kg, with an axillary temperature $\geq 37.5^{\circ}\text{C}$. Results of 28-day cure rate (PCR-corrected), median parasite clearance time (PCT), and fever clearance time (FCT) are reported in table 3 below.

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

Study B2303 was conducted in Africa in 452 infants and children, aged 3 months to 12 years, weighing 5 kg to <35 kg, with fever ($\geq 37.5^{\circ}\text{C}$ axillary or $\geq 38^{\circ}\text{C}$ rectally) or history of fever in the preceding 24 hours. This study compared crushed tablets and dispersible tablets. Results of 28-day cure rate (PCR-corrected), median parasite clearance time (PCT), and fever clearance time (FCT) for crushed tablets are reported in table 3 below.

Table 3 Clinical efficacy by weight for pediatric studies

Study No. Weight category	Median PCT ¹ [25 th , 75 th percentile]	PCR-corrected 28-day cure rate ² n/N (%) in evaluable patients
Study A2403	24 hours [24, 36]	145/149 (97.3)
5 - <10 kg	35 hours [24, 36]	103/107 (96.3)
10 - <15 kg	24 hours [24, 36]	41/43 (95.3)
15 -25 kg		
Study B2303 ^{CT}	36 hours [24, 36]	65/69 (94.2)
5 - <10 kg	35 hours [24, 36]	174/179 (97.2)
10 - <15 kg	35 hours [24, 36]	134/140 (95.7)
15 -<25 kg	26 hours [24, 36]	30/31 (96.8)
25-35 kg		

¹ mITT population² Efficacy cure rate based on blood smear microscopy^{CT} ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG tablets administered as crushed tablets

Study B2306, was a multi-centre, open- label, single-arm study conducted in 20 infants in Africa, Benin and Burkina Faso to evaluate the efficacy, safety and pharmacokinetics of dispersible tablets in infants aged >28 days and <5 kg of body weight, who were treated with one dispersible tablet (20 mg artemether/120 mg lumefantrine) given twice-daily for three days and followed up for six weeks (core follow-up) and at the age of 12 months (long-term follow-up).

Dispersible tablets were well tolerated with reported adverse events of mild to moderate severity. In the per protocol population, PCR-corrected cure rate at days 28 and 42 was 100% (95% CI: 79.4, 100). For important exposure results, see section 5.2. Although neurotoxicity was not

AR LIFESCIENCES**819/B, RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST. GANDHINAGAR, GUJARAT, INDIA.**

observed in the patients in Study B2306, artemether has been associated with neurotoxicity in studies in rats and dogs, see section 5.3.

QT/QTc Prolongation:

Adults and children with malaria

For information on the risk of QT/QTc prolongation in patients see section 4.4

Healthy adults

In a healthy adult volunteer parallel group study including a placebo and moxifloxacin control group (n=42 per group), the administration of the six dose regimen of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG was associated with prolongation of QTcF. The mean changes from baseline at 68, 72, 96, and 108 hours post first dose were 7.45, 7.29, 6.12 and 6.84 msec, respectively. At 156 and 168 hours after first dose, the changes from baseline for QTcF had no difference from zero. No subject had a >30 msec increase from baseline nor an absolute increase to >500 msec. Moxifloxacin control was associated with a QTcF increase as compared to placebo for 12 hours after the single dose with a maximal change at 1 hour after dose of 14.1 msec. In the adult/adolescent population included in clinical trials, 8 patients (0.8%) receiving ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG experienced a QTcB >500 msec and 3 patients (0.4%) a QTcF >500 msec. Prolongation of QTcF interval >30 msec was observed in 36% of patients.

In clinical trials conducted in children with the 6-dose regimen, no patient had post-baseline QTcF >500 msec whereas 29.4% had QTcF increase from baseline >30 msec and 5.1% >60 msec. In clinical trials conducted in adults and adolescents with the 6-dose regimen, post-baseline QTcF prolongation of >500 msec was reported in 0.2% of patients, whereas QTcF increase from baseline >30 msec was reported in 33.9% and >60 msec in 6.2% of patients.

In the infant/children population included in clinical trials, 3 patients (0.2%) experienced a QTcB >500 msec. No patient had QTcF >500 msec. Prolongation of QTcF intervals >30 msec was observed in 34% of children weighing 5-10 kg, 31% of children weighing 10-15 kg and 24% of children weighing 15-25 kg, and 32% of children weighing 25-35 kg.

5.2 Pharmacokinetic properties

Pharmacokinetic characterisation of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG is limited by the lack of an intravenous formulation, and the very high inter-and

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

intra-subject variability of artemether and lumefantrine plasma concentrations and derived pharmacokinetic parameters (AUC, C_{max}).

Absorption

Artemether is absorbed fairly rapidly and dihydroartemisinin, the active metabolite of artemether, appears rapidly in the systemic circulation with peak plasma concentrations of both compounds reached about 2 hours after dosing. Mean C_{max} and AUC values of artemether ranged between 60.0–104 ng/ml and 146–338 ng·h/ml, respectively, in fed healthy adults after a single dose of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG, 80 mg artemether/480 mg lumefantrine. Mean C_{max} and AUC values of dihydroartemisinin ranged between 49.7–104 ng/mL and 169–308 ng·h/ml, respectively. Absorption of lumefantrine, a highly lipophilic compound, starts after a lag-time of up to 2 hours, with peak plasma concentration (mean between 5.10–9.80 µg/ml) about 6–8 hours after dosing. Mean AUC values of lumefantrine ranged between 108 and 243 µg·h/ml. Food enhances the absorption of both artemether and lumefantrine: in healthy volunteers the relative bioavailability of artemether was increased more than two- fold, and that of lumefantrine sixteen-fold compared with fasted conditions when ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG was taken after a high-fat meal.

Food has also been shown to increase the absorption of lumefantrine in patients with malaria, although to a lesser extent (approximately two- fold), most probably due to the lower fat content of the food ingested by acutely ill patients. The food interaction data indicate that absorption of lumefantrine under fasted conditions is very poor (assuming 100% absorption after a high- fat meal, the amount absorbed under fasted conditions would be <10% of the dose). Patients should therefore be encouraged to take the medication with a normal diet as soon as food can be tolerated.

Distribution

Artemether and lumefantrine are both highly bound to human serum proteins *in vitro* (95.4% and 99.7%, respectively). Dihydroartemisinin is also bound to human serum proteins (47–76%).

Biotransformation

Artemether is rapidly and extensively metabolised (substantial first-pass metabolism) both *in vitro* and in humans. Human liver microsomes metabolise artemether to the biologically active main metabolite dihydroartemisinin (demethylation), predominantly through the isoenzyme CYP3A4/5. This metabolite has also been detected in humans *in vivo*.

Glucuronidation of dihydroartemisinin is predominately catalysed by UGT1A9 and UGT2B7.

Dihydroartemisinin is further converted to inactive metabolites.

The pharmacokinetics of artemether in adults is time-dependent. During repeated administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG, plasma artemether levels decreased significantly, while levels of the active metabolite (dihydroartemisinin) increased, although not to a statistically significant degree. The ratio of day 3/day 1 AUC for artemether was between 0.19 and 0.44, and was between 1.06 and 2.50 for dihydroartemisinin. This suggests that there was induction of the enzyme responsible for the metabolism of artemether. Artemether and dihydroartemisinin were reported to have a mild inducing effect on CYP3A4 activity. The clinical evidence of induction is consistent with the *in vitro* data described in section 4.5

Lumefantrine is N-debutylated, mainly by CYP3A4, in human liver microsomes. *In vivo* in animals (dogs and rats), glucuronidation of lumefantrine takes place directly and after oxidative biotransformation. In humans, the exposure to lumefantrine increases with repeated administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG over the 3-day treatment period, consistent with the slow elimination of the compound (see section 5.2 Elimination). Systemic exposure to the metabolite desbutyl- lumefantrine, for which the *in vitro* antiparasitic effect is 5 to 8 fold higher than that for lumefantrine, was less than 1% of the exposure to the parent drug. Desbutyl- lumefantrine data is not available specifically for an African population. *In vitro*, lumefantrine significantly inhibits the activity of CYP2D6 at therapeutic plasma concentrations (see sections 4.3 and 4.5).

Elimination

Artemether and dihydroartemisinin are rapidly cleared from plasma with a terminal half-life of about 2 hours. Lumefantrine is eliminated very slowly with an elimination half-life of 2 to 6 days. Demographic characteristics such as sex and weight appear to have no clinically relevant effects on the pharmacokinetics of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG. Limited urinary excretion data are available for humans. In 16 healthy volunteers, neither lumefantrine nor artemether was found in urine after administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG, and only traces of dihydroartemisinin were detected (urinary excretion of dihydroartemisinin amounted to less than 0.01% of the artemether dose).

In animals (rats and dogs), no unchanged artemether was detected in faeces and urine due to its rapid and extensive first-pass metabolism, but numerous metabolites (partly identified) have been

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

detected in faeces, bile and urine. Lumefantrine was excreted unchanged in faeces and with traces only in urine. Metabolites of lumefantrine were eliminated in bile/faeces.

Dose proportionality

No specific dose proportionality studies were performed. Limited data suggest a dose-proportional increase of systemic exposure to lumefantrine when doubling the ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG dose. No conclusive data is available for artemether.

Bioavailability/bioequivalence studies

Systemic exposure to lumefantrine, artemether and dihydroartemisinin was similar following administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG as dispersible tablets and crushed tablets in healthy adults.

Systemic exposure to lumefantrine was similar following administration of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG dispersible tablets and intact tablets in healthy adults. However, exposure to artemether and dihydroartemisinin was significantly lower (by 20-35%) for the dispersible than for the intact tablet. These findings are not considered to be clinically relevant for the use of the dispersible tablets in the paediatric population since adequate efficacy of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG dispersible tablets was demonstrated in this population. The dispersible tablet is not recommended for use in adults.

Older people

No specific pharmacokinetic studies have been performed in elderly patients. However, there is no information suggesting that the dosage in patients over 65 years of age should be different than in younger adults.

Paediatric population

In paediatric malaria patients, mean C_{max} (CV%) of artemether (observed after first dose of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG) were 223 (139%), 198 (90%) and 174 ng/ml (83%) for body weight groups 5-<15, 15-<25 and 25-<35 kg, respectively, compared to 186 ng/ml (67%) in adult malaria patients. The associated mean C_{max} of DHA were 54.7 (108%), 79.8 (101%) and 65.3 ng/mL (36%), respectively compared to 101 ng/ml (57%) in adult malaria patients. AUC of lumefantrine (population mean, covering the six doses of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG) were 577, 699 and 1150

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

$\mu\text{g} \cdot \text{h/ml}$ for paediatric malaria patients in body weight groups 5-<15, 15-<25 and 25-<35 kg, respectively, compared to a mean AUC of 758 $\mu\text{g} \cdot \text{h/ml}$ (87%) in adult malaria patients. The elimination half-lives of artemether and lumefantrine in children are unknown.

Infants weighing <5 kg

Study B2306 (see section 5.1) showed that the C_{max} of artemether and DHA in infants with uncomplicated *P. falciparum* malaria weighing <5 kg and older than 28 days of age who were treated with artemether/lumefantrine dispersible tablets, was on average 2- to 3-fold higher than that in pediatric patients with a body weight ≥ 5 kg and children up to 12 years of age treated with the same dose of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG tablets. The mean C_{max} of lumefantrine was similar to that observed in pediatric patients with a body weight ≥ 5 kg.

Race/Ethnicity

Pharmacokinetics of artemether, DHA and lumefantrine in the Japanese population was found to be consistent with other populations.

Hepatic and Renal impairment

No specific pharmacokinetic studies have been performed either in patients with hepatic or renal insufficiency or elderly patients. The primary clearance mechanism of both artemether and lumefantrine may be affected in patients with hepatic impairment. In patients with severe hepatic impairment, a clinically significant increase of exposure to artemether and lumefantrine and/or their metabolites cannot be ruled out. Therefore, caution should be exercised in dosing patients with severe hepatic impairment. Based on the pharmacokinetic data in 16 healthy subjects showing no or insignificant renal excretion of lumefantrine, artemether and dihydroartemisinin, no dose adjustment for the use of ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG in patients with renal impairment is advised.

5.3 Preclinical safety data

General toxicity

The main changes observed in repeat-dose toxicity studies were associated with the expected pharmacological action on erythrocytes, accompanied by responsive secondary haematopoiesis.

Neurotoxicity

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

Studies in dogs and rats have shown that intramuscular injections of artemether resulted in brain lesions. Changes observed mainly in brainstem nuclei included chromatolysis, eosinophilic cytoplasmic granulation, spheroids, apoptosis and dark neurons. Lesions were observed in rats dosed for at least 7 days and dogs for at least 8 days, but lesions were not observed after shorter intramuscular treatment courses or after oral dosing. The estimated artemether 24 h AUC after 7 days of dosing at the no observed effect level is approximately 7- fold greater or more than the estimated artemether 24 h AUC in adult humans. The hearing threshold was affected at 20 dB by oral artemether administration to dogs at a dose of about 29 times the highest artemether clinical dose (160 mg/day) based on body surface area comparisons. Most nervous system disorder adverse events in the studies of the 6-dose regimen were mild in intensity and resolved by the end of the study.

Mutagenicity

Artemether and lumefantrine were not genotoxic/clastogenic based on *in vitro* and *in vivo* testing.

Carcinogenicity

Carcinogenicity studies were not conducted.

Reproductive toxicity studies

Embryotoxicity was observed in rat and rabbit reproductive toxicity studies conducted with artemether, a derivative of artemisinin. Artemisinins are known to be embryotoxic. Lumefantrine alone caused no sign of reproductive or development toxicity at doses up to 1,000 mg/kg/day in rats and rabbits, doses which are at least 10 times higher than the daily human dose based on body surface area comparisons.

Reproductive toxicity studies performed with the artemether- lumefantrine combination caused maternal toxicity and increased post-implantation loss in rats and rabbits.

Artemether caused increases in post- implantation loss and teratogenicity (characterised as a low incidence of cardiovascular and skeletal malformations) in rats and rabbits. The embryotoxic artemether dose in the rat yields artemether and dihydroartemisinin exposures similar to those achieved in humans based on AUC.

Fertility

Artemether- lumefantrine administration yielded altered sperm motility, abnormal sperm, reduced epididymal sperm count, increased testes weight, and embryotoxicity; other reproductive effects (decreased implants and viable embryos, increased preimplantation loss) were also observed. The

BALMAL 20/120 TABLETS**ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG****INDIA**

no adverse effect level for fertility was 300 mg/kg/day. The relevance to this finding in humans is unknown.

Juvenile toxicity studies

A study investigated the neurotoxicity of oral artemether in juvenile rats. Mortality, clinical signs and reductions in body weight parameters occurred most notably in younger rats. Despite the systemic toxicity noted, there were no effects of artemether on any of the functional tests performed and there was no evidence of a direct neurotoxic effect in juvenile rats.

Very young animals are more sensitive to the toxic effect of artemether than adult animals. There is no difference in sensitivity in slightly older animals compared to adult animals. Clinical studies have established the safety of artemether and lumefantrine administration in patients weighing 5 kg and above.

Cardiovascular Safety Pharmacology

In toxicity studies in dogs at doses ≥ 600 mg/kg/day, there was some evidence of prolongation of the Q Tc interval (safety margin of 1.3- fold to 2.2- fold for artemether using calculated free Cmax), at higher doses than intended for use in man. In vitro hERG assays showed a safety margin of >100 for artemether and dihydroartemisinin. The hERG IC₅₀ was 8.1 μ M for lumefantrine and 5.5 μ M for its desbutyl metabolite.

Based on the available non-clinical data, a potential for QTc prolongation in the human cannot be discounted. For effects in the human see sections 4.3, 4.4 and 5.1.

6. PHARMACEUTICAL PARTICULARS**6.1 List of excipients**

Sr. No.	Ingredients Name	Specification
1	Maize Starch	BP
2	Dicalcium Phosphate	BP
3	Micro Crystalline Cellulose	BP
4	P.V.P.K -30	BP
5	Methyl Paraben	BP
6	Propyl Paraben	BP
7	Purified Water	BP
8	Purified Talcum	BP
9	Magnesium stearate	BP
10	Colloidal Silicon Dioxide	BP
11	Sodium Starch Glycolate	BP

AR LIFESCIENCES**819/B, RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST.GANDHINAGAR, GUJARAT, INDIA.**

BALMAL 20/120 TABLETS

ARTEMETHER & LUMEFANTRINE TABLETS 20MG/120MG

INDIA

6.2 Incompatibilities

None Known

6.3 Shelf life

36 Months

6.4 Special precautions for storage

Store in a cool & dry place, below 30°C. Keep
all medicines out of reach of children.

6.5 Nature and contents of container

Each Alu/PVC blister contains 24 tablets and such 1 blister pack in a monocarton with pack insert and such
30 monocarton are packed in a outercarton.

6.6 Special precautions for disposal and other handling

No special requirements.

7- Marketing Authorization Holder:

GROHE PHARMACEUTICAL LTD

25, Emma Abimbola Cole Street, Lekki

Phase 1, Lagos State, Nigeria.

8- Marketing Authorization Number (s):

Product license / registration Number (s)

9- Manufacturer Name:

AR LIFESCIENCES

819/B, Rakanpur Industrial Area, Tal. Kalol, Dist.Gandhinagar,Gujarat, India

Email:info. systochem@yahoo.com

AR LIFESCIENCES

819/B,RAKANPUR INDUSTRIAL AREA, TAL. KALOL, DIST.GANDHINAGAR, GUJARAT, INDIA.