

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED	
BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

1.3 Product Information

1.3.1 Summary of Product Characteristics (SmPC)

Enclosed.

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

1. Name of the medicinal product

Vadis Metronidazole Suspension

2. Qualitative and quantitative composition

Each 5 ml contains

Metronidazole BP 200 mg

3. Pharmaceutical form

Suspension

4. Clinical particulars

4.1 Therapeutic indications

Metronidazole Oral Suspension is indicated in the prophylaxis and treatment of infections in which anaerobic bacteria have been identified or are suspected as the pathogen.

Metronidazole Oral Suspension is active against a wide range of pathogenic micro-organisms, notably *Trichomonas vaginalis*, *Entamoeba histolytica*, *Giardia lamblia*, *Balantidium coli* and other species of bacteroides, fusobacteria, eubacteria, clostridia, gardnerella vaginalis and anaerobic cocci.

It is indicated in

Adults, Children and Newborns with a gestation age of over 40 weeks for:

- The treatment of septicaemia, bacteraemia, brain abscess, necrotising pneumonia, osteomyelitis, puerperal sepsis, pelvic abscess, pelvic cellulitis, peritonitis and post-operative wound infections from which one or more pathogenic anaerobes have been isolated.
- The prevention of post-operative infections caused by anaerobic bacteria particularly species of bacteroides and anaerobic streptococci.

Adults and Children over 10 years only for:

- Bacterial vaginosis (also known as non-specific vaginitis, anaerobic vaginitis or *Gardnerella* vaginitis).
- Acute dental infections (e.g. acute pericoronitis and acute apical infections).
- Anaerobically infected leg ulcers and pressure sores.

Adults and Children for:

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

- The treatment of urogenital trichomoniasis in the female (trichomonal vaginitis) and in the male.
- All forms of amoebiasis (intestinal and extra-intestinal disease and that of symptomless cyst passers)
- Giardiasis
- Acute ulcerative gingivitis.

Children for

- Eradication of *Helicobacter pylori*

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

A: **Prophylaxis:** against anaerobic infection- chiefly in the context of abdominal (especially colorectal) and gynaecological surgery.

Dosage: 400mg at 8 hourly intervals during the 24 hours preceding the operation followed by postoperative intravenous or rectal administration until the patient is able to take Metronidazole Oral Suspension by mouth.

Children < 12 years: 20 – 30mg/kg as a single dose given 1 – 2 hours before surgery.

Newborns with a gestation age <40 weeks: 10mg/kg body weight as a single dose before operation.

Elderly: Caution is advised in the elderly, particularly at high doses, although there is limited information available on modification of drug.

Anaerobic infections: The duration of a course of Metronidazole treatment is about 7 days but it will depend upon the seriousness of the patient's condition as assessed clinically and bacteriologically.

B: ***Treatment of established anaerobic infection:***

800mg followed by 400mg at 8 hourly intervals.

Children > 8 weeks to 12 years of age: The usual daily dose is 20 – 30mg/kg/day as a single dose or divided into 7.5mg/kg every 8 hours. The daily dose may be increased to 40mg/kg, depending on the severity of the infection. Duration of treatment is usually 7 days.

Children < 8 weeks of age: 15mg/kg as a single dose daily or divided into 7.5mg/kg every 12 hours.

In newborns with a gestation age <40 weeks, accumulation of metronidazole can occur during the first week of life, which is why the concentrations of metronidazole in serum should preferably be monitored after a few days therapy.

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

C: *Treatment of Protozoal and Other Infections:*

(See Table).

	Duration of dosage in days	Adults and children over 10 years**	Children*		
			7-10 years	3-7 years	1-3 years
Urogenital Trichomoniasis Where re-infection is likely, in adults the consort should receive a similar course of treatment concurrently	7	200mg three times daily	40mg/kg orally as a single dose or 15 – 30mg/kg/day divided in 2 – 3 doses not to exceed 2000mg/dose		
	or				
	5 - 7	400mg twice daily			
	or				
	1	2000mg as a single dose			
Bacterial Vaginosis	5 - 7	400mg twice daily			
	or				
	1	2000mg as a single dose			
Amoebiasis	5	800 mg three times daily	400 mg three times daily	200 mg four times daily	200 mg three times daily
(a) Invasive intestinal disease in susceptible subjects					
(b) Intestinal disease in less susceptible subjects and chronic amoebic hepatitis	5-10	400 mg three times daily	200 mg three times daily	100 mg four times daily	100 mg three times daily
(c) Amoebic liver abscess also other forms of extra-intestinal	5	400 mg three times daily	200 mg three times daily	100 mg four times daily	100 mg three times daily

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

amoebiasis					
(d) Symptomless cyst passers	5-10	400-800 mg three times daily	200-400 mg three times daily	100-200 mg four times daily	100-200 mg three times daily
	Alternatively, doses may be expressed by body weight 35 to 50mg/kg daily in 3 divided doses for 5 to 10 days, not to exceed 2400mg/day				
Giardiasis	3 or	2000mg once daily	1000mg once daily	600-800mg once daily	500mg once daily
	5 or	400mg three times daily			
	7 - 10	500mg twice daily			
			Alternatively, as expressed in mg per kg of body weight: 15 – 40mg/kg/day divided in 2 – 3 doses.		
Acute Ulcerative Gingivitis	3	200mg three times daily	100mg three times daily	100mg twice daily	50mg three times
Acute Dental Infections	3-7	200mg three times daily			
Leg Ulcers and Pressure Sores	7	400mg three times daily			

Dosage is given in terms of metronidazole or metronidazole equivalent.

* Children and babies weighing less than 10Kg should receive proportionally smaller doses.

** Metronidazole is well tolerated by the elderly, but a pharmacokinetic study suggests cautious use of high dosage regimen in this age group.

Eradication of Helicobacter pylori in paediatric patients:

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED	
BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

As a part of combination therapy, 20mg/kg/day not to exceed 500mg twice daily for 7 – 14 days. Official guidelines should be consulted before initiating therapy.

Method of administration

For oral administration only.

4.3 Contraindications

Known hypersensitivity to Metronidazole, nitroimidazoles and/or hydroxybenzoates or any of the excipients.

4.4 Special warnings and precautions for use

Regular clinical and laboratory monitoring (especially leucocyte count) are advised if administration of Metronidazole for more than 10 days is considered to be necessary and patients should be monitored for adverse reactions such as peripheral or central neuropathy (such as paraesthesia, ataxia, dizziness, convulsive seizures).

There is the possibility that after *Trichomonas vaginalis* has been eliminated a gonococcal infection might persist.

The elimination half-life of metronidazole remains unchanged in the presence of renal failure. The dosage of metronidazole therefore needs no reduction. Such patients however, retain the metabolites of metronidazole. The clinical significance of this is not known at present.

In patients undergoing haemodialysis, metronidazole and metabolites are efficiently removed during an eight-hour period of dialysis. Metronidazole should therefore, be re-administered immediately after haemodialysis. No routine adjustment in the dosage of Metronidazole need be made in patients with renal failure undergoing intermittent peritoneal dialysis (IPD) or continuous ambulatory peritoneal dialysis (CAPD).

Metronidazole is mainly metabolised by hepatic oxidation. Substantial impairment of metronidazole clearance may occur in the presence of advanced hepatic insufficiency.

Significant cumulation may occur in patients with hepatic encephalopathy and the resulting high plasma concentrations of metronidazole may contribute to the symptoms of encephalopathy.

Metronidazole should be administered with caution to patients with hepatic encephalopathy. The daily dosage may be reduced to one third and may be administered once daily.

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

Metronidazole should be used with caution in patients with active or chronic severe peripheral and central nervous system disease due to the risk of neurological aggravation.

Metronidazole Oral Suspension contains glucose, sucrose and sorbitol. Each 5ml contains 1.1g glucose, 0.6g sucrose and 0.6g sorbitol. Patients with hereditary fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine. The glucose and sucrose content should be taken into account in patients with diabetes mellitus. It may also be harmful to teeth.

Methyl, ethyl and propyl hydroxybenzoates are contained in this product which may cause allergic reactions (possibly delayed).

Patients should be warned that metronidazole may darken urine.

Due to inadequate evidence on the mutagenicity risk in humans, the use of Metronidazole for longer treatment than usually required should be carefully considered.

Cases of severe hepatotoxicity/acute hepatic failure, including cases with a fatal outcome with very rapid onset after treatment initiation in patients with Cockayne syndrome have been reported with products containing metronidazole for systemic use. In this population, metronidazole should therefore be used after careful benefit-risk assessment and only if no alternative treatment is available. Liver function tests must be performed just prior to the start of therapy, throughout and after end of treatment until liver function is within normal ranges, or until the baseline values are reached. If the liver function tests become markedly elevated during treatment, the drug should be discontinued.

Patients with Cockayne syndrome should be advised to immediately report any symptoms of potential liver injury to their physician and stop taking metronidazole.

4.5 Interaction with other medicinal products and other forms of interaction

Patients should be advised not to take alcohol during metronidazole therapy and for at least 48 hours afterwards because of the possibility of a disulfiram-like (Antabuse effect) reaction.

Psychotic reactions have been reported in patients who were using metronidazole and disulfiram concurrently. Some potentiation of anticoagulant therapy has been reported when metronidazole has been used with the warfarin type oral anti-coagulants. Dosage of the anticoagulant may require reducing. Prothrombin time should be monitored. No interactions have been reported of the heparin type.

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

Lithium retention accompanied by evidence of possible renal damage has been reported in patients treated simultaneously with lithium and metronidazole. Lithium treatment should be tapered or withdrawn before administering metronidazole. Plasma concentration of lithium, creatinine and electrolytes should be monitored in patients under treatment with lithium while they receive metronidazole.

Patients receiving phenobarbital or phenytoin metabolise metronidazole at a much greater rate than normally, reducing the half life to approximately three hours.

Increased serum carbamazepine levels and toxicity have been seen in patients given concomitant metronidazole.

Aspartate amino transferase assays may give spuriously low values in patients taking metronidazole, depending on the method used.

Clinicians who contemplate continuous therapy for the relief of chronic conditions, for periods no longer than those recommended, are advised to consider the possible therapeutic benefit against the risk of peripheral neuropathy.

Metronidazole reduces the clearance of 5-fluorouracil and can therefore result in increased toxicity of 5-fluorouracil.

Patients receiving ciclosporin or tacrolimus with metronidazole are at risk of elevated ciclosporin / tacrolimus serum levels. Serum ciclosporin / tacrolimus and serum creatinine should be closely monitored when coadministration is necessary.

Plasma levels of busulfan may be increased by metronidazole which may lead to severe busulfan toxicity.

4.6 Pregnancy and lactation

There is inadequate evidence of the safety of metronidazole in pregnancy. Metronidazole should not therefore be given during pregnancy or during lactation unless the physician considers it essential, in these circumstances short, high dosage regimes are not recommended.

A significant amount of metronidazole is found in breast milk and breast feeding should be avoided after a large dose. This could give a bitter taste to the milk.

4.7 Effects on ability to drive and use machines

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

Patients should be warned about the potential for drowsiness, dizziness, confusion, hallucinations, convulsions or transient visual disorders, and advised not to drive or operate machinery if these symptoms occur.

4.8 Undesirable effects

The frequency of adverse events listed below is defined using the following convention:

very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data).

Frequency, type and severity of adverse reactions in children are the same as in adults.

Serious adverse reactions occur very rarely with standard recommended regimens. However, clinicians who contemplate continuous therapy for the relief of chronic conditions, for periods longer than those recommended are advised to consider the possible therapeutic benefit against the risk of peripheral neuropathy.

Blood and lymphatic system disorders:

Very rare:	agranulocytosis, neutropenia, thrombocytopenia and pancytopenia, often reversible on drug withdrawal, although fatalities have occurred.
Not known:	A moderate leucopenia has been reported in some patients but the white cell count has always returned to normal before or after treatment has been completed.

Immune system disorders:

Rare:	Anaphylaxis
Not known:	urticaria, angioedema and fever

Metabolism and nutrition disorders:

Not known:	anorexia
------------	----------

Psychiatric disorders:

Very rare:	psychotic disorders, including confusion and hallucinations
Not known:	depressed mood

Nervous system disorders:

Very rare:

- Encephalopathy (eg. confusion, fever, headache, hallucinations, paralysis, light sensitivity, disturbances in sight and movement, stiff neck) and subacute cerebellar syndrome (eg. ataxia, dysarthria, gait

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

impairment, nystagmus and tremor) have been reported very rarely which may resolve on discontinuation of the drug

- Drowsiness, dizziness, convulsions, headache, ataxia, inco-ordination of movement

Not known:

- During intensive and/or prolonged metronidazole therapy a few instances of peripheral neuropathy or transient epileptiform seizures have been reported. In most cases neuropathy disappeared after treatment was stopped or when dosage was reduced.

- Aseptic meningitis has been reported

Eye disorders:

Very rare:	transient visual disorders such as diplopia and myopia have been reported
Not known:	Optic neuropathy/neuritis has been reported

Ear and labyrinth disorders:

<i>Not known:</i>	hearing impaired/hearing loss (including sensorineural), tinnitus
-------------------	---

Gastrointestinal disorders:

Not known:	Unpleasant taste in the mouth, oral mucositis, furred tongue, nausea, vomiting, gastrointestinal disturbances such as epigastric pain and diarrhoea.
------------	--

Hepatobiliary disorders:

Very rare:

- Abnormal liver function tests, increase in liver enzymes (AST, ALT, alkaline phosphatase), cholestatic or mixed hepatitis, and hepatocellular liver injury, jaundice and pancreatitis, reversible on drug withdrawal have been reported.
- Cases of liver failure requiring liver transplant have been reported in patients treated with metronidazole in combination with other antibiotic drugs.

Skin and subcutaneous tissue disorders:

Very rare:	skin rashes, pustular eruptions, pruritus, flushing
Not known:	Erythema multiforme may occur, which may be reversed on drug withdrawal. Stevens-Johnson syndrome or toxic epidermal necrolysis, fixed drug eruption.

Musculoskeletal, connective tissue and bone disorders:

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

Very rare:	myalgia, arthralgia
<i>Renal and urinary disorders:</i>	
Very rare:	darkening of the urine (due to metronidazole metabolite)

Metronidazole Oral Suspension contains glycerol, which can cause headache, gastro-intestinal disturbance and diarrhoea.

The parahydroxybenzoates used in Metronidazole Oral Suspension may cause immediate or delayed hypersensitivity reactions.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

Single oral doses of metronidazole, up to 12g have been reported in suicide attempts and accidental overdoses. Symptoms were limited to vomiting, ataxia and slight disorientation. There is no specific antidote for metronidazole overdosage. In cases of suspected massive overdose, symptomatic and supportive treatment should be instituted.

5. Pharmacological properties

5.1 Pharmacodynamic properties

The selective action of this compound against anaerobes and anoxic and hypoxic cells is due to the mode of action. The nitro group of metronidazole acts as electron acceptor and is thus reduced to a chemically reactive drug form. This produces biochemical lesions in the cells, thus causing death. The major site of action is believed to be DNA, where it causes loss of the helical structure and inhibits synthesis.

5.2 Pharmacokinetic properties

It is readily absorbed from the gastro-intestinal tract and widely distributed in body tissues. Half life in plasma is about 8-10 hours. About 10% is bound to plasma proteins.

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

It penetrates well into body tissues and fluids, including vaginal secretions, seminal fluid, saliva and breast milk. Therapeutic concentrations are also achieved in cerebrospinal fluid.

Unchanged metronidazole and several metabolites are excreted in the urine, the liver is the main site of metabolism and the major metabolites are as a result of side chain oxidation, forming glucuronides.

5.3 Preclinical safety data

Metronidazole has been shown to be carcinogenic in the mouse and in the rat following chronic oral administration however similar studies in the hamster have given negative results. Epidemiological studies have provided no clear evidence of an increased carcinogenic risk in humans.

Metronidazole has been shown to be mutagenic in bacteria in vitro. In studies conducted in mammalian cells in vitro as well as in rodent or humans in vivo, there was inadequate evidence of a mutagenic effect of metronidazole, with some studies reporting mutagenic effects, while others studies were negative.

6. Pharmaceutical particulars

6.1 List of excipients

Sodium benzoate, Sucrose, Glycerine, Methyl paraben, Propyl Paraben, Guar Gum, Tween 80, Tetrazine yellow, Mint oil, Sodium Hydroxide

6.2 Incompatibilities

None known

6.3 Shelf life

36 months

6.4 Special precautions for storage

Store below 25°C and protect from light.

6.5 Nature and contents of container

60 ml

FIRST VADIS PHARMACEUTICAL INDUSTRIES LIMITED

BRAND NAME	VADIS METRONIDAZOLE SUSPENSION
GENERIC NAME	METRONIDAZOLE SUSPENSION

6.6 Special precautions for disposal and other handling

Keep out of the reach of children.

Shake the bottle well before use.

If a dose of under 5ml is required, the suspension should be administered using an oral dosing device.

7. Marketing authorisation holder

First Vadis Pharmaceutical,

Industries Limited

Plot IN/2 Phase 2 Extension,

Emene Industrial Layout Enugu State

Manufacturer
First Vadis Pharmaceutical
Industries Limited,
Enugu State.

8. Marketing authorisation number(s)

NAFDAC REG. NO.: 04 – 9718

9. Date of first authorisation/renewal of the authorisation

10. Date of revision of the text
