



SUMMARY OF PRODUCT CHARACTERISTICS(SmPC)

1.0 NAME OF THE MEDICINAL PRODUCT

Zuresi Diazepam 5 mg and 10 mg 1000 tablets

2.0 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each Tablet contains:

Diazepam BP 5 mg and 10 mg

Excipients Q.S.

Excipient(s) with known effect:

For the full list of excipients, see section 6.1.

3.0 PHARMACEUTICAL FORM

Uncoated Tablet

4.0 Clinical particulars

4.1 Therapeutic indications

Adults The short-term relief (2-4 weeks) only, of anxiety which is severe, disabling, or subjecting the individual to unacceptable distress, occurring alone or in association with insomnia or short-term psychosomatic, organic or psychotic illness • Cerebral palsy. • Muscle spasm. • As an adjunct to certain types of epilepsy (eg myoclonus). • Symptomatic treatment of acute alcohol withdrawal. • As oral premedication for the nervous dental patient. • For premedication before surgery Children • Control of tension and irritability in cerebral spasticity in selected cases 2) • As an adjunct to the control of muscle spasm in tetanus • Oral premedication

4.2 Posology and method of administration

Posology

As an anxiolytic, the lowest effective dose should be employed; dosage regimes should not exceed beyond 4 weeks and treatment should be gradually withdrawn. Patients who have received benzodiazepines for a long time may require an extended withdrawal period. Longterm chronic use

is not recommended. Adults: Anxiety states, obsessive-compulsive neuroses, and other psychiatric disorders: 5- 30mg daily in divided doses. Insomnia associated with anxiety: 5-15mg before retiring. Cerebral palsy: 5-60mg daily in divided doses. Upper motor neuron spasticity: 5-60mg daily in divided doses. Muscle spasm of varied aetiology, fibrositis, cervical spondylosis: 5-15mg daily in divided doses. Adjunct to the management of some types of epilepsy: 2-60 mg daily in divided doses. Alcohol withdrawal: 5-20mg, repeated, if necessary, in 2 to 4 hours. Oral premedication in dental patients: 5mg the night before, 5mg on waking and 5mg two hours before the appointment. Oral Premedication before surgery: 5mg-20mg. Children: Alternative presentations of diazepam are recommended for paediatric usage in order to obtain suitable doses of less than 5mg. Spastic children with minimal brain damage: 5-40mg daily in divided doses. Oral Premedication before surgery: 2mg-10mg Elderly and debilitated patients: Doses should be half the above recommended doses. Renal and hepatic impairment The use of diazepam in hepatic impairment may precipitate coma, therefore the dose should be reduced or an alternative drug considered. In severe renal impairment the dose should be reduced.

Method of administration

For Oral Route Administration

4.3 Contraindications

Diazepam is contra-indicated for patients with: • Known hypersensitivity to benzodiazepines and any other ingredients in diazepam tablets • Severe hepatic insufficiency (elimination half-life of diazepam may be prolonged) • Phobic or obsessional states; chronic psychosis, hyperkinesia (paradoxical reactions may occur) • Acute pulmonary insufficiency; respiratory depression, acute or chronic severe respiratory insufficiency (ventilatory failure may be exacerbated) • Myasthenia gravis (condition may be exacerbated) • Sleep apnoea (condition may be exacerbated) • Diazepam should not be used as monotherapy in patients with depression or those with anxiety and depression as suicide may be precipitated in such patients. • Pregnancy (unless there are compelling reasons). Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

4.4 Special warnings and precautions for use

The concomitant use of Zuresi diazepam with alcohol and/or CNS depressants should be avoided. Such concomitant use has the potential to increase the clinical effects of diazepam possibly including severe sedation, clinically relevant respiratory and/or cardio-vascular depression.

Duration of Treatment - The duration of treatment should be as short as possible depending on the indication. The patient must be evaluated after a period of no more than 4 weeks and then regularly thereafter in order to assess the need for continued treatment, especially if the patient is free of symptoms. In general, treatment must not last any longer than 8-12 weeks, including the tapering off process. Extension beyond these periods should not take place without re-evaluation of the situation. It may be useful to inform the patient when treatment is started that it will be of limited duration and to explain precisely how the dosage will be progressively decreased. Moreover it is important that the patient should be aware of the possibility of rebound phenomena, thereby minimizing anxiety over such symptoms should they occur while diazepam is being discontinued. There are indications that, in the case of benzodiazepines with a short duration of action, withdrawal phenomena can become manifest within the dosage interval, especially when the dosage is high.

4.5 Interaction with other Medicinal Products and other forms of interaction

Use of diazepam may lead to the development of physical and psychic dependence. The risk of dependence increases with the dose and duration of treatment, and in patients with a history of alcoholism and drug abuse or in patients with marked personality disorders. Regular monitoring in such patients is essential, routine repeat prescriptions should be avoided and treatment should be withdrawn gradually.

As sudden discontinuation of benzodiazepines may result in convulsions, particular care should be taken in patients with epilepsy, other patients who have had a history of seizures or in alcohol or drug dependants.

- Tolerance - Limits of tolerance in patients with organic cerebral changes (particularly arteriosclerosis) or cardio-respiratory insufficiency may be very wide; care must be taken in adapting the dosage with such patients.
- Some loss of efficacy to the hypnotic effects of diazepam may develop after repeated use for a few weeks.
- Alcohol should be avoided during treatment with diazepam (additive CNS depression).
- Amnesia - diazepam may induce anterograde amnesia. The condition occurs most often several hours after ingesting the product and therefore to reduce the risk patients should ensure that they will be able to have uninterrupted sleep of 7-8 hours. Anterograde amnesia may occur using therapeutic doses, the risk increases with higher doses.
- In cases of loss of bereavement, psychological adjustment may be inhibited by benzodiazepines.
- Diazepam should be used with caution in patients with a history of alcohol or drug abuse as these are patients predisposed to habituation and dependence.

- Hypo-albuminaemia may predispose patient to higher incidence of sedative side effects.
- Extreme caution should be used in prescribing diazepam to patients with personality disorders.

Paediatric population

Alternative presentations of diazepam are recommended for paediatric usage in order to obtain suitable doses of less than 5mg. Safety and effectiveness of diazepam in paediatric patients below the age of 6 months have not been established.

4.6 Interaction with other medicinal products and other forms of interaction

Alcohol Diazepam should not be used together with alcohol (CNS inhibition enhanced sedative effects: impaired ability to drive/ operate machinery). Sodium oxybate Avoid concomitant use (enhanced effects of sodium oxybate). HIV-protease inhibitors Avoid concomitant use (increased risk of prolonged sedation) – see below for zidovudine.

Take into account Pharmacodynamic interactions If diazepam is used with other centrally acting agents, careful consideration has to be given to the pharmacology of the agents employed, particularly with compounds that may potentiate or be potentiated by the action of diazepam, such as neuroleptics, anxiolytics/sedatives, hypnotics, antidepressants, anticonvulsants, sedating antihistamines, antipsychotics, anaesthetics for general anaesthesia and narcotic analgesics. Such concomitant use may increase sedative effects and cause depression of respiratory and cardiovascular functions. Concomitant use of narcotic analgesics may promote psychic dependency due to enhancement of euphorogenic effects. Anti-epileptic drugs Pharmacokinetic studies on potential interactions between diazepam and antiepileptic drugs have produced conflicting results. Both depression and elevation of drug levels, as well as no change, have been reported. Phenobarbital taken concomitantly may result in an additive CNS effect. Increased risk of sedation and respiratory depression. Phenobarbital is a known inducer of CYP3A4 and increases hepatic metabolism of diazepam. Reduced effect of diazepam. Special care should be taken in adjusting the dose in the initial stages of treatment. Side effects may be more evident with hydantoins or barbiturates. Diazepam has been reported to be displaced from protein-binding sites by sodium valproate (increased serum levels: increased risk of drowsiness). Narcotic analgesics Enhancement of the euphoria may lead to increased psychological dependence. Other drugs enhancing the sedative effect of diazepam Cisapride, lofexidine, nabilone, disulfiram and the muscle-relaxants – baclofen, Tizanidine, suxamethonium and tubocurarin. Compounds that affect hepatic enzymes (particularly cytochrome P450): • Inhibitors (eg cimetidine: isoniazid: erythromycin: omeprazole: esomeprazole) reduce clearance and may potentiate the action of benzodiazepines. Itraconazole, ketoconazole, and to a lesser extent fluconazole and voriconazole are potent inhibitors of the cytochrome P450 isoenzyme CYP3A4 and may increase plasma levels of benzodiazepines. The

effects of benzodiazepines may be increased and prolonged by concomitant use. A dose reduction of the benzodiazepine may be required.

Rifamycins (rifampicin) Rifampicin is a potent inducer of CYP3A4 and substantially increases the hepatic metabolism and clearance of diazepam. In a study with healthy subjects administered 600 mg or 1.2 g rifampicin daily for 7 days, the clearance of diazepam was increased by about fourfold. Coadministration with rifampicin gives rise to substantially decreased concentrations of diazepam. Reduced effect of diazepam. The concomitant use of rifampicin and diazepam should be avoided.

Antihypertensives, vasodilators & diuretics: Enhanced hypotensive effect with ACE inhibitors, alpha-blockers, angiotensin-II receptor antagonists, calcium channel blockers, adrenergic neurone blockers, beta-blockers, moxonidine, nitrates, hydralazine, minoxidil, sodium nitroprusside and diuretics. Enhanced sedative effect with alpha-blockers or moxonidine.

Dopaminergics Possible antagonism of the effect of levodopa.

Antacids Concurrent use may delay absorption of diazepam

Antiviral agents (atazanavir, ritonavir, delavirdine, efavirenz, indinavir, nelfinavir, saquinavir) Antiviral agents may inhibit the CYP3A4 metabolic pathway for diazepam. Increased risk of sedation and respiratory depression. Therefore, concomitant use should be avoided.

Zidovudine Increased zidovudine clearance by diazepam. Oral contraceptives Inhibition of oxidative metabolism of diazepam. Increased effects of diazepam. Co-administration of diazepam and combined oral contraceptives has been known to cause breakthrough bleeding. The mechanism of this reaction is unknown. Breakthrough bleeding, but no contraceptive failures have been reported.

Theophylline A proposed mechanism is competitive binding of theophylline to adenosine receptors in the brain. Counteraction of the pharmacodynamic effects of diazepam, e.g. reduction of sedation and psychomotor effects.

Caffeine Concurrent use may result in reduced sedative and anxiolytic effects of diazepam.

Grapefruit juice Inhibition of CYP3A4 may increase the plasma concentration of diazepam (possible increased sedation and amnesia). C_{max} increased by 1.5 times and AUC by 3.2 times. Possible increased effect of diazepam. This interaction may have little significance in healthy individuals, but it is not clear if other factors such as old age or liver cirrhosis increase the risk of adverse effects with concurrent use.

Clozapine Mechanism: Pharmacodynamic synergism. Effect: Severe hypotension, respiratory depression, unconsciousness and potentially fatal respiratory and/or cardiac arrest. Therefore, concomitant use is not recommended and should be avoided.

Pharmacokinetic interactions Diazepam is mainly metabolised to the pharmacologically active metabolites N-desmethyldiazepam, temazepam and oxazepam. The oxidative metabolism of diazepam is mediated by CYP3A4 and CYP2C19 isoenzymes. Oxazepam and temazepam are further conjugated to glucuronic acid. Inhibitors of CYP3A4 and/or CYP2C19 can give rise to increased concentrations of diazepam while enzyme inducing drugs such as rifampicin, hypericum

perforatum and certain antiepileptics can result in substantially decreased plasma concentrations of diazepam. Carbamazepine Carbamazepine is a known inducer of CYP3A4 and increases hepatic metabolism of diazepam. This can result in up to three-fold greater plasma clearance and a shorter half-life of diazepam. Reduced effect of diazepam. Phenytoin Phenytoin is a known inducer of CYP3A4 and increases hepatic metabolism of diazepam. Reduced effect of diazepam. The metabolism of phenytoin may be increased or decreased or remain unaltered by diazepam in an unpredictable way. Increased or decreased serum concentration of phenytoin. Phenytoin concentrations should be monitored more closely when diazepam is added or discontinued. Azoles (fluconazole, itraconazole, ketoconazole, voriconazole) Increased plasma concentration of benzodiazepines, due to inhibition of the CYP3A4 and/or CYP2C19 metabolic pathway. Fluconazole: Co-administration with 400 mg fluconazole on the first day and 200 mg on the second day increased the AUC of a single 5 mg oral dose of diazepam 2.5-fold and prolonged the half-life from 31 hours to 73 hours. Voriconazole: A study with healthy subjects found that 400 mg voriconazole twice daily on the first day and 200 mg twice daily on the second day increased the AUC of a single 5 mg oral dose of diazepam 2.2-fold and prolonged the half-life from 31 hours to 61 hours. Increased risk of undesired effects and toxicity of benzodiazepine. Concomitant use should be avoided or the dose of diazepam reduced. Fluvoxamine Fluvoxamine inhibits both CYP3A4 and CYP2C19 which leads to inhibition of the oxidative metabolism of diazepam. Co-administration with fluvoxamine results in an increased half-life and an approximately 190% increased plasma concentrations (AUC) of diazepam. Drowsiness, reduced psychomotor performance and memory. Preferably, benzodiazepines that are metabolised via a non-oxidative pathway should be used instead. Corticosteroids Chronic use of corticosteroids may cause increased metabolism of diazepam due to induction of cytochrome P450 isoenzyme CYP3A4, or of enzymes responsible for glucuronidation. Reduced effects of diazepam. Cimetidine Cimetidine inhibits the hepatic metabolism of diazepam, reducing its clearance and prolonging its half-life. In one study where 300 mg cimetidine was administered four times daily for 2 weeks, the combined plasma level of diazepam and its active metabolite, desmethyldiazepam, was found to be increased by 57%, but reaction times and other motor and intellectual tests remained unaffected. Increased action of diazepam and increased risk of drowsiness. Reduction of the diazepam dose may be necessary. Omeprazole Omeprazole inhibits the CYP2C19 metabolic pathway for diazepam. Omeprazole prolongs the elimination half-life of diazepam and increases the plasma concentrations (AUC) of diazepam approximately between 30% - 120%. The effect is seen in CYP2C19 extensive metabolisers but not in slow metabolisers, with a low clearance of diazepam. Increased action of diazepam. Reduction of the diazepam dose may be necessary. Esomeprazole Esomeprazole inhibits

the CYP2C19 metabolic pathway for diazepam. Co- administration with ezomeprazole results in an extended half-life and an increase in plasma concentrations (AUC) of diazepam by approximately 80%. Increased effect of diazepam. Reduction of the diazepam dose may be necessary. Isoniazid Isoniazid inhibits the CYP2C19 and CYP3A4 metabolic pathway for diazepam. Coadministration with 90 mg isoniazid twice daily for 3 days resulted in a a prolonged elimination half-life of diazepam and in a 35% increased plasma concentration (AUC) of diazepam. Increased effect of diazepam. Itraconazole Increased plasma concentration of diazepam due to inhibition of the CYP3A4 metabolic pathway. In a study with healthy subject given 200 mg itraconazole daily for 4 days increased the AUC of a single 5 mg oral dose of diazepam by about 15%, but there was no clinically significant interaction as determined by psychomotor performance tests. Possible increased effect of diazepam. Fluoxetine Fluoxetine inhibits the metabolism of diazepam via CYP2C19 and other pathways, resulting in elevated plasma concentrations and decreased clearance of diazepam. Increased effect of diazepam. Concomitant use should be monitored closely. Disulfiram Reduced metabolism of diazepam leading to prolonged half-life and increased plasma concentration of diazepam. The elimination of the N-desmethyl metabolites of diazepam is slowed down which can give rise to marked sedative effects. Increased risk of CNS inhibition such as sedation. Cisapride Accelerated absorption of diazepam. Temporary increase of the sedative effects of orally administered diazepam. Levodopa Concomitant use with diazepam resulted in reduced effects of levodopa in a small number of case reports. Ketamine Due to similar oxidative processes, diazepam competitively inhibits ketamin metabolism. Pre-medication with diazepam leads to prolonged half-life of ketamine with enhanced effect as a result. Increased sedation.

Additional information on special populations None known

4.7 Pregnancy and Lactation

The safety of diazepam in human pregnancy has not been established. It should not be used in the first and third trimesters. There may be a small increase in the risk of congenital malformation, particularly oral cleft with the use of benzodiazepines in the first trimester but a causal relationship has not been established. If the product is prescribed to a woman of childbearing potential, she should be warned to contact her physician regarding discontinuance of the product if she intends to become or suspects that she is pregnant. Pregnancy If, for compelling medical reasons, the product is administered during the late phase of pregnancy, or during labour at high doses, effects on the neonate, such as hypothermia, hypotonia ("Floppy Infant Syndrome"), irregularities in the heart rate, poor suckling and moderate respiratory depression, can be expected, due to the pharmacological action of the compound. Moreover, infants born to mothers who took

benzodiazepines chronically during the latter stages of pregnancy may have developed physical dependence and may be at some risk for developing withdrawal symptoms in the postnatal period.

Studies in animals have shown reproductive toxicity. Lactation Benzodiazepines are found in the breast milk, Reports have demonstrated milk: plasma concentration ratios to vary between 0.2 and 2.7. There is therefore a risk of accumulation in the breastfeeding child. Benzodiazepines should not be given to breast feeding mothers. Fertility Studies in animals have shown a decrease in pregnancy rate and reduced number of surviving offspring in rats at high doses. There are no human data.

4.8 Effects on ability to drive and use machines

Sedation, amnesia and impaired muscular function may adversely effect the ability to drive or use machines. If insufficient sleep occurs, the likelihood of impaired alertness may be increased (see also Interactions). Impaired function and sedation may occur the following morning and for several days after. Patients should be warned that effects on the central nervous system may persist into the day after administration even after a single dose. This medicine can impair cognitive function and can affect a patient's ability to drive safely. This class of medicine is in the list of drugs included in regulations under 5a of the Road Traffic Act 1988. When prescribing this medicine, patients should be told: • The medicine is likely to affect your ability to drive • Do not drive until you know how the medicine affects you • It is an offence to drive while under the influence of this medicine • However, you would not be committing an offence (called 'statutory defence') if: • The medicine has been prescribed to treat a medical or dental problem and you have taken it according to the instructions given by the prescriber and in the information provided with the medicine and • It was not affecting your ability to drive safely

4.9 Undesirable effects

Blood and Lymphatic System Disorders: Leukopenia. There have been rare reports of granulocytopenia, pancytopenia, agranulocytosis, or thrombocytopenia. Patients with liver disease, including hepatic echinococcosis, appear to be more at risk of bone marrow suppression. **Immune System Disorders:** Hypersensitivity reactions, including rash and urticaria. **Postmarketing Adverse Reactions:** In addition to adverse events reported from clinical trials, the following events have been identified during world-wide post- approval use of Albendazole. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to Albendazole. **Blood and Lymphatic System Disorders:** Aplastic anemia, bone marrow suppression, neutropenia. **Hepatobiliary Disorders:** Elevations of hepatic enzymes, hepatitis, acute liver failure. **Skin and Subcutaneous Tissue Disorders:** Erythema multiforme, Stevens-Johnson syndrome. **Renal and Urinary Disorders:** Acute renal failure.

4.9 Overdose

Features The symptoms of diazepam overdose are mainly an intensification of the therapeutic effects (ataxia, drowsiness, dysarthria, sedation, muscle weakness, profound sleep, hypotension, bradycardia, nystagmus) or paradoxical excitation. In most cases only observation of vital functions is required. Extreme overdosage may

lead to coma, areflexia, cardio-respiratory depression and apnoea, requiring appropriate countermeasures (ventilation, cardiovascular support). Benzodiazepine respiratory depressant effects are more serious in patients with severe chronic obstructive airways disease. Severe effects in overdose also include rhabdomyolysis and hypothermia. Management Maintain a clear airway and adequate ventilation. Consider activated charcoal (50g for an adult, 1g/kg for a child) in adults who have taken more than 100mg or children who have taken more than 1mg/kg within one hour, provided they are not too drowsy. Monitoring level of consciousness, respiratory rate, pulse oximetry and blood pressure in symptomatic patients. Consider arterial blood gas analysis in patients who have a reduced level of consciousness (GCS < 8; AVPU scale P or U) or have reduced oxygen saturations on pulse oximetry. Correct hypotension by raising the foot of the bed and by giving an appropriate fluid challenge. Where hypotension is thought mainly due to decreased systemic vascular resistance, drugs with alpha-adrenergic activity such as noradrenaline or high dose dopamine (10-30 micrograms/kg/min) may be beneficial. The dose of inotrope should be titrated against blood pressure. If severe hypotension persists despite the above measures, then central venous pressure monitoring should be considered. Supportive measures are indicated depending on the patient's clinical state. Benzodiazepines are not significantly removed from the body by dialysis. Flumazenil, a benzodiazepine antagonist, is not advised as a routine diagnostic test in patients with reduced conscious level. It may sometimes be used as an alternative to ventilation in children who are naive to benzodiazepines, or in patients with COPD to avoid the need for ventilation. It is not necessary or appropriate in cases of poisoning to fully reverse the benzodiazepine effect. Flumazenil has a short half-life (about an hour) and in this situation an infusion may therefore be required. Flumazenil is contraindicated when patients have ingested multiple medicines, especially after co- ingestion of a benzodiazepine and a tricyclic antidepressant or any other drug that causes seizures. This is because the benzodiazepine may be suppressing seizures induced by the second drug; its antagonism by flumazenil can reveal severe status epilepticus that is very difficult to control. Contraindications to the use of flumazenil include features suggestive of a tricyclic antidepressant ingestion including a wide QRS, or large pupils. Use in patients post- cardiac arrest is also contraindicated. It should be used with caution in patients with a history of seizures, head injury, or chronic benzodiazepine use. Occasionally a respirator may be required but generally few problems are encountered, although behavioural changes are likely in children. If excitation occurs, barbiturates should not be used. Effects of overdose are more severe when taken with centrally-acting drugs, especially alcohol, and in the absence of supportive measures, may prove fatal.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamics properties

Diazepam is a benzodiazepine tranquillizer with anticonvulsant, sedative, muscle relaxant and amnesic properties.

5.2 Pharmacokinetic properties

Diazepam is readily and completely absorbed from the GI tract, peak plasma concentrations occurring within about 30-90 minutes of oral administration. Diazepam crosses the bloodbrain barrier and is highly lipid soluble. Diazepam has a biphasic half-life with an initial rapid distribution phase followed by a prolonged terminal elimination phase of 1 or 2 days; its action is further prolonged by the even longer half-life of 2-5 days of its principle active metabolite, desmethyldiazepam (nordiazepam), the relative proportion of which increases in the body on long-term administration. Diazepam is extensively metabolised in the liver and, in addition to desmethyldiazepam, its active metabolites include oxazepam and temazepam. It is excreted in the urine, mainly in the form of its metabolites, either free or in conjugated form. Diazepam is very extensively bound to plasma proteins. The plasma half-life of diazepam is prolonged in neonates, in the elderly, and in patients with kidney or liver disease. In addition to crossing the blood-brain barrier, diazepam and its metabolites also cross the placental barrier and are excreted in breast milk

5.3 Preclinical Safety Data

There are no pre-clinical data of relevance to the prescriber

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Maize Starch

Lactose

Magnesium Stearate

Purified Talc

Sodium Starch glycolate

6.2 Incompatibilities

Not applicable.

6.3 Shelf-life

36 months

6.4 Keep tightly closed, in a dry place at or below 30°C.

6.5 Nature and contents of container

White colour 350cc jars and white cap containing 1000 tablets

6.6 No special requirements.

7 APPLICANT/MANUFACTURER

7.1 Name and Address

EVERGREEN MORNING STAR PHARMACEUTICALS INDUSTRY LIMITED
NDIOWERRE ORLU L.G.A
ORLU, IMO STATE

7.2 Tel number

+234 814 514 1892

7.3 Email

vcaamerican@gmail.com